Colonoscopy in Rural Communities: Can Family Physicians Perform the Procedure with Safe and Efficacious Results?

Jeffrey K. Edwards, MD, and Thomas E. Norris, MD

Background: Colonoscopy is becoming increasingly necessary for many patients in screening, diagnosing, and treating colorectal problems. Because the majority of rural doctors are family physicians, providing colonoscopy for the enlarging group of patients with valid indications in rural areas is difficult, unless rural family physicians perform the procedure. Subspecialists in academic settings have been responsible for most of the previously reported studies regarding colonoscopy. We have studied the safety and efficacy of the procedure when performed by rural family physicians.

Methods: A total of 200 sequential colonoscopies performed by family physicians in a rural setting were prospectively collected. Outcomes were measured based on current recommendations and benchmarks, including rate of reaching the cecum, time to reach the cecum, time to completion of the study, pathologic lesions found, and complications.

Results: The rate of reaching the cecum was 96.5%, and the average time to the cecum was 15.9 minutes. The average time to study completion was 34.4 minutes. The rate of neoplastic polyps and cancer found was 22.5% and 2.5%, respectively. There were no serious complications.

Conclusions: Adequately trained family physicians can provide safe and technically competent colonoscopy in a rural setting. Their results compare favorably to the currently reported comparative benchmarks from other endoscopists. (J Am Board Fam Pract 2004;17:353–8.)

Colon cancer is a preventable but potentially fatal disease. In the last 2 decades, research has established that most colon cancers arise from neoplastic polyps within the colon. If these polyps are found early and removed, colon cancer can be prevented. There is currently a growing national movement to screen patients at risk for colon cancer, with a consequential growing need for surveillance of patients with a history of polyps or colon cancer. Many physicians and patients prefer colonoscopy as the “reference standard” screening study for colon polyps and cancer as well as polyp surveillance. Unfortunately, there is a lack of qualified colonoscopists in rural areas. The majority of rural doctors are family physicians. If colonoscopies are to be offered widely in rural areas, family physicians will need to perform them.

We present 200 prospectively collected sequential colonoscopy cases performed by rural family physicians, and we provide a comparison with regard to the quality and safety of the procedure as found by a current review of the literature.^{1–23} It has been reported that a competent colonoscopist can reach the cecum more than 90% of the time,^{1,3–6,10–12,15–18,21,22} in a reasonable amount of time.^{1,5,6,8,16,18,20–22} Colonoscopy can be completed with a minimal amount of risk to the patient^{3,6,7,10–12,16,18,20,23} and with a satisfactory rate of detection of pathologic lesions.^{3,8,10,11,12,16,24}

Methods

Data were collected at the time of the procedure on 200 consecutive colonoscopies performed at St. Mary's Hospital in Cottonwood, Idaho. St. Mary's Hospital is a rural, 14-bed health care facility in north central Idaho that serves the largest county, in terms of area, in the state. This service area includes a patient population of 12,000 to 15,000 people. All colonoscopies completed by family physicians at St. Mary's Hospital over a 2-year period from December 2000 through January 2002 were included in the study. These cases included both

Submitted, revised, 6 April 2004.

From the St. Mary's Hospital, Cottonwood, ID (JKE); and the School of Medicine, University of Washington, Seattle (TEN). Address correspondence to Thomas E. Norris, MD, School of Medicine, University of Washington, Seattle, WA 98195-6340 (e-mail: tnorris@u.washington.edu).

SMH COLONOSCOPY STUDY

Patient Name: _____ (Last) _____ (First)

Date: _____ Age: _____ Sex: M F

Physician: _____

Following to be completed by Physician

INDICATORS FOR COLONOSCOPY

☐ Abdominal Pain	☐ History of Colitis
☐ Anemia	☐ History of Colon Cancer
☐ Bowel Obstruction	☐ History of Polyps
☐ Colon Cancer Screening	☐ Melena
☐ Diarrhea	☐ Rectal Mass
☐ Family History of Colon Cancer	☐ Other _____
☐ Hemocult Positive Stools	

POST PROCEDURE INFORMATION

Was cecum reached? Yes _____ No _____

Any Complications? Yes _____ No _____ (If yes- example) _____

Bowel Prep: Excellent _____ Good _____ Fair _____ Poor _____

Sedation reversal medication required: Yes _____ No _____

Following to be completed by Endoscopy Room Technician

Time to reach the cecum: _____ Time to complete procedure: _____

Number of polyps removed: _____ Number of biopsies done: _____

POST COLONOSCOPY DIAGNOSIS

☐ Diverticula
☐ Diverticulitis
☐ Polyps
☐ Cancer
☐ Stenosis
☐ Normal Colon
☐ Hemorrhoids
☐ Colitis
☐ Other _____

Figure 1. Standardized data collection form.

inpatient (urgent) and outpatient (nonurgent) study participants.

The data were collected prospectively during the study period. A standardized data collection form was completed at the time of each procedure. Figure 1 shows the standardized form used to collect colonoscopy data. The physician completed this form immediately after the colonoscopy (before pathology reports were available), and endoscopy room staff recorded procedure times. Study patients were then called 1 to 3 days after the procedure to review for any problems unless they were inpatients already.

Pathology reports were collected for all procedures in which tissue specimens were obtained. Only in situations in which the pathologist confirmed biopsy results were polyps of neoplastic origin reported (adenomatous, tubular, or villous) or the case designated as colon cancer. Pathology reports of hyperplastic polyps and other nonmalignant findings were excluded from the analysis. Microsoft Access (Microsoft Corp., Redmond, WA) was the database used for storage and analysis of the

data. Analysis of the data included only the calculation of totals (sums) and percentages, using all the cases included in the study. No statistical calculations were required or performed. Data were analyzed by category, including demographics, procedure indication, time to cecum, time to completion of colonoscopy, complications, and findings. After data collection, outcome results from this descriptive study were compared with others reported in the literature.

All procedures were completed by 1 of 4 rural family physicians. The 3 younger physicians had been trained to do colonoscopy in their residencies. A younger physician proctored the oldest physician (who had been doing flexible sigmoidoscopy) until he was proficient with colonoscopy.

The colonoscopies were all performed in a dedicated hospital endoscopy room at St. Mary's Hospital. In all cases presented in this report, Certified Registered Nurse Anesthetists provided intravenous conscious sedation. It is noted that in other settings, other trained personnel could provide this service. Intravenous sedation used included a combination of midazolam, fentanyl, and propofol. Continuous cardiorespiratory monitoring was done during each procedure. An Olympus video colonoscope, model CFQ140L, was used for all procedures.

Results

Two hundred consecutive sequential colonoscopy procedures were performed over a 2-year period at St. Mary's Hospital from December 2000 through January 2002 by 4 family physicians. Of the 200 patients, 91 were women and 109 were men. The age range of patients was from 16 to 90 years, with an average patient age of 62 years. The number of colonoscopies completed per physician varied from 23 to 108. Table 1 shows the estimated number of

Table 1. Estimated Number of Colonoscopies and the Total Years Physicians Had Been Performing Colonoscopy before Beginning the Study

Physician	Number of Cases	Years Performing Procedure
A	<50	3
B	50 to 100	8
C	100 to 200	12
D	>500	15

Table 2. Most Common Indications for 200 Colonoscopies (Some Cases Had Multiple Indications)

Indication	Number of Cases	Percentage of Cases
Rectal bleeding	52	26
Hx of polyps	49	25
FHx of colon cancer	44	22
Stools positive for FOBT	35	18

Hx, history; FHx, family history; FOBT, fecal occult blood testing.

colonoscopies and years of experience performing colonoscopy by a physician before beginning the study. The most frequent indication for colonoscopy was rectal bleeding. Some patients had multiple indications for colonoscopy, whereas others were for colon cancer screening only. Twelve patients had procedures completed as inpatients for urgent reasons (ie, acute gastrointestinal bleeding). Table 2 shows the most common indications for colonoscopy.

The success rate for reaching the cecum was found to vary among physicians from 91 to 100%. The average rate for reaching the cecum overall was 96.5% (193 of 200) as shown in Table 3. Of the cases in which the cecum was not reached, 2 of 7 (29%) were the result of an obstructing colon cancer, whereas 4 of 7 (57%) were caused by a tortuous colon, and 1 was secondary to an anastomosis stenosis.

The average time to reach the cecum per physician varied from 6.5 to 23.8 minutes and is shown in Table 4. The overall average time to reach the cecum was 15.9 minutes. The physician with the most experience had the shortest average time to the cecum, whereas the physician with the least experience had the longest. The overall average time for completion of the procedure was 34.4 minutes, including biopsies and polypectomies.

Table 3. Rate to Reach the Cecum

Physician	Number of Colonoscopies	Times Cecum Reached	Percentage
A	23	21	91
B	45	43	96
C	24	24	100
D	108	105	97
Total	200	193	96.5

Table 4. Average Time to Cecum and Procedure Completion by Physician

Physician	Average Time to Reach Cecum (minutes)	Average Time to Complete Procedure (minutes)
A	23.8	51.1
B	16.6	36.5
C	16.9	33.7
D	6.5	16.3
Average	15.9	34.4

During the study, 45 of 200 (22.5%) cases were found to have neoplastic polyps that were confirmed by pathological examination. Table 5 demonstrates the frequency of neoplastic polyps found based on the most frequent preprocedure indications. Five of 200 (2.5%) patients had confirmed colon cancer. The most common preprocedure diagnosis, for patients subsequently found to have colon cancer, was hemoccult-positive stools [3 of 35 (8.5%)]. The 2 other patients with colon cancer had a preprocedure diagnosis of anemia or rectal bleeding.

Complications were considered to be adverse events that required intervention and occurred during the procedure or after, if related to the procedure. Complications included use of reversal agents with sedation, cardiorespiratory problems with sedation, bowel perforation, hospital admission, emergency department visits, and bleeding requiring transfusion that may have been related to the procedure. A review of the patients' charts 2 years after completion of the study was completed to look for any possible delayed complications related to the procedure or missed colon cancers. There was 1 case of sedation-related bradycardia that required a single dose of atropine. One patient was admitted to the hospital for abdominal pain for

Table 5. Frequency of Neoplastic Polyps Based on Most Common Procedure Indications

Indication	Number of Cases (Total)	Percentage of Cases
Stools positive for FOBT	9 (35)	26
Hx of polyps	13 (49)	27
FHx of colon cancer	8 (44)	18
Rectal bleeding	11 (52)	21

Hx, history; FHx, family history; FOBT, fecal occult blood testing.

Table 6. Review of Reported Rates of Reaching the Cecum More Than 90% of the Time

Author	Year	n	Rate to Cecum (%)
Marshall et al ⁴	1993	418	96
Church ¹⁵	1994	2,907	93.6
Marshall and Barthel ⁵	1995	423	93.4
Hopper et al ¹²	1996	713	93
Chak et al ¹	1996	315	94.3
Pierzchajlo et al ¹¹	1997	751	91.5
Wexner et al ¹⁶	1998	2,069	96.5
Tassios et al ¹⁷	1999	430	91
Kim et al ²¹	2000	909	96.4
Wexner et al ¹⁸	2001	13,580	92
Anderson et al ²²	2001	755	91.6
Thomas-Gibson et al ¹⁰	2002	505	93
Nelson et al ⁶	2002	3,196	97
Current study	2004	200	96.5

observation overnight but required no intervention. One patient was admitted to the hospital after colonoscopy for observation for several hours because of other medical problems, the length of procedure, and the number of polyps removed. This patient required no intervention. One patient required placement of a rectal tube to relieve retained air within the colon, and symptoms resolved. There were no bowel perforations. No cases required sedation reversal medications.

Discussion

This study demonstrates that the quality of colonoscopy performed by adequately trained rural family physicians compares favorably with benchmarks reported in the literature.¹⁻⁴ Although benchmarks have been poorly defined in the past, standards are now becoming more apparent. The literature suggests that the colonoscopist should be able to reach the cecum more than 90% of the time (Table 6). They should be able to perform a colonoscopy in a reasonable amount of time (Tables 7 and 8). They are required to find and diagnose all significant pathologic lesions. Finally, they must be able to complete the procedure with minimal risk of complications and patient discomfort.

The reported success rate for reaching the cecum has, in the past, varied from 57% to more than 95%.³ Our success rate to reach the cecum was 96.5%. Our rate is similar to that of other pub-

Table 7. Review of Reported Time to Reach the Cecum

Author	Year	n	Mean Time to Cecum (minutes)
Marshall ⁵	1995	34	9*
Chak et al ¹	1996	297	10.5*
Kim et al ²¹	2000	909	6.9
Anderson et al ²²	2001	755	5.6*
Nelson et al ⁶	2002	3196	10.5
Current study	2004	200	15.9

* Median rather than mean.

lished studies shown in Table 6. The majority of studies reported in the last 10 years confirm that skilled endoscopists can reach the cecum more than 90% of the time. However, when considering endoscopists in training, the rate of successful cecal intubation is usually lower than 90%.^{1,3,5,8,14,17,19} The 4 physicians in our study all had cecal intubation rates greater than 90%, suggesting that they meet the most widely studied standard in achieving technical competence in colonoscopy.

Confirmation of reaching the cecum can sometimes be a difficult task. All colonoscopies in our study were either videotaped or photographs were taken of cecal anatomy for confirmation, but these cases were not reviewed by outside staff. One study has considered the photograph documentation of cecal landmarks as confirmation of a complete colonoscopy.²⁵ They found that it was difficult for reviewers to consistently agree on whether adequate visual documentation had been obtained. Since completion of our study, there have been no missed colon cancers reported in the study group.

The procedure time was divided into 2 primary measurements: (1) time to reach the cecum and (2) time to procedure completion. We found that our average time to the cecum was 15.9 minutes. However, there were cases in the study group in which

Table 8. Review of Reported Time to Complete Colonoscopy

Author	Year	n	Mean Time to Completion (minutes)
Wexner et al ¹⁶	1998	1,023	34.7
Wexner et al ¹⁸	2001	13,580	22.7
Nelson et al ⁶	2002	3,196	30.6
Current study	2004	200	34.4

polyps were removed before reaching the cecum; this would obviously lead to an increase in the time to reach the cecum. Further analysis was not performed on these cases.

Table 7 shows the time reported to reach the cecum in recent studies. It has been stated that a “reasonable standard” would be cecal intubation within 15 minutes or less.¹ The 4 physicians in the current study showed a large amount of variation in their average time to reach the cecum (6.5 to 23.8 minutes). This seemed to be related to the number of previous procedures completed before beginning the study. A similar variation is seen when one compares the time to procedure completion among these same physicians (16.3 to 51.1 minutes). Table 8 shows recently reported studies that measured total procedure times for colonoscopy. The reported studies also show a considerable amount of variation in reported average times (22.7 to 34.7 minutes).

Various factors affect the amount of time required to reach the cecum and complete the colonoscopy procedure (including biopsies and polypectomies). Factors affecting these outcome measurements include physician experience, patient anatomy, quality of bowel preparation, pathology encountered, and reporting differences. For example, the study with the shortest reported time to reach the cecum subtracted the time consumed with polyp removal before reaching the cecum.²² At this time, more research is required to clarify these variables comparing colonoscopists and their procedure time.

Our study group included a heterogeneous collection of subjects. Patients were included regardless of their indication for colonoscopy. The indications ranged from asymptomatic screening to surveillance follow-up for polyps or prior colon cancer. Among this diverse patient population, we found a prevalence of neoplastic polyps of 22.5% and a prevalence of colon cancer of 2.5%. These findings are consistent with what has been reported previously.²⁴

The study group included a wide variety of patients based on age, indications (both diagnostic and therapeutic) and complicating medical conditions. The group also included both an inpatient (urgent) and outpatient (nonurgent) population. Some patients were believed to be too “high risk” and were referred to a larger medical center with subspecialist management. However, the number

of patients who were not offered colonoscopy by the family physicians was not followed. Despite this, there were no serious complications in the study group. One adverse cardiopulmonary event (1 of 200, 0.5%) required intervention. There were no colon perforations. Our rate of procedure-related morbidity was consistent with morbidity reported by others.^{6,7} The reported rate of mortality as a result of colonoscopy is greater than 1 in 5000, and major morbidity is approximately 0.4%.^{6,7} The rate of colon perforation is reported to vary from 0.14 to 0.65% for diagnostic procedures and 0.15 to 3.0% for therapeutic colonoscopy (including polypectomy).²³

The primary weakness of the study is the relatively small sample size; however, we continue to collect sample data prospectively and look forward to reporting this in the future. It is possible that patient complications that occurred after the day of the procedure could have been missed if the patient presented to another hospital for evaluation and treatment. However, this is unlikely because of the rural locality of the study hospital and the distance to the next largest medical facility. In addition, this patient study group is part of a rural primary care 5-clinic system with ongoing longitudinal follow-up that uses central data management.

Conclusion

As the number of indications for the use of colonoscopy increase, rural patients face mounting geographic and distance-related obstacles to obtaining the endoscopy services that they need. This study provides an appraisal of the quality of 200 consecutive colonoscopies provided by 4 rural family physicians with various degrees of experience for a heterogeneous group of patients. Outcomes were measured and compared with other published results. We found that well-trained rural family physicians could safely provide diagnostic and therapeutic colonoscopy for their patients.

Further research regarding quality measures for colonoscopy and setting “standards” regarding compliance with these measures is needed to allow objective comparison between colonoscopists. Research in this area will advance the quality of colonoscopy provided to all our patients, both urban and rural.

We thank Cheri Holthaus and Pat Forsman, of St. Mary's Hospital, who assisted greatly with data input and management.

References

1. Chak A, Cooper GS, Blades EW, Canto M, Sivak MV Jr. Prospective assessment of colonoscopic intubation skills in trainees. *Gastrointest Endosc* 1996;44:54-7.
2. Quality and outcomes assessment in gastrointestinal endoscopy. American Society for Gastrointestinal Endoscopy. *Gastrointest Endosc* 2000;52:827-30.
3. Bond JH, Frakes JT. Who should perform colonoscopy? How much training is needed? *Gastrointest Endosc* 1999;49:657-9.
4. Marshall JB, Barthel JS. The frequency of total colonoscopy and terminal ileal intubation in the 1990s. *Gastrointest Endosc* 1993;39:518-20.
5. Marshall JB. Technical proficiency of trainees performing colonoscopy: a learning curve. *Gastrointest Endosc* 1995;42:287-91.
6. Nelson DB, McQuaid KR, Bond JH, Lieberman DA, Weiss DG, Johnston TK. Procedural success and complications of large-scale screening colonoscopy. *Gastrointest Endosc* 2002;55:307-14.
7. Dafnis G, Ekblom A, Pahlman L, Blomqvist P. Complications of diagnostic and therapeutic colonoscopy within a defined population in Sweden. *Gastrointest Endosc* 2001;54:302-9.
8. Harper M, Pope JB, Mayeaux EJ, Davis TJ, Myers A, Lirette A. Colonoscopy experience at a family practice residency: a comparison to a gastroenterology and general surgery services. *Fam Med* 1997;29:575-9.
9. Worthington DV. AAFP position paper. Colonoscopy: procedural skills. *Am Fam Physician* 2000;62:1177-82.
10. Thomas-Gibson S, Thapar C, Shah SG, Saunders BP. Colonoscopy at a combined district general hospital and specialist endoscopy unit: lessons from 505 consecutive examinations. *J R Soc Med* 2002;95:194-7.
11. Pierzchajlo RP, Ackermann RJ, Vogel RL. Colonoscopy performed by a family physician. A case series of 751 procedures. *J Fam Pract* 1997;44:473-80.
12. Hopper W, Kyker KA, Rodney WM. Colonoscopy by a family physician: a 9 year experience of 1048 procedures. *J Fam Pract* 1996;43:561-6.
13. Carr KW, Worthington JM, Rodney WM, Gentry S, Sellers A, Sizemore J. Advancing from flexible sigmoidoscopy to colonoscopy in rural family practice. *Tenn Med* 1998;91:21-6.
14. Cass OW, Freeman ML, Peine CJ, Zera RT, Onstad GR. Objective evaluation of endoscopy skills during training. *Ann Intern Med* 1993;118:40-4.
15. Church JM. Complete colonoscopy: how often? And if not, why not? *Am J Gastroenterology* 1994;89:556-60.
16. Wexner SD, Forde KA, Sellers G, Geron N, Lopes A, Weiss EG, et al. How well can surgeons perform colonoscopy? *Surg Endosc* 1998;12:1410-4.
17. Tassios PS, Lada SD, Grammenos I, Demertzis K, Raptis SA. Acquisition of competence in colonoscopy: the learning curve of trainees. *Endoscopy* 1999;31:702-6.
18. Wexner SD, Garbus JE, Singh JJ. A prospective analysis of 13,580 colonoscopies. Reevaluation of credentialing guidelines. *Surg Endosc* 2001;15:251-61.
19. Mitchell RM, McCallion K, Gardiner KR, Watson RG, Collins JS. Successful colonoscopy; completion rates and reasons for incompleteness. *Ulster Med J* 2002;71:34-7.
20. Cotton PB, Connor P, McGee D, et al. Colonoscopy: practice variation among 69 hospital based endoscopists. *Gastrointest Endosc* 2003;57:352-7.
21. Kim WH, Young JC, Park JY, Min PK, Kang JK, Park IS. Factors affecting insertion time and patient discomfort during colonoscopy. *Gastrointest Endosc* 2000;52:600-5.
22. Anderson JC, Messina CR, Cohn W, et al. Factors predictive of a difficult colonoscopy. *Gastrointest Endosc* 2001;54:558-62.
23. Putcha RV, Burdick JS. Management of iatrogenic perforation. *Gastroenterol Clin North Am* 2003;32:1289-309.
24. Rex DK. Colonoscopy: a review of its yield for cancers and adenomas by indication. *Am J Gastro* 1995;90:353-63.
25. Marshall JB, Brown DN. Photodocumentation of total colonoscopy: how successful are endoscopists? Do reviewers agree? *Gastrointest Endosc* 1996;44:243-8.