

Risk Indices Associated with the Insulin Resistance Syndrome, Cardiovascular Disease, and Possible Protection with Yoga: A Systematic Review

Kim E. Innes, MSPH, PhD, Cheryl Bourguignon, RN, PhD, and Ann Gill Taylor, MS, EdD

Objective: To conduct a systematic review of published literature regarding the effects of yoga, a promising mind-body therapy, on specific anthropometric and physiologic indices of cardiovascular disease (CVD) risk and on related clinical endpoints.

Methods: We performed a literature search using 4 computerized English and Indian scientific databases. The search was restricted to original studies (1970 to 2004) evaluating the effects of yoga on CVD or indices of CVD risk associated with the insulin resistance syndrome (IRS). Randomized controlled trials (RCTs), nonrandomized controlled trials, uncontrolled (pre and post) clinical trials, and cross-sectional (observational) studies were included if they met specific criteria. Data were extracted regarding study design, setting, population size and characteristics, intervention type and duration, comparison group or condition, outcome assessment, data analysis and presentation, follow-up, and key results, and the quality of each study was evaluated according to specific predetermined criteria.

Results: We identified 70 eligible studies, including 1 observational study, 26 uncontrolled clinical trials, 21 nonrandomized controlled clinical trials, and 22 RCTs. Together, the reported results of these studies indicate beneficial changes overall in several IRS-related indices of CVD risk, including glucose tolerance and insulin sensitivity, lipid profiles, anthropometric characteristics, blood pressure, oxidative stress, coagulation profiles, sympathetic activation, and cardiovascular function, as well as improvement in several clinical endpoints.

Conclusions: Collectively, these studies suggest that yoga may reduce many IRS-related risk factors for CVD, may improve clinical outcomes, and may aid in the management of CVD and other IRS-related conditions. However, the methodologic and other limitations characterizing most of these studies preclude drawing firm conclusions. Additional high quality RCTs are needed to confirm and further elucidate the effects of standardized yoga programs on specific indices of CVD risk and related clinical endpoints. (J Am Board Fam Pract 2005;18:491–519.)

Cardiovascular disease (CVD) is the leading cause of death and disability in the United States and other industrialized nations,^{1–4} as well as in a growing number of developing countries.^{5–7} Leading to

premature morbidity and mortality, and to preventable losses of employment, earnings, and quality of life, CVD is clearly of pressing clinical and economic significance, underscoring the need for effective primary prevention efforts that target common, modifiable risk factors. Prominent among these are the physiologic and anthropometric risk factors associated with the insulin resistance syndrome (IRS), and the neuroendocrine and psychosocial alterations that may both predispose to and result from these IRS-related abnormalities.

The IRS, also referred to as syndrome X or the metabolic syndrome, is a cluster of metabolic and hemodynamic abnormalities that both collectively and independently predict the development of atherosclerosis and CVD.^{4,8–11} Core features of the IRS are insulin resistance, glucose intolerance, atherogenic dyslipidemia, high blood pressure, and

Submitted 5 January 2005; revised 20 May 2005; accepted 31 May 2005.

From the Center for the Study of Complementary and Alternative Therapies, University of Virginia Health Systems, Charlottesville, VA

Funding: This work was made possible by grant S-T32-AT00052 from the National Center for Complementary and Alternative Medicine (NCCAM). The contents of this paper are solely the responsibility of the authors and do not necessarily represent the official views of the NCCAM, or the National Institutes of Health.

Conflict of interest: none declared.

Corresponding author: Kim E. Innes, MSPH, PhD, Center for the Study of Complementary and Alternative Therapies, The Blake Center, Suite G113, PO Box 800905, University of Virginia Health Systems, Charlottesville, VA 22908-0905 (e-mail: kei6n@virginia.edu).

visceral adiposity.^{8,10–12} Other abnormalities associated with the IRS include impaired fibrinolysis and increased coagulability,^{10,13–15} chronic inflammation,^{10–12,15} endothelial dysfunction,^{10,16,17} and oxidative stress.¹⁶ Insulin resistance (ie, resistance to insulin-stimulated glucose uptake) is thought to be the primary underlying defect and a cardinal feature linking the IRS with CVD.^{8,15,16,18–20}

Increased sympathetic activity, enhanced cardiovascular reactivity, and reduced parasympathetic tone have also been strongly implicated in the pathogenesis of IRS^{10,21–29} and in the development and progression of atherosclerosis^{27,28,30,31} and cardiovascular disease.^{21,25,28–36} In addition, recent research offers compelling evidence that chronic psychological stress and negative affective states contribute significantly to the pathogenesis and progression of insulin resistance,^{37–41} glucose intolerance,³⁸ hypertension,^{42–46} dyslipidemia,^{38,41,47} and other IRS-related conditions^{28,37,48–55} and ultimately, increase risk for CVD morbidity and mortality.^{28,32,43,44,46,52,56–60}

Not only can IRS-related conditions be exacerbated by lifestyle variables, such as smoking, lack of exercise, and poor diet, but these conditions can interact with one another in a destructive manner,⁶¹ likely accounting for their synergistic effect on CVD risk.^{9,62} Thus, a vicious cycle is initiated, which, as time goes on, becomes increasingly difficult to treat, highlighting the importance of early intervention. Given that IRS-related conditions affect a large percentage of the U.S. population,^{1,4,63} are now reaching epidemic proportions worldwide,^{1,64} and are powerful predictors of CVD morbidity and mortality,^{1,65} investigating potential cost-effective strategies for reducing IRS-related risk factors for CVD is clearly warranted.

In light of the strong influence of psychosocial factors on the development of both IRS and CVD, the role of sympathetic activation in the pathogenesis of insulin-resistant states, and the mutually exacerbating effects of these and other IRS-related risk factors, mind-body therapies may have considerable potential in the prevention and treatment of CVD. Of particular interest in this regard is yoga, an ancient mind-body discipline that has been widely used in India for the management of hypertension, diabetes, and related chronic insulin resistance conditions^{66,67–69} and may hold promise as a therapeutic intervention and health promotion measure for Western populations as well. Originat-

ing in India over 4000 years ago, the practice of yoga has been rising steadily in western industrialized countries during recent decades.^{70–72} Of the 7 major branches of Hindu yoga, Hatha (or forceful) yoga, Raja (or classical) yoga, and Mantra yoga are perhaps the best known and most widely practiced forms. Mantra yoga, emphasizing the use of specific sounds or chants to achieve mental and spiritual transformation, has been popularized in the West by Maharishi Mahesh Yogi, the founder of Transcendental Meditation (TM). Both Hatha yoga and Raja yoga emphasize specific postures (asanas), including both active and relaxation poses, as well as breath control (pranayama), concentration (dharana), and meditation (dhyana). Hatha yoga, the branch of yoga most commonly practiced in the Western industrialized world and which itself includes many different styles (eg, Iyengar, Kundalini, Ashtanga, Integral, and Bikram yoga), also incorporates mantras or chants, cleansing exercises (kriyas), and specific hand gestures (mudras).

Yoga is an economical, noninvasive practice that has become increasingly popular in the United States as a means of potentially relieving stress, enhancing health, and improving fitness.^{70–73} With no appreciable side effects and multiple collateral lifestyle benefits,^{70,72,74,75} yoga seems safe, is simple to learn, and can be practiced even by elderly, ill, or disabled individuals.⁷⁶ Requiring little in the way of equipment or professional personnel, yoga also seems easy and inexpensive to maintain, with some studies indicating excellent long-term adherence,^{77–79} even in Western populations.⁷⁹ Most importantly, there is a growing body of research suggesting that practice of yoga may reduce IRS-related risk factors for CVD, and may attenuate signs, reduce complications, and improve the prognosis of those with frank or underlying disease.^{67–72,75,78,80}

However, despite yoga's growing popularity and apparent promise as a safe and cost-effective intervention for populations with CVD or at risk for CVD, systematic reviews are lacking. In this article, we critically review the published scientific literature regarding the effects of yoga on IRS-related anthropometric and physiologic indices of CVD risk and on cardiovascular clinical outcomes, taking into account the major limitations and biases of these studies. We also briefly discuss mechanisms that may mediate the reported effects of yoga on CVD risk.

Methods

There seems to be mounting evidence that yoga may indeed reduce IRS-related indices of CVD risk and aid in the management of CVD and other IRS-related chronic disorders. To examine this evidence systematically, we first conducted a thorough search of the published medical literature. The search was restricted to English language articles published after 1970 and available in U.S. libraries, and to original studies specifically evaluating the effects of yoga (alone or in combination with other interventions) on cardiovascular disease or IRS-related risk factors for cardiovascular disease. Specifically, original studies were included in the review if they reported outcome data and evaluated the effects of yoga and yoga-based interventions on measures of insulin resistance, lipid profiles, body weight or composition, blood pressure, oxidative stress, coagulation/fibrinolytic profiles, or on markers of sympathetic activation and cardiovascular function. We also included studies assessing the influence of yoga and yoga-based programs on relevant clinical endpoints, including medication use, CVD-related events and hospitalizations, and measures of CVD progression.

Because this review is focused on the influence of yoga on physiologic and anthropometric indices of CVD risk, those studies exclusively evaluating psychosocial risk factors for CVD were excluded. Case studies, abstracts from conference proceedings, and anecdotal reports were also eliminated, as were studies with intervention group sizes of less than 4 subjects. Studies evaluating only the effects of TM programs on CVD risk profiles and clinical outcomes were likewise excluded (for a comprehensive review of research on the use of this intervention in the treatment and prevention of coronary heart disease, see King et al⁸¹). Articles were identified using MEDLINE, PubMed, and PsycINFO, 3 commercially indexed, scientific databases. Because the majority of studies investigating the effects of yoga therapy have been conducted in the Indian subcontinent and these databases offer only incomplete capture of articles published in Indian medical journals, IndMED, a bibliographic database of over 75 major Indian biomedical journals, was also searched. Search words included yoga, yogic, relaxation, mind-body, and meditation. In addition, the citation sections of all identified articles were scanned to identify relevant pa-

pers not indexed in PsycINFO, PubMed, MEDLINE, or IndMED.

Each eligible study (ie, each study identified that met our inclusion criteria) was classified into one of 5 design categories: randomized controlled trials, nonrandomized controlled trials, uncontrolled (pre and post) trials, cross-sectional (observational) studies, and studies assessing dynamic change in specific physiologic indices during 1 or 2 yoga sessions. In randomized controlled trials (RCT), a member of the investigative team randomly allocates a yoga or yoga-based program to one group (the experimental or yoga group) and no intervention, usual care, or another intervention to one or more other groups (controls). Both groups are then evaluated to determine change over time in specific indices of CVD risk. In nonrandomized controlled studies (controlled clinical trials [CCT]), a member of the investigative team allocates a yoga-based intervention to one group, and no intervention or another intervention to one or more other groups, but the allocation to a given treatment is not performed randomly. Both groups are again assessed to determine change over time in specific indices of CVD risk relative to baseline. In uncontrolled (pre and post) studies, all subjects participate in a yoga or yoga-based program, and change over time (baseline to post-intervention) in specific indices of CVD risk is evaluated. In cross-sectional (observational) studies, the investigator does not manipulate assignment of the intervention, but rather compares, at a single point in time, indices of risk between groups with varying experience in the practice of yoga. An additional study design category included studies that monitored temporary physiologic changes occurring during 1 or 2 yoga sessions; all but one of these studies⁸² included a comparison group and/or condition, although none were randomized.

We evaluated study quality using criteria based on those used in recent systematic reviews regarding the effects of another popular mind-body therapy.^{83,84} Criteria included (1) adequate sample size; (2) explicit eligibility criteria and/or adequate description of study population; (3) single, well-defined intervention; (4) appropriate control group(s) or comparison condition(s); (5) randomization of treatment allocation; (6) blinding of outcome assessment; (7) adequate accounting for confounders; (8) statistical methods well described and appropriate; (9) outcome measures well-defined and point

estimates and measures of variability presented; (10) adequate follow-up/drop-out rate reported; and (11) conclusions supported by findings.

Outcomes evaluated were categorized into several different domains of CVD risk indices and clinical outcomes. For each domain, we summarized findings from relevant studies. Because of the heterogeneity of outcomes, study designs, settings, and interventions, we did not conduct a meta-analysis.

Results and Discussion

The literature search identified a total of 70 eligible studies on physiologic/anthropometric risk factors for cardiovascular disease and/or related clinical endpoints, including 1 cross-sectional study, 8 studies (7 controlled, 1 uncontrolled) examining changes occurring during 1 or 2 yoga sessions, 25 uncontrolled clinical trials, 15 nonrandomized controlled trials (including one study that also examined changes occurring during a single yoga session), and 22 RCTs (Table 1). The majority (63%) of identified studies were published between 1990 and 2004, and most were small in size, with over 40% having study populations numbering under 25 subjects. Almost 70% of the studies were conducted in India; in contrast, less than 5% of studies identified were performed in the United States. Yoga-based interventions used in these studies ranged in duration from 2 days to 12 months, and included a single (N = 18) or multiple components of yoga practice (N = 35), and comprehensive yoga-based programs that included a special diet, lifestyle education, nonyogic exercise, stress management, or other interventions (N = 17). The distribution of studies by study population and intervention (yoga alone versus multiple interventions) is given in Table 2. The majority of the studies identified for this review were conducted in healthy young to middle-aged adults (50%) or in adults with or at risk for cardiovascular disorders (30%). In contrast, few studies have evaluated the influence of yoga on CVD risk indices in populations with other chronic disorders (N = 9), or in healthy children (N = 3) or elderly adults (N = 2) (Table 2). Table 3 lists the characteristics of eligible studies identified. The findings of these studies regarding the effects of yoga-based interventions on specific physiologic and anthropometric indices of CVD risk and relevant clinical endpoints are reviewed below.

Table 1. Characteristics of eligible studies investigating the effects of yoga and yoga-based programs on specific physiological and anthropometric indices of CVD risk and related clinical endpoints

Study Design	Total		Sample Size						Location				Year Published				
	N	%	<25	25-40	41-60	>60	India	US	Canada	Eurc	Australia	Japan	Thailand	2000	1990-9	1980-9	1970-9
Cross-sectional studies	1	1.4				1	1							1			
Studies of changes during 1 to 2 yoga sessions	1	1.4	1									1		1			
Not including comparison group or condition	7	10.1	4	3			4			2	1			3	4		
Including comparison group or condition																	
Uncontrolled clinical trials (>2 yoga sessions) (Intervention 2 full days to 6 months)	25	36.2	15	2	5	3	21	1	1	2				7	7	6	5
Non-randomized controlled trials (>2 yoga sessions)* (Intervention 3 weeks to 12 months)	15	21.7	7	2	3	3	11	1		3				3	3	4	5
Randomized controlled trials (>2 yoga sessions) (Intervention 8 days to 12 months)	22	31.9	3	10	4	5	11	1		8	1	1	1	8	8	4	2
Total number of studies evaluated	70		29	17	12	12	47	3	1	15	1	2	1	22	22	14	12
% Total			41.4	24.3	17.1	17.1	67.1	4.3	1.4	21.4	1.4	2.9	1.4	31.4	31.4	20.0	17.1

*Including one study that also evaluated change during 1 to 2 yoga sessions¹³⁸; grand totals are adjusted to count this study only once.

Table 2. Studies investigating the effects of yoga and yoga-based programs on specific physiologic and anthropometric indices of CVD risk and related clinical endpoints, by study population

Study Population	Study Design								Total		
	Cross-Sectional Studies		Changes over 1 to 2 Yoga Sessions		Uncontrolled Trials		Non-randomized Controlled Trials			Randomized Controlled Trials	
	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions		Yoga Alone	Multiple Interventions
Healthy young and middle-aged adults	1		8		11	3	7	1	5		35*
Healthy elderly adults									1	1	2
Healthy children and adolescents									3		3
Adults with hypertension/CVD or at risk for CVD					2	2	6	1	4	6	21
Adults with diabetes mellitus					3	1			1		5
Adults with cancer						1					1
Adults with serious mental illness					1						1
Children and adults with asthma						1					1
Children and adults with epilepsy									1		1
Totals	1		8		17	8	13	2	15	7	70
Total yoga alone or with biofeedback	53*										
Total multiple interventions	17										

*Including one study that also evaluated change during 1 to 2 yoga sessions¹³⁸; grand totals are adjusted to count this study only once.

Table 3. Characteristics and relevant endpoints of eligible studies. Under endpoint categories, a plus indicates a beneficial change in at least one marker; a negative sign indicates no change in any of the markers in a specific category

Reference, Location	Sample Size (Yoga, Controls)	Duration	Comparison Group/Condition	Intervention	Endpoints Examined*						
					IR indices	Lipids	Body Size/Comp	Blood Pressure	Coagulation	Oxidative Stress	Symp/PS Markers
Cross-sectional Studies											
Healthy adults											
India ⁹⁶	105 (35, 35, 35)	NA	Raja M <5 years; no M experience	Raja M experience ≥5 years		X+		X+			X+
Changes during 1–2 yoga sessions											
Healthy adults											
Germany ¹¹⁰	25 (12, 13)	1 session	Relaxed reading (controls)	Hatha yoga (experienced practitioners)				X–			X–
Australia ¹⁴²	23 (12, 11)	2 sessions	Reading	15 km run (elite runners) or M (experienced meditators)							X+
India ¹⁴⁵	7	1 session	Non-targeted thinking	Yoga M (silent chanting)							X+
India ¹⁴⁴	27	1 session	2 non-yoga resting postures	Shavasana							X+
Japan ⁸²	8	1 session	Ss own controls	Yoga (rest, AS, PR, M)							X+
Italy ³⁶	23	1 session	Talking, free breathing	Yoga mantras, slow controlled breathing							X+
India ¹⁴³	35	2 sessions	Supine rest	Yoga-based guided relaxation/M							X+
India ¹³⁸	24 (10, 14)	1 session	Naïve controls during rest	SKY yoga by trained subjects						X+	
Uncontrolled Clinical Trials											
Healthy Adults											
India ⁸⁹	12	6 months	Ss own controls	Hatha yoga course							X–
India ¹⁴⁷	20	10 full d	Ss own controls 2 camps compared	Vispanna Yoga; strict diet	X+	X+	X+				X+
India ⁹⁰	10	4 months	Ss own controls	Yoga (AS, PR, M)	X+	X+		X+			X+
India ⁹⁴	73	3 months	Ss own controls	Yoga (PR, AS, SH)	X–	X–					
India ¹¹⁶	17	3 months	Ss own controls	Yoga (AS, PR)				X+			X+

India ¹³⁶	7	4 months	Ss own controls	Yoga (AS, PR, M)			X+	
India ¹¹⁷	8	2 months	Ss own controls	SH + autogenic phrases with biofeedback			X+	X+
India ¹³⁹	75	6 weeks	Ss own controls	Yoga PR				X+
India ¹⁰³	8	30 days	Ss own controls	Santhi Kriya yoga			X-	X+
India ¹⁰⁴	40	3 months	Ss own controls	Residential program: yoga (AS, PR, M) + education			X+	X+
India ¹⁰⁵	6	4 weeks	Ss own controls	Yoga (PR, AS)				X+
India ¹¹¹	8	2 weeks	Ss own controls	Sarvangasana			X-	X+
India ¹⁴⁰	26	2 full days	Ss own controls	Yoga + stress management				X+
U.S. ¹⁰⁸	10	8 weeks	Ss own controls	Hatha yoga classes (PR, AS, SH)			X-	
Adults with hypertension/CAD								
UK ⁹⁷	14	6 weeks	Ss own controls	Relaxation, M w/biofeedback			X+	X+
India ¹¹⁸	44	30 days	Ss own controls	Yoga + diet			X+	
India ¹¹⁹	25	6 months	Ss own controls	Shavasana yoga			X+	X+
India ⁶⁸	20	78 sessions	Ss own controls	Yoga (AS, PR, M, SH) + education			X+	X+
Adults with uncomplicated DM								
India ⁹¹	52	6+ months	Ss own controls	Yoga (AS, SH)			X+	X+
India ⁹²	149	40 days	Ss own controls	Residential yoga + diet			X+	X+
India ⁸⁵	19	40 days	Ss own controls	Yoga (AS, PR, M)				
India ⁸⁶	24	40 days	Ss own controls	Supervised yoga AS, PR			X+	X+
Adults with SMI								
India ¹⁵⁷	37	3 months	Ss own controls	SKY yoga				
Canada ¹⁵⁸	69	8 weeks	Ss own controls	MBSR program (relaxation, yoga, M, daily practice)				X- X-/+
Children & adults with asthma								
India ¹⁵¹	9	7 days	Ss own controls	Residential yoga (AS, PR) + diet				X+
Non-randomized controlled trials								
Healthy adults								
India ¹¹²	28 (14, 14)	6 months	Non-yogic exercise	Yoga training (AS, PR)			X-	X+

Table 3. Continued

Reference, Location	Sample Size (Yoga, Controls)	Duration	Comparison Group/Condition	Intervention	Endpoints Examined*							
					IR indices	Lipids	Body Size/Comp	Blood Pressure	Coagulation	Oxidative Stress	Symp/PS Markers	Clin. End Points
India ⁹³	10 (4, 4, 2)	6 months	Exercise; Ss own controls	2 yoga AS routines	X+		X+				X+	
India ¹⁵³	20 (10, 10)	2.5 months	Similar sedentary controls	Yoga PR, AS							X+	
India ⁸⁸	180 (45, 45, 45, 45)	12 weeks	Untrained controls	PR: Ujjayi, Bhastrika, or combination	X+	X+						
Germany ⁹⁸	154 (118, 36) [§]	3 months	Non-participants matched for analyses	Residential program: Kriya yoga + diet		X+	X+	X+			X+	
India ¹³⁷	60 (30, 30)	10 weeks	Sedentary controls	Yogic PR training						X+		
India ¹³⁸	24 (10, 14)	5 months	No yoga practice	SKY Yoga						X+		
India ¹¹³	16 (8, 8)	1 (Ph I) to 2 years (Ph II)	Exercise only (N = 6, Phase I; N = 8, Phase II)	Yoga PR + exercise (N = 6, Phase I; N = 8, Phase II)				X−			X−	
Adults with hypertension/CAD												
UK ¹²¹	40 (20, 20)	3 months	Usual care; also meet but rest on couch	Yoga relaxation/M with biofeedback				X+				X+
UK ¹²²	47 (27, 20)	9 weeks	Usual care; also meet but rest on couch	Training in yoga relaxation & M + home practice 2×/wk				X+				X+
USA ¹²⁴	19 (14, 5)	6 months	Usual care	Buddhist M, PR, and relaxation training/practice				X+			X+	
India ^{123†}	60 (20, 10, 20, 10)	4 to 6 weeks	Group II: 2 weeks SH, 2 weeks drugs, 2 weeks both (2 weeks between each)	I, IV: 4 weeks SH III: 4 weeks SH + meds				X+			X+	
India ¹⁰⁹	14 (7, 7)	6 months (17 sessions)	Usual care, met 1×/week	Yoga, PR, relaxation (Phase I) + thermal biofeedback (Phase II)			X−	X+				X+
India ¹²⁰	20 (10, 10)	3 weeks	Postural tilt table	Postural yoga asanas				X+			X+	
India ⁸⁰	113 (71, 42)++	12 months, (21 sessions)	Usual care: Meds, diet, moderate exercise	Yoga + diet, walking, stress management, meds	X+	X+	X+					X+

Randomized controlled trials						
Healthy adults						
India ¹²⁵	18 (6, 6, 6)	12 weeks	20 minutes rest (controls)	20 min. TM, SH		X+
India ¹⁰⁶	40 (20, 20)	12 months	No yoga training	Yoga (AS, PR, PR, mudras, kriyas)	X+	
Japan ¹²⁶	90 (45, 45)	3 weekly trainings	Jacobson progressive muscle relaxation	Hatha yoga classes (PR, AS, relaxation)	X+	X+
UK ¹¹⁴	26 (12, 14)	6 weeks	Aerobic training (cycling)	Yoga training (AS, PR, relaxation)	X-	X+
India ¹²⁷	54 (28, 26)	10 months	No yoga until month 6; yoga month 6-10	Hatha yoga AS, PR, KR; mo. 1-10	X+	X+
USA ⁸⁷	46 (20, 9, 14)	12 months	Usual care; diet, exercise, educ, supplements	Yoga M, AS + walking, diet	X+	X+
India ¹²⁸	30 (15, 15)	3 months	Flexibility, aerobic exercise	Hatha yoga (AS, PR), Omkar meditation	X+	X-
Adults with Hypertension or CVD or at risk for CVD						
UK ¹⁵²	32 (16, 16)	6 weeks	Usual care, anti-HT drugs	Yoga relaxation, M training with BF+		X+
UK ¹³⁰	34 (17, 17)	6 weeks	Placebo therapy (rest supine)	Yoga relaxation, PR + biofeedback	X+	
UK ¹⁰²	192 (99, 93)	8 weeks	Usual care + lifestyle education	PR, relaxation, M stress management + lifestyle education	X+	X+
UK ⁷⁹	161 (86, 75)	8 weeks	Usual care + lifestyle education (0, 8 month, 4 year followup)	PR, M, relaxation + stress management, education	X+/-	X+
UK ¹²⁹	99 (49, 54)	8 weeks	Same medication/placebo schedule; no training	Relaxation, M with biofeedback + education, home practice	X+	X+
Holland ¹⁰¹	55 (18, 17)	8 weeks, + Cont. 1 year	Sit, relax 2x/days + support, non-specific counseling	Yoga PR, AS, PMR, stress management	X-	X-
India ⁹⁹	93 (52, 18)	Camp + 14 weeks practice	Standard care, diet control, exercise	4 d yoga camp + diet; yoga practice (AS, PR, SH, M)	X+	X+

Table 3. Continued

Reference, Location	Sample Size (Yoga, Controls)	Duration	Comparison Group/Condition	Intervention	Endpoints Examined*							
					IR indices	Lipids	Body Size/ Comp	Blood Pressure	Coagulation	Oxidative Stress	Symp/PS Markers	Clin. End Points
India ⁷⁸	42 (21, 21)	4 days training + 1 year	Usual care, regular evaluations	Yoga: (PR, AS, K, M) + diet, exercise, regular evaluations		X+	X+					X+
India ¹⁰⁷	33 (11, 11, 11)	11 weeks	Hypertension medication; no tx	Yoga (M, PR, AS) training/practice			X+	X+			X+	
Thailand ¹⁰⁰	44 (22, 22)	4 months	Usual care, including drug therapy	Yoga AS, SH + imagery education, stress management, support groups		X+				X+		
India ¹³¹	40 (10, 10, 10, 10)	8 days	Conversation regarding neutral topics	SH, Broota method, or Jacobsen PMR				X+				
Adults with Type II DM UK ⁹⁵	21 (11, 10)	12 weeks	Usual care	Yoga classes (PR, AS, SH)	X+							X+
Adults/Children with epilepsy India ¹⁴⁶	32 (10, 10, 12)	6 months	Sham yoga exercise, usual care	Sahaja yoga M						X+	X+	
Healthy Children India ¹⁴¹	28 (14, 14)	6 months	Games (jog, races, rapid stretches+)	Yoga AS							X+	
India ¹¹⁵	24 (12, 12)	3 months	No yoga practice/ training	PR exercises				X-			X+	

*+, positive (beneficial) findings for 1 or more endpoints within a given category; -, no beneficial effects demonstrated.

[†]Including 10 normotensive adults (Group I).

[‡]Including a subset of patients with diabetes mellitus.

[§]36 to 72 analyzed in matched analyses.

Abbreviations and explanation of terms: AS, yoga asanas or postures; BF, biofeedback; BP, blood pressure; Clin, clinical; comp, composition; DM, diabetes mellitus; IR, insulin resistance (markers of); KR, kriyas or cleansing exercises; M, meditation; MBSR, mindfulness-based stress-reduction program; PMR, progressive muscle relaxation; PR, pranayama or yogic breathing exercises; SH, Shavasana or corpse pose, a traditional yoga relaxation pose; Symp/PS, markers of sympathetic/parasympathetic activation and cardiovascular function, including heart and respiratory rate, neuroendocrine markers, markers of cardiovascular reactivity to and recovery from stress, and indices of heart rate variability and baroreflex sensitivity; Santhi Kriya yoga: 3 stages, totaling 50 minutes: (1) 10 minutes PR while in standing and sitting postures; (2) 10 min chanting, meditation, and breathing exercises while in shavasana (corpse pose); and (3) 30 min of relaxation and concentrative meditation (dahara vidya); SKY, Sudharshan kriya yoga, developed by Pundit Ravishankar from Bangalore, India, uses specific yoga breathing exercises (Ujjayii, Bhastrika, cyclical breathing); sessions typically close with Yoga Nidra (tranquil state) in a supine posture.

Observed Effects of Yoga on Core Indices of the IRS

Since the 1970s, at least 51 eligible, published studies from 7 countries have investigated the potential influence of yoga and yoga-based programs on one or more core indices of the IRS, including measures of insulin resistance, lipid profiles, body weight and composition, and blood pressure. Our literature search identified 13 eligible studies examining the influence of yoga-based interventions on markers of insulin resistance (including 8 uncontrolled clinical trials, 2 nonrandomized controlled trials, and 2 RCTs), 14 studies evaluating the effects of yoga on lipid profiles (including 1 cross-sectional study, 5 uncontrolled clinical trials, 3 nonrandomized controlled trials, and 5 RCTs), and 18 studies assessing the influence of yoga on body weight or composition (including 9 uncontrolled trials, 3 nonrandomized controlled trials, and 6 RCTs). In addition, we identified 37 eligible studies evaluating the effects of yoga on blood pressure, including 1 cross-sectional study, 1 study examining changes during a single yoga session, 11 uncontrolled studies, 12 nonrandomized controlled studies, and 12 RCTs. Findings of these studies are summarized in Tables 4 and 6 and are discussed briefly below.

Markers of Insulin Resistance

Of the 13 studies evaluating the effects of yoga on markers of insulin resistance, most documented significant, postintervention improvement in one or more indices following the practice of yoga either alone or in combination with other therapies (Table 4). Interventions ranged in length from 40 days^{85,86} to 12 months,^{80,87} and all but one⁸⁸ incorporated yoga asanas or postures. Seven of the 8 uncontrolled studies of healthy young adults,^{89,90} patients with type II diabetes mellitus (DM II)^{85,86,91,92} or hypertension⁶⁸ reported significant improvement postintervention in indices of insulin resistance relative to baseline values. Documented changes included reductions in fasting^{68,80,85,86,88–90,92,93} and postprandial glucose,^{85,86,89,91,92} and in fasting glycohemoglobin;^{85–87} in the one uncontrolled study yielding negative results, a trend toward reduced glucose was observed in male subjects despite low baseline levels⁹⁴ (Table 4). Likewise, nonrandomized controlled studies of healthy young adults⁸⁸ and adults with coronary artery disease (CAD)⁸⁰ reported reductions in fasting glucose among subjects receiving a yoga-based intervention versus controls re-

ceiving enhanced usual care⁸⁰ or no active intervention⁸⁸; descriptive findings of an additional small controlled study of healthy young men suggested similar improvement following a prescribed yoga program.⁹³ In a small RCT of diabetic British adults, Monro et al demonstrated a decline in both fasting glucose and glycohemoglobin among participants attending structured yoga classes relative to usual care controls.⁹⁵ Similarly, an RCT of American seniors demonstrated a substantial (19%) reduction in fasting insulin among high-risk subjects completing a comprehensive integrative medicine program including yoga meditation and asanas, although small sample sizes precluded statistical comparison with controls.⁸⁷ Overall, yoga practice was associated with a 5.4 to 33.4% reduction in fasting glucose, 24.5 to 27.0% reductions in postprandial glucose, and 13.3 to 27.3% reduction in glycohemoglobin, with the percentages varying by study population and design (Table 5).

Although the evidence for a beneficial effect of yoga on indices of insulin resistance is relatively consistent across studies, most studies (62%) had no comparison groups^{68,85,86,89–92,94} and many had small sample sizes^{68,85,86,89,90,93,95} or reduced power due to stratified analyses.^{87,92,94} Others lacked detailed information on eligibility criteria or study population characteristics,^{80,85,86,88,89,93} and blinding of the outcome assessment was reported in only one study.⁸⁷ The study sample selection process was unclear in several studies,^{86,89,90,93} and in others, potential selection bias, differential loss to follow-up⁸⁰ and uncontrolled confounding factors such as demographic, anthropometric, and lifestyle characteristics^{85,91–95} may have been important. The intervention was not well described⁸⁸ or varied in duration⁹¹ in some studies; others included multiple interventions,^{68,80,87,92} rendering detection of the effect of a specific component difficult. Analytic methods were not well described in most studies^{68,85,86,88,90–95}; no statistical analysis was presented in some,^{68,89,91,93} and many had methodological problems, including lack of adequate point estimates for key outcome measures,^{68,89,91,93} failure to adjust for multiple comparisons,^{68,87,88,90,91,94} confusing data presentation,⁶⁸ and other potential analytic problems.^{87,88} Among controlled studies, additional limitations included lack of randomization,^{80,88,93} failure to describe treatment allocation methods,⁸⁰ and or absence of intergroup comparisons.⁹³

Table 4. Summarized findings of studies (1970–2004) investigating the effects of yoga and yoga-based programs on insulin resistance, lipid profiles, and body size and composition, stratified by study design and intervention (yoga, including yoga-based meditation, alone vs. in combination with other interventions). Studies demonstrating beneficial effects are in bold type.

Findings, by Endpoint	Study Design						
	Cross- Sectional (Yoga Alone)	Uncontrolled Clinical Trials (>2 sessions)		Non-Randomized Controlled Clinical Trials (>2 Sessions)		Randomized Clinical Trials (>2 sessions)	
		Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions
Markers of insulin resistance							
Fasting glucose							
Reduced		[85, 86, 89, 90]	[68, 92]	[88, 93]	[80]	[95]	
No change		[94]*					[87]
Post prandial glucose							
Reduced		[85, 86, 89, 91]	[92]				
No change							[87]
Reduced fasting insulin							[87](NS)
Reduced fasting glycated hemoglobin		[85, 86]				[95]	
Lipid profiles							
Total Cholesterol							
Reduced	[96]	[94](NS)	[89, 90, 97] [†]	[68]	[88]	[80, 98]	[78, 79, 99]
Increased							[100]
Unchanged							[79] [‡] [101]
Triglycerides							
Reduced	[96]						[78, 99, 100]
Increased		[94]					
Increased high density lipoproteins			[68]				
Low density lipoprotein (LDL)							
Reduced						[80, 98]	[78, 99]
Increased							[100]
Anthropometric markers							
Reduced BMI			[92]		[98]		[100]
Reduced		[89, 91, 103]	[104]	[93]	[80]	[106, 107]	[78, 99]
Unchanged		[97] [†] [108]		[109]			[101]
Body composition							
Reduced waist, hip circumference						[106]	
Reduced skin fold thickness [§]						[106]	
↓ % body fat, ↑ % lean mass		[105]					
No change (% body fat, fat mass)		[108]					

*Sexes analyzed separately; trend toward reduced fasting glucose in men ($N = 53$, $0.5 < P < .1$), despite low baseline fasting glucose levels ($\bar{x} = 64.4 \pm 6.2$).

[†]With biofeedback.

[‡]In individuals at risk for CVD, cholesterol was lower at 0 and 8 months post-intervention, but not at 4 year follow-up

[§]Triceps, subscapular, thigh, calf, etc, skin folds.¹⁰⁶

Lipid Profiles

Of the 14 identified studies that examined the potential effects of yoga on blood lipid concentrations, most offered evidence suggesting that the practice of yoga and yoga-based programs may improve lipid profiles. In a cross-sectional study of healthy middle-aged men with similar lifestyle characteristics, Vyas and colleagues found those with both short-term and long-term experience in

Raja yoga meditation had reduced cholesterol and triglyceride levels relative to those with no yoga experience.⁹⁶ Similarly, 4 of 5 uncontrolled studies^{68,89,90,97} and all 3 of the controlled, nonrandomized studies^{80,88,98} identified demonstrated significant positive changes in blood lipid levels following yoga-based interventions that ranged from 6 weeks⁹⁷ to 12 months⁸⁰ in duration. These studies included investigations of both healthy adults and

Table 5. Observed percent change with yoga in selected physiologic and anthropometric indices of CVD risk*

Findings, by Endpoint and Population	Uncontrolled Clinical Trials (>2 sessions)	Non-Randomized Controlled Clinical Trials (>2 sessions)	Randomized Clinical Trials (>2 sessions)
Markers of insulin resistance			
Reduced fasting glucose			
Healthy Adults	6.8% [90]	6.4–28.0% [88, 93]	
Adults with hypertension/CVD	5.4–6.4% [68]		
Adults with type II diabetes (DM II)	25.1–33.4% [85, 86, 92]		6.9% [95]
Reduced postprandial glucose			
Adults with type II diabetes (DM II)	24.5–27.0% [85, 86, 91, 92]		
Reduced fasting Insulin			
Adults at risk for CVD			19.2% [87]
Reduced fasting glycated hemoglobin			
Adults with type II diabetes (DM II)	13.3–27.3% [85, 86]		15.5% [95]
Lipid profiles			
Reduced total cholesterol			
Healthy adults	9.3% [90]	8.0–8.2% [88] [98]	
Adults with hypertension/CVD+ or DM II	5.8–10.1% [68, 97]	25.2% [80]	14.8–21% [78, 99]
Adults with other chronic disorders			
Reduced triglycerides			
Adults with hypertension/CVD+ or DM II			22.0–28.5% [78, 99, 100]
Reduced low density lipoprotein (LDL)			
Healthy adults		12.8% [98]	
Adults with hypertension/CVD+ or DM II		26.0% [80]	19.1–23.3% [78, 99]
Anthropometric markers			
Reduced body weight			
Healthy adults	1.5–9.1% [103, 104]		10.1% [106]
Adults with hypertension/CVD+ or DM II			5.1–13.6% [78, 99, 107]
Blood pressure			
Reduced diastolic BP			
Healthy adults	7.7–8.8% [90, 104, 116]	7.3–15.9% [93 98]	4.9–12.4% [125, 127, 128]
Adults with or at risk for CVD	8.5–16.7% [68, 97, 117–119]	12.7–17.5% [80, 120–123]	8.2–24.2% [79, 87, 107, 129–131]
Reduced systolic BP			
Healthy adults	2.6–7.3% [90, 104, 116]	18% [98]	6.8–9.3% [125, 127, 128]
Adults with or at risk for CVD	7.1–17.0% [68, 97, 117–119]	11.1–20.8% [80, 120–123]	4.4–21.3% [79, 87, 107, 129–131]
Reduced baseline respiratory rate			
Healthy adults	3.8–14.4% [103, 104, 116, 139, 140]		
Healthy children/adolescents			5.3% [141]
Adults with hypertension		60.3% [123]	
Reduced baseline heart rate			
Healthy adults	4.3–38.8% [90, 104, 111, 116]	4.3–8.2% [93, 98]	5.0–11.6% [114, 115, 125–127]
Healthy children/adolescents			17.0–30.2% [115, 141]
Adults with hypertension/CVD		9.5–23.5% [120, 123]	8.4% [107]
Adults with other chronic disorders	14.9% [151]		

*Including studies reporting improvement in these indices; studies not reporting point estimates or presenting sufficient data to allow reliable calculation of point estimates were excluded; weighted averages were calculated for studies presenting only stratified data (e.g., by sex).

CVD, cardiovascular disease.

patients with hypertension or CAD (Table 3). Observed changes included reductions in cholesterol and low density lipoprotein (LDL),^{68,80,88–90,97,98}

and increases in high density lipoprotein (HDL) levels⁶⁸ relative to baseline levels^{68,89,90,97} and/or control values.^{80,88,98} In agreement with these find-

ings, several RCTs investigating the effects of yoga in combination with diet,^{78,80,99} education,⁷⁹ stress management,^{79,80,100} and other therapies^{78,80,100} have likewise demonstrated significant improvement in lipid profiles relative to controls receiving enhanced usual care, exercise, and/or dietary interventions (Tables 3 and 4). Of the 5 RCTs identified, all adults with hypertension,¹⁰¹ CVD,^{78,100} or risk factors for CVD,^{79,99} 3 documented improvement in all indices of dyslipidemia examined,^{78,79,99} and one reported significantly greater reductions in triglycerides, but not in cholesterol or LDL.¹⁰⁰ In a follow-up of an earlier British study,¹⁰² Patel and colleagues reported maintenance of lipid profile changes at 8 months postintervention, but no differences between the yoga relaxation/meditation and the control groups at 4 years.⁷⁹ Only one RCT, a study of 35 hypertensive Dutch adults (18 intervention group, 17 controls), reported no differences between the intervention and control groups.¹⁰¹ Of those studies demonstrating positive effects, yoga practice was associated with a 5.8% to 25.2% decrease in total cholesterol, 22.0% to 28.5% reduction in triglycerides, and a 12.8% to 26.0% reduction in LDL, with the observed magnitude of the effects differing by study population and design (Table 5).

Again, the vast majority of both uncontrolled and controlled studies suggest that the practice of yoga may have a positive influence on lipid profiles. However, many of the studies evaluated have methodological or other limitations that hinder interpretation of findings, including absence of comparison groups or conditions,^{68,89,90,94,97} low power because of small sample sizes^{68,89,90,97} or other factors,^{94,101} lack of evidence for blinded outcome assessment,^{68,79,80,88–90,94,96–101} possible uncontrolled confounding factors,^{94,96–101} potential selection bias,^{96,98,100} loss to follow-up,⁸⁰ lack of randomization^{88,98} or information on the randomization process,^{78,79,99,100} unclear statistical analyses,^{68,79,88,90,94,97,98} and multiple^{68,78–80,99–101} or poorly defined interventions.^{88,97} Other methodological issues include lack of adequate point estimates,^{68,88,89,98} intergroup comparisons,¹⁰⁰ or adjustment for multiple comparisons,^{68,88,90,94,98,100} possible floor effects,⁹⁴ and inadequate description of study populations^{80,88,89,99,101} and of sampling or treatment allocation process.^{80,89,90}

Body Weight and Composition

Between 1970 and 2004, 18 clinical trials in 6 countries examined the effects of yoga on anthropometric indices of CVD risk (Table 4). Six of the 8 uncontrolled clinical trials and 3 of the 4 nonrandomized controlled studies identified reported improvement in body weight^{80,89,91–93,98,103,104} or composition¹⁰⁵ after yoga-based interventions ranging from 4 weeks^{103,105} to 12 months.⁸⁰ Likewise, 5 of 6 randomized controlled trials, including 4 conducted in India^{78,99,106,107} and one completed in Thailand,¹⁰⁰ demonstrated improvement in body weight and/or composition relative to usual care,^{78,100,107} diet and exercise,⁹⁹ and no intervention controls.¹⁰⁶ Studies reporting improvement in anthropometric characteristics included investigations of healthy populations,^{89,93,98,103–106} as well as those with hypertension¹⁰⁷ and/or other CVD risk factors,⁹⁹ CAD,^{78,80,100} or diabetes.^{91,92} Of those studies reporting improvement in anthropometric indices, yoga practice was associated with a 1.5% to 13.6% reduction in body weight (Table 5). All studies reporting a positive change in these indices used interventions incorporating active yoga asanas, whereas only one¹⁰⁸ of those showing no effect, an uncontrolled 8-week study of 10 U.S. university students, included active postures as a central component in the intervention.

Limitations of these studies include lack of control groups,^{89,91,92,97,103–105,108} reduced power because of a low number of study subjects^{89,93,97,103,105,108,109} or other factors,^{92,101} possible selection bias^{98,105,108} or influence of uncontrolled confounding^{91–93,97–101,103,107,109} lack of appropriate statistical analyses or presentation,^{89,91,93,98} inadequate description of analytic methods,^{91–93,97,98,105,106} study population,* treatment allocation,⁸⁰ or sample selection process,^{89,93,106} loss to follow-up,^{80,105,108} multiple comparisons,[†] failure to compare control and intervention groups,^{93,100,109} lack of randomization^{93,98,109} or description of randomization methods,^{78,99,100,106,107} and multiple interventions.^{78,80,92,99–101,104} In addition, only one study⁷⁸ reported blinded outcome assessment.

*See References 80, 89, 91, 93, 99, 101, 104, 106, 107, 109

†See References 91, 98, 100, 104–106, 108, 109

Blood Pressure

Of the core indices of IRS, blood pressure is the endpoint that has been most extensively studied with respect to the influence of yoga-based interventions. Our search identified 37 studies from 6 countries, including 24 controlled studies. Over 75% of these studies report improvement in blood pressure with yoga or yoga-based interventions (Table 6). In a cross-sectional study of healthy mid-life men with similar lifestyle characteristics, Vyas and colleagues found those with both short-term and long-term experience in Raja yoga meditation had reduced diastolic blood pressure compared with those who were naïve to meditation.⁹⁶ Of the 36 remaining studies evaluating change in baseline blood pressure with yoga and yoga-based programs, only 8 studies (22%) found no evidence of improvement, including a small German study of changes in healthy young female practitioners during a single yoga session,¹¹⁰ 2 (20%) of the 10 uncontrolled studies,^{103,111} 2 (17%) of the 12 nonrandomized controlled trials,^{112,113} and 3 (25%) of the 13 RCTs.^{101,114,115} Of the 4 uncontrolled and nonrandomized controlled trials with negative findings, all were small studies of healthy young Indian men that demonstrated no change in blood pressure, but an improvement in heart rate^{111,112} and/or other indices of cardiovascular function^{103,111,113}; 2 of the 4 studies included active yoga asanas.^{103,112} RCTs yielding negative results included a 6-week study of healthy British elderly, and a 3-month study of healthy Indian boys that demonstrated no impact on blood pressure, but a significant reduction in heart rate and increase in heart rate variability (see below) in participants completing a yoga program compared with those receiving an aerobic exercise program¹¹⁴ or no intervention.¹¹⁵ In a third study of hypertensive Dutch elderly, the authors found a multimodal program that included training in yoga relaxation techniques to have no effect on 24-hour ambulatory blood pressure.¹⁰¹

In contrast, 28 eligible intervention studies published between 1970 and 2004 reported a beneficial effect of yoga and yoga-based programs on blood pressure change (Table 6). These include 8 uncontrolled studies of healthy^{90,104,116,117} and hypertensive adults,^{97,118,119} all but one of which⁹⁷ were conducted in India. Most of these studies were small, with only 3^{104,118,119} including 25 participants or

more. Among nonrandomized controlled trials, 9 studies of healthy adults^{93,98} and adults with hypertension^{109,120–124} and/or CAD⁸⁰ have demonstrated significant reductions in blood pressure among participants who completed a yoga-based intervention relative to controls receiving an exercise,¹²⁵ enhanced usual care,^{109,121–123} or no intervention.^{98,125} Likewise, 11 RCTs of healthy adults^{125–128} and patients with hypertension or other CVD risk factors^{79,87,102,107,129–131} have demonstrated significant blood pressure declines among subjects receiving a yoga-based intervention versus controls receiving usual care,^{107,129} enhanced usual care with diet and exercise,^{87,128} social contact,¹³¹ placebo therapy,^{129,130} or no active intervention.^{107,127} In a study of 95 female Japanese university students, Cusumano and colleagues demonstrated comparable, significant reductions over time in blood pressure among women receiving 3 weekly training sessions in either Hatha yoga or progressive muscle relaxation.¹²⁶ Overall, these studies demonstrated a 4.9% to 24.2% decline in diastolic blood pressure and a 2.6% to 21.3% decline in systolic blood pressure with yoga, with the magnitude of change varying with the study design and sample population (Table 5).

Yoga-based programs used in the 28 studies reporting positive findings ranged in duration from 30 days¹¹⁸ to 6 months¹¹⁹ in the uncontrolled studies, from 3 weeks¹²⁰ to 6 months^{93,109,124} in the controlled nonrandomized studies, and from 8 days¹³¹ to 12 months⁸⁷ in the RCTs. Interventions associated with blood pressure improvement included yoga routines both alone (N = 20) and in combination with other therapies (N = 8).[‡] Of those studies using yoga alone, 10 incorporated active yoga asanas[§] and 10 used relaxation postures alone^{117,119,123,131} or in combination with meditation.^{97,121,122,124,125,130}

Collectively, the findings of both controlled and uncontrolled studies suggest that even the relatively short-term practice of yoga or yoga-based programs may reduce blood pressure. However, because of potential biases and limitations characterizing most of these studies, caution is nonetheless warranted in drawing firm conclusions. For example, interpreting the positive findings of many of

[‡]See References 68, 79, 87, 98, 102, 104, 118, 129.

[§]See References 90, 93, 104, 107, 109, 116, 120, 126–128.

Table 6. Summarized Findings of Studies (1970–2004) Investigating the Effects of Yoga and Yoga-Based Programs on Blood Pressure, Stratified by Study Design and Intervention (Studies Reporting Beneficial Effects Are in Bold)

Findings, by Endpoint	Cross-Sectional 1 to 2 Sessions (Yoga Alone)	Study Design					
		Uncontrolled Clinical Trials (>2 sessions)			Non-Randomized Controlled Clinical Trials (>2 sessions)		
		Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions
Blood Pressure							
Reduced systolic and/or diastolic pressure	[96]	[97, 117]* [90, 116, 119] [103, 111]	[68, 104, 118]	[121, 122]* [93, 109, 120] [123]* [124] [112] [113] [¶]	[98] [†] [80]	[130]* [107, 125–128, 131]	[79, 87, 102, 129]
No change in systolic or diastolic pressure						[114, 115]	[101]
Systolic blood pressure							
Reduced							
Unchanged							
Diastolic blood pressure							
Reduced	[96]	[97, 117]* [90, 116, 119] [103, 111]	[68, 104, 118]	[121, 122]* [93, 109, 120] [123] [‡] [112] [113] [¶]	[80, 98]	[130]* [107, 125, 127, 128, 131] [114]	[87] [¶] [79, 129]
Unchanged							
Reduced mean blood pressure	[96]	[97, 117]* [90, 116, 119] [103, 111]	[68, 104, 118]	[109, 121, 122]* [123] [‡] [120] [113] [¶] [93, 112] [124]	[80, 98]	[130]* [107, 125, 127, 128, 131] [114]	[79, 87, 102, 129]
Unchanged						[126, 128]	[101]

*With biofeedback.

[†]Only among individuals with high initial SBP.

[‡]In both hypertensive and normotensive subjects.

[§]At 4 year follow up of individuals at risk for CVD.

[¶]Yoga PR in addition to exercise vs exercise alone.

^{||}P = .07 to .08.⁸⁷

Abbreviation: PR, pranayama.

these studies is hindered by absence of comparison groups,^{68,90,97,104,116–119} small sample sizes,^{||} reduced power due to stratified analyses^{87,118,127} or other factors,^{101,131} possible selection bias or confounding by lifestyle or other factors,^{||} absence of detailed information regarding the study population,^{**} the intervention,^{97,121,124} statistical^{††} or randomization methods, treatment allocation,^{80,124} or subject selection process,^{90,93,117,119,123} potential uncontrolled variation in exposure to the intervention,^{96,118} or loss to follow-up.^{80,116,121,122} Few studies reported blinding of outcome assessment,^{87,128–130} and several studies used yoga in combination with other interventions,^{‡‡} making it difficult to detect an effect specific to yoga. Other methodologic limitations are lack of randomization^{93,98,109,121–125} and inadequacies in statistical analyses or presentation, including lack of appropriate point estimates,^{68,93,98} adjustment for multiple comparisons,^{§§} and appropriate comparisons between^{93,109,120,122,123,128,131} or within^{68,93,117,123} groups.

Observed Effects of Yoga on Indices of Coagulopathy and Oxidative Stress

Procoagulant changes and damage caused by oxidative stress are thought to mediate many of the atherosclerotic and thrombotic changes that are associated with IRS^{13,14,132–134} and to play a pivotal role in the development and progression of CVD^{13,132–134} and diabetes.¹³⁵ Although few published studies have examined the possible effects of yoga and yoga-based interventions on coagulation profiles or oxidative stress, findings of these studies, summarized in Table 7, suggest that yoga may be instrumental in improving these important indices of CVD risk. An uncontrolled Indian study of 7 healthy young men naive to yoga demonstrated a significant decline in fibrinogen and a significant increase in fibrinolytic activity following a 4-month

yoga training course, in addition to a pronounced, although nonsignificant increase in platelet aggregation and activated partial thromboplastin time, and a rise in blood platelets.¹³⁶ Likewise, in a non-randomized controlled study of healthy German adults, Schmidt et al documented a significant fall in fibrinogen among participants completing a 3-month residential Kriya yoga program relative to community controls matched on age, gender, and baseline fibrinogen levels (N = 76 subjects, including 38 controls).⁹⁸ Although limited, these data suggest that yoga may foster beneficial changes in the coagulation and fibrinolytic systems, at least in healthy adults.

We identified 5 eligible studies investigating the influence of yoga on measures of oxidative stress, including 2 uncontrolled clinical trials, 2 nonrandomized controlled clinical trials, and one RCT (Table 7). All but one of these studies¹⁰⁰ were conducted in India, and all provide evidence that yoga may reduce oxidative stress in both healthy populations and those with chronic IRS-related disorders. In uncontrolled studies of Indian adults with uncomplicated diabetes mellitus,⁸⁵ or hypertension,⁶⁸ investigators demonstrated significant reductions in malonyldialdehyde (MDA), a circulating product of lipid peroxidation, following a 40-day yoga training course⁸⁵ and a 12-week yoga lifestyle intervention,⁶⁸ respectively. In agreement with these findings, nonrandomized controlled trials have shown significant reductions in MDA¹³⁷ and other markers of oxidative stress^{137,138} in healthy young men completing a 10-week¹³⁷ to 5-month¹³⁸ yoga training program relative to matched untrained controls. Observed changes in other oxidative stress indices include increases in antioxidants¹³⁸ and antioxidative enzymes,¹³⁸ and reductions in free radicals.¹³⁷ Offering further support for a direct effect of yoga on oxidative stress, Sharma et al documented significant improvement in several of these indices among trained participants during a single yoga session when compared with controls during rest¹³⁸ (Table 7). Similarly, in an RCT of 44 Thai patients with CAD, Jatuporn et al found those receiving a 4-month yoga-based lifestyle modification program but no medication showed significant increases in blood antioxidants relative to usual care controls, although MDA levels did not differ between groups.¹⁰⁰

Together, the 7 studies reviewed above offer support for a possible beneficial influence of yoga

^{||}See References 68, 90, 93, 97, 109, 116, 117, 119, 120, 124, 125.

^{††}See References 93, 96–98, 101, 107, 109, 118–120, 124–127.

^{**}See References 80, 93, 101, 104, 107, 109, 117, 119, 121, 123, 130, 131.

^{††}See References 68, 79, 90, 93, 97, 98, 102, 116–119, 121–123, 125, 131.

^{‡‡}See References 68, 79, 80, 87, 101, 102, 104, 118, 129, 130.

^{§§}See References 68, 87, 90, 98, 104, 109, 116, 123, 125, 127.

Table 7. Cross-Sectional Studies and Clinical Trials Investigating the Effects of Yoga and Yoga-Based Programs on Coagulation Profiles and Indices of Oxidative Stress, Stratified by Study Design and Intervention: Summarized Findings (Studies Reporting Beneficial Effects Are in Bold)

Findings, by Endpoint	Study Design							
	Cross-Sectional (Yoga Alone)	1 to 2 Sessions (Yoga Alone)	Uncontrolled Clinical Trials (>2 sessions)		Non-Randomized Controlled Clinical Trials (>2 sessions)		Randomized Controlled Trials (>2 sessions)	
			Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions
Coagulation/Fibrinolytic Profiles								
Reduced fibrinogen			[136]				[98]	
Increased fibrinolytic activity			[136]					
Prolonged platelet aggregation time			[136] (NS)					
Prolonged activated partial thromboplastin time			[136] (NS)					
Markers of Oxidative stress								
Increased Glutathione		[138]*			[138]			[100]
Increased Vitamins E, C								[100]
Increased total antioxidants								[100]
Super oxide dismutase (SOD)								
Increased		[138]			[138]			
No change					[137]			
Catalase								
Increased					[138]			
No change		[138]						
Reduced free radicals					[137]			
Malonyldialdehyde (MDA)								
Reduced			[85]	[68]	[137]			
No change								[100]

* $P < .1$; Yoga PR in addition to exercise vs exercise alone.

Abbreviations: NS, $P > .05$ (N = 7 subjects).

on oxidative stress and coagulation profiles, at least in certain populations. Limitations of these studies include lack of appropriate comparison groups,^{68,85,136} small sample sizes,^{68,85,136,138} possible selection bias or uncontrolled confounding by lifestyle and other factors,^{85,98,100,136–138} lack of randomization,^{98,137} inadequate information regarding the study population,^{85,138} subject selection,^{136,138} intervention,¹³⁸ or analytic methods,^{68,85,98,100,137} multiple interventions,^{68,100} and problems with data analysis or presentation.^{68,98,100,137} In addition, blinding of outcome assessment was not well-reported in any of the 7 studies identified.

Effects of Yoga on Markers of sympathetic/parasympathetic activation and cardiovagal function

Since 1970, numerous studies have investigated the effects of yoga on markers of sympathetic/parasympathetic activation and cardiovagal function. Our search identified a total of 42 studies from 6 countries, including 2 cross-sectional studies, 15 uncontrolled clinical trials, 16 nonrandomized controlled

trials, and 9 RCTs. Findings of these studies are summarized in Table 8. Although some studies have yielded inconsistent results, over 85% offer some evidence that yoga promotes a reduction in sympathetic activation, enhancement of cardiovagal function, and a shift in autonomic nervous system balance from primarily sympathetic to parasympathetic. Key changes, reviewed briefly below, include significant reductions in respiratory and heart rate, in cortisol concentrations, catecholamine levels, and renin activity, in skin conductance, and in cardiovascular response to stress, as well as significant increases in heart rate variability and baroreflex sensitivity.

Of the 27 studies investigating the effect of yoga on heart and/or respiratory rate, only 2 studies showed no change in either index, including a non-randomized controlled study of 10 young athletes¹¹³ and an RCT of healthy young men.¹²⁸ However, both of these latter studies demonstrated improvement in other indices of cardiovascular function. In contrast, 25 eligible intervention studies reported a beneficial effect of yoga and yoga-based programs on baseline respiratory rate, base-

Table 8. Cross-Sectional Studies and Clinical Trials Investigating the Effects of Yoga and Yoga-Based Programs on Markers of Sympathetic Activation and Reactivity, Cardiovascular Function, and Cardiovascular Recovery from Stress, Stratified by Study Design and Intervention: Summarized Findings (Studies Reporting Beneficial Effects Are in Bold)

Findings, by Endpoint	Study Design							
	Cross-Sectional (Yoga Alone)	1 to 2 Sessions (Yoga Alone)	Uncontrolled Clinical Trials (>2 Sessions)		Non-Randomized Controlled Clinical Trials (>2 Sessions)		Randomized Controlled Trials (>2 Sessions)	
			Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions
Respiratory and Heart Rate								
Respiratory rate								
Reduced			[103, 116, 139]	[104, 140]	[123]*		[141]	
Unchanged		[143]					[128]	
Heart rate (HR)								
Reduced baseline	[96]	[145]	[117] [†] [90, 111, 116]	[104, 151]	[123]* [93]	[98, 145]	[107, 114, 115, 125–127, 141]	
Reduced during/after yogic exercise		[142, 143]						
Unchanged			[103]	[140]	[113]		[128] [‡]	
Neuroendocrine markers								
Epinephrine/norepinephrine								
Increased plasma				[147]	[93]			
Reduced urinary				[68]	[120]	[98]		
Reduced urinary VMA				[68]			[146]	
Cortisol								
Reduced		[82]		[147] [158] [§]				
No change		[110]	[157]					
Increased excretion					[89]	[98]		
Increased basal skin resistance/ reduced skin conductance		[143, 145]	[117] [†]	[68, 104]			[125, 146]	
Reduced renin activity					[120, 124]			[102]
↓ Cardiovascular reactivity to stress								
Reduced DBP response to HGT				[151]				
Reduced DBP, SBP response to CPT							[152] [¶]	
Reduced DBP rise with exercise							[152] [¶]	
Reduced HR rise with exercise			[105]		[112, 153]	[127]		
↓ Cardiovascular recovery time after stress								
Faster HR return to baseline after exercise		[144]			[153]			
Faster DBP, SBP return to baseline after exercise								
Cold pressor test (CPT)							[152] [¶] [152] [¶]	
Increased Heart rate variability								
Increased (power of mid frequency/high frequency (HF) component)		[36, 143]		[140] [†]			[114]	
Increased (HF component of α index)							[114]	
Reduced (power of low frequency component)		[143]		[140] [†]				
Increased RRIV							[115]	
Other measures of ↑ baroreflex sensitivity								
Reduced HR, blood pressure response to head-up/head-down tilt			[111]		[120]			

*In both hypertensive and normotensive subjects.

[†]Only in subjects with Occupational Stress Index scores greater than the median.[¶]With biofeedback.^{||}17-hydroxy-corticosteroid.[§]Only in subjects with initial cortisol levels above the median; cortisol levels in those below the median did not change.[‡]Yoga PR in addition to exercise vs. exercise alone.

Abbreviations: CPT, cold pressor test; DBP, diastolic blood pressure; HGT, hand grip test; HR, heart rate; RRIV, R-R (heart rate) interval variation; SBP, systolic blood pressure; VMA, vanillylmandelic acid, a metabolite of norepinephrine.

line heart rate, and/or heart rate after exercise (Table 8). Seven studies (of 8 total) have documented a 3.8% to 60.3% reduction in respiratory rate with yoga relative to baseline^{103,104,116,139,140} and/or controls receiving usual care¹²³ or an aerobic exercise program¹⁴¹ (Table 5). These investigations include studies in healthy adults^{103,104,116,139,140} and children,¹⁴¹ as well as in hypertensive adults.¹²³ Similarly, 21 of 24 total clinical trials reported a 4.3% to 38.8% decline in heart rate relative to baseline,^{¶¶} engagement in a relaxing, nonyogic activity^{104,142–145} and/or controls receiving usual care,¹²³ an aerobic exercise program,¹⁴¹ or no intervention.^{98,115,125,127} These trials included studies of healthy^{¶¶} and hypertensive adults^{107,123} and of healthy children.^{115,141} In addition, in a cross-sectional study of healthy middle-aged men, Vyas and colleagues found those who had been practicing yoga meditation for 5 years or more to have significantly lower baseline heart rates than those who with no meditation experience.⁹⁶ Of those clinical trials demonstrating positive effects, yoga practice was associated with a 3.8 to 60.3% reduction in respiratory rate and a 4.3 to 38.8% decline in heart rate (Table 5).

Although findings regarding the effects of yoga on cortisol levels have been inconsistent, probably due in part to differing collection times and measurement methods, those regarding the influence of yoga on other neuroendocrine indices of sympathetic activation have been remarkably consistent overall (Table 8). Of the 13 studies investigating the influence of yoga practice on catecholamine levels, renin activity, and/or basal skin conductivity, 11 have demonstrated significant reductions in these markers of sympathoadrenal activation. These studies include investigations of healthy populations^{98,104,125,143,145} and those with hypertension^{68,102,117,120,124} or epilepsy.¹⁴⁶ Only 2 small uncontrolled studies of healthy Indian adults, conducted by the same group of investigators in the 1970s, have yielded contradictory results.^{93,147} With respect to catecholamine levels, 4 of the 6 eligible studies identified demonstrated significant declines in 24 hour urinary catecholamines among yoga program participants when compared with baseline⁶⁸ or to controls receiving usual care,¹⁴⁶

sham yoga,¹⁴⁶ or no intervention.⁹⁸ Studies evaluating the possible effects of yoga on renin activity and basal skin conductance have yielded even more consistent findings; of the 3 studies examining the former, and 7 studies investigating the latter endpoints, all documented significant reductions in these markers with yoga practice alone^{***} or in combination with lifestyle education.^{68,102}

Consistent with these results, findings from several studies suggest that yoga practice can lead to improvement in both cardiovascular reactivity to stress and cardiovascular recovery from stress (Table 8). Cardiovascular reactivity to stress, a factor strongly associated with insulin resistance^{148,149} is also a major independent predictor of hypertension, stroke, myocardial infarction, and cardiovascular mortality³¹; cardiovascular recovery from stress is a marker of vagal rebound that has likewise been strongly associated with CVD risk.¹⁵⁰ Documented changes associated with the practice of yoga and yoga-based programs in the 7 eligible studies identified include: (1) significantly reduced blood pressure and heart rate responses to stress relative to baseline^{105,151} or controls receiving nonyogic exercise,¹¹² usual care,¹⁵² or no intervention^{127,153}; and (2) accelerated cardiovascular recovery time from stress compared with nonyoga resting poses¹⁴⁴ or controls receiving usual care¹⁵² or no intervention.¹⁵³

Finally, 7 recent studies in Indian^{111,115,120,140} and European^{36,114} populations suggest that participating in yoga and yoga-based programs can improve heart rate variability and baroreflex sensitivity in both healthy^{36,111,114,115,140,143} and hypertensive populations¹²⁰ (Table 8). Low heart rate variability (HRV) and baroreflex sensitivity reflect impaired cardiovagal adaptability and suggest excessive sympathetic and/or insufficient parasympathetic tone.^{60,154,155} These alterations are, in turn, strong independent predictors of cardiovascular morbidity and mortality^{21,29,36} and are thought to mediate in part the effect of IRS-related abnormalities on these outcomes.^{21,156} In contrast, high HRV and baroreflex sensitivity are generally considered to indicate good cardiovagal adaptability and sympathovagal balance,¹⁵⁵ permitting greater responsiveness and sensitivity to changing environmental demands.

¶¶See References 90, 93, 104, 111, 116, 117, 125, 126.

¶¶See References 90, 93, 98, 104, 111, 114, 116, 117, 125–127, 142–145.

***See References 104, 117, 120, 124, 125, 143, 145, 146.

Collectively, this research suggests that even the short-term practice of yoga may produce marked reductions in sympathoadrenal activation, enhance cardiovagal tone, and promote sympathovagal balance. Several studies have demonstrated pronounced changes in markers of sympathetic/parasympathetic activity within a single yoga session (Table 8); others have documented significant beneficial alterations in baseline measurements in as little as 2 days in uncontrolled trials,¹⁴⁰ 3 weekly sessions in nonrandomized controlled trials,¹²⁶ and 8 days¹³¹ in RCTs. Interventions associated with improvement in markers of sympathetic/parasympathetic activity and cardiovagal function include yoga routines both alone (N = 29) and in combination with other therapies (N = 10); of those studies using yoga alone, 16 incorporated active yoga asanas, 7 used relaxation postures alone or in combination with meditation, and 5 used meditation or breathing exercises alone (Table 3).

Although evidence for a beneficial influence of yoga on sympathovagal balance is remarkably consistent overall across this diverse array of studies, many studies have limitations that hinder conclusive interpretation of findings. Many had no comparison groups,^{†††} small sample sizes^{†††} and/or comparison groups that differed in baseline characteristics.^{96,98,110,120,142,146,153} Over 50% of the studies identified had 25 or fewer subjects, and over 20% had less than 15. Others lacked detailed information on eligibility criteria and population characteristics,^{§§§} treatment allocation,^{113,124} and analytic methods,^{|||} and blinding of the outcome assessment was well reported in only a few studies.^{115,128} In others, loss to follow-up,^{105,114,116,146,157,158} variable exposure to or experience with the intervention,^{82,96,110–112,118,142–144} potential selection bias,^{¶¶¶} and uncontrolled confounding factors^{****} may have been important. The yoga intervention was not well described in some studies,^{112,124,147,152} and several used multiple inter-

ventions.^{†††} Other methodological problems included lack of randomization^{†††} or description of the randomization process,^{§§§§} inadequate data analysis or presentation,^{68,89,93,98,112,117,123} multiple comparisons,^{|||} or absence of appropriate intergroup comparisons.^{¶¶¶¶}

Effects of Yoga on Clinical Endpoints

Several studies in Indian,^{68,78,80,91,92,109,119} European,^{79,95,121,122,129,130} and American populations⁸⁷ have suggested that yoga may also be helpful in the management of CVD and other IRS-related chronic conditions. Our search identified a total of 14 eligible studies evaluating the influence of yoga and yoga-based interventions on clinical endpoints, including 4 uncontrolled trials, 4 nonrandomized controlled studies, and 6 RCTs (Table 9). In uncontrolled Indian studies of adults with hypertension^{68,119} and/or diabetes,^{68,91,92} completion of a simple^{91,119} or comprehensive yoga program^{68,92} was associated with a markedly reduced need for medication relative to baseline. In agreement with these findings, controlled studies of adults with hypertension^{109,121,122,130} risk factors for CVD⁷⁹ and/or CAD,⁸⁰ and diabetes⁹⁵ showed a decline in need for drug therapy among those enrolled in a yoga-based intervention compared with controls receiving usual care,⁹⁵ enhanced usual care,^{79,109,121,122} usual care plus a diet and exercise program,⁸⁰ or placebo therapy.¹³⁰ Controlled studies in Indian,⁷⁸ British,⁷⁹ and American⁸⁷ populations also suggest that yoga-based programs may attenuate signs, reduce complications, and improve the prognosis of those with frank or underlying disease. For example, in an RCT of Indian men with CAD, those enrolled in a 12-month comprehensive yoga program showed retardation of coronary atherosclerosis, increased regression and reduced progression of vascular lesions, and reduced anginal episodes relative to usual care controls⁷⁸; another, nonrandomized controlled study of CAD patients yielded similar results.⁸⁰ In a group of generally healthy

†††See References 68, 82, 89, 90, 103–105, 111, 116–118, 139, 140, 147, 151, 157, 158.

†††See References 36, 68, 82, 89, 90, 93, 103, 105, 110, 111, 113, 115–117, 120, 124, 125, 142, 145, 147, 151, 153.

§§§See References 82, 89, 90, 93, 104, 107, 112, 115, 117, 123, 139–142, 147, 151–153.

|||See References 68, 82, 90, 93, 98, 102, 105, 111–113, 116–118, 123, 125, 144, 147.

¶¶¶See References 82, 89, 90, 93, 105, 110, 117, 145, 146, 153.

****See References 93, 96, 98, 103, 107, 110–113, 115, 118, 120, 124–127, 139, 153, 157.

††††See References 68, 102, 104, 118, 140, 147, 151, 158.

****†See References 93, 98, 110, 112, 113, 120, 123–125, 153.

§§§§See References 102, 107, 114, 115, 127, 141, 152.

||||||See References 68, 90, 98, 104, 105, 110, 111, 113, 115, 116, 123, 125, 127, 139, 140, 142, 144, 151, 152, 157, 158.

¶¶¶¶¶See References 93, 112, 114, 115, 120, 123, 128, 141, 143, 153.

Table 9. Clinical Trials (>2 Sessions) Investigating the Effects of Yoga and Yoga-Based Programs on Clinical Outcomes, Stratified by Study Design and Intervention (Yoga, Including Yoga-Based Meditation Alone vs. in Combination with Other Interventions) (Studies Reporting Beneficial Effects Are in Bold)

Findings, by Endpoint	Study Design					
	Uncontrolled Clinical Trials		Non-Randomized Controlled Trials		Randomized Controlled Trials	
	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions	Yoga Alone	Multiple Interventions
Clinical endpoints						
Reduced need for medication						
Hypertension/CAD	[119]	[68]	[109, 121, 122]*	[80]	[130]*	[79][†]
Diabetes	[91]	[92] [68]			[95]	[78]
Reduced revascularization procedures related complications [†]						[78]
Number of subjects on treatment for:						
Hypertension (no difference) [†]						[79][†]
Other CVD complications (reduced) [†]						[79][†]
Reduced ECG evidence of ischemia [†]						[79][†]
Reduction in anginal episodes						[79][†] [78]
Reduced carotid intimal media thickness						[87]
Reduced progression of CAD				[80]		[78]
Increased regression of CAD				[80]		[78]
Reduced number of serious coronary events						[129](NS)[79][†]
Reduced number of hospitalizations				[80]		

*With biofeedback.

[†]At 4 year follow up of individuals at risk for CVD; no difference in number of Ss currently on treatment for hypertension.

Abbreviations: CAD, coronary artery disease; CVD, cardiovascular disease.

American seniors, those completing a 12-month comprehensive yoga-based program demonstrated a decline in carotid intimal media thickness, an indicator of carotid atherosclerosis, relative to those receiving usual care or a comprehensive medical, diet, and exercise intervention⁸⁷; the decline was correlated inversely with adherence, suggesting a direct relation between the practice of this program and atherosclerotic change. There is also some evidence that the clinical benefits observed following yoga-based programs might persist long term. In a 4-year follow-up of an earlier RCT in hypertensive adults, Patel et al found that those who had participated in an 8-week comprehensive yoga relaxation program were less likely than usual care controls to be receiving treatment for CVD-related complications, to have experienced a serious coronary event, or to have electrocardiographic evidence of ischemia.⁷⁹

Although promising, several of these studies have methodologic and other limitations, including absence of control groups,^{119,68,91,92} small sample sizes,^{68,95,109,119} lack of randomization,^{109,121,122} loss to follow-up,^{80,121,122} possible uncontrolled confounding factors,^{91,92,95,109,119} unclear statistical analyses,^{68,91} or absence of intergroup comparisons,^{109,122} multiple comparisons,^{68,87,91,109} appar-

ent lack of blinded outcome assessment,^{****} and multiple interventions.^{68,78–80,87,92,129,130} Several lacked adequate descriptions of the study population,^{80,91,109,119,121,130} sample selection process,¹¹⁹ intervention,¹²¹ treatment allocation,⁸⁰ or statistical methods.⁺⁺⁺⁺ In addition, reporting bias may have resulted in omission of negative findings, possibly inflating the apparent positive effect of yoga on clinical endpoints.

Yoga, Psychosocial Risk Factors for CVD, and Possible Underlying Mechanisms

Although methodologic and other limitations of many studies preclude drawing firm conclusions, the almost 70 investigations reviewed above nonetheless offer evidence that yoga may have a beneficial influence on many IRS-related risk factors for CVD, may improve clinical outcomes, and may aid in the management of CVD and other IRS-related conditions. In addition, a growing number of studies from both Western and non-Western countries suggest that yoga therapy may be effective in reducing psychosocial risk factors for CVD as well.

****See References 68, 79, 80, 91, 92, 109, 119, 121, 122.

++++See References 68, 79, 91, 92, 95, 119, 121, 122.




Figure 1. Hypothesized Pathways by Which Yoga Intervention May Enhance Cardiovascular Risk Profiles

For example, yoga has been reported to decrease perceived stress^{109,159,160} and reactivity to stressors,¹¹⁰ enhance stress-related coping,¹⁶¹ reduce symptoms of depression^{127,128,162–165} and anxiety,^{66,127,161,162,166,167} and decrease anger,^{162,168} tension, and fatigue.^{162,169} Likewise, recent studies in India^{66,159,170} and the United States¹⁷¹ have indicated that yoga practice may improve sleep in both healthy^{66,159} and chronically ill populations.^{170,171}

Although the mechanisms underlying the putative beneficial effects of yoga therapy on cardiovascular risk profiles are not yet well understood, the observed changes probably occur primarily through 2 pathways. First, by reducing the activation and reactivity of the sympathoadrenal system and the hypothalamic pituitary adrenal (HPA) axis and promoting feelings of well-being, yoga may alleviate the effects of stress and foster multiple positive downstream effects on neuroendocrine status, metabolic function and related inflammatory responses (Figure 1, pathway 1). Second, by directly stimulating the vagus nerve, yoga may enhance parasympathetic output and thereby shift the autonomic nervous system balance from primarily sympathetic to parasympathetic, leading to positive changes in cardiac-vagal function, in mood and energy state, and in related neuroendocrine, metabolic, and inflammatory responses (Figure 1, pathway 2). Yoga may also provide a positive source of social support, a factor strongly associated with CVD risk,^{32,38} may aid in improving health-related

attitudes and lifestyle choices, in part by enhancing psychological well-being,³⁸ and in this way may play an important role in CVD prevention and health promotion.³⁸

Summary and Conclusions

In summary, a substantial body of published research suggests that yoga may be instrumental in improving core indices of the IRS, including glucose tolerance and insulin sensitivity, lipid profiles, anthropometric characteristics, and blood pressure in both healthy populations and those with chronic IRS-related conditions. In addition, more limited data indicate that yoga may improve coagulation profiles, at least in healthy populations, and reduce oxidative stress in both healthy adults and populations with serious chronic disorders. There is also mounting evidence to suggest that yoga may reduce sympathetic activation and enhance cardio-vagal function, factors strongly associated with both IRS and CVD risk. In addition, findings from several studies, including several recent controlled trials,^{78,80,87} indicate that yoga may be useful in the management of CVD and related conditions, and in the prevention of CVD in those at risk.

However, despite the growing popularity and apparent therapeutic potential of yoga in the prevention and management of CVD, and the now considerable body of published literature on this topic, relatively few rigorous, controlled studies have been conducted on the effects of yoga on

physiologic and anthropometric indices of cardiovascular risk; even fewer have been conducted in western populations. Since 1970, approximately 70% of clinical trials investigating the effects of yoga practice on CVD disease risk and clinical outcomes have been conducted in Indian populations, for whom yoga is an integral part of a long-standing cultural and spiritual tradition. Interpretation of many existing studies is also limited by small sample sizes, lack of appropriate control groups, inadequate description of methods, selection bias, failure to adjust for lifestyle characteristics and other potential confounders, exposure to multiple interventions, inadequacies in statistical analysis and presentation, or other methodological problems. In many of the controlled trials, treatment allocation was not randomized, and direct statistical comparisons were not made to control groups, potentially biasing the findings. In addition, the large variation in the nature, duration, intensity, and delivery methods of the yoga-based interventions used, even among studies using yoga practice alone, renders comparison across studies difficult. Publication bias may also play a role in the selective reporting of positive results,¹⁷² particularly in non-Western countries, where yoga is more widely embraced and more likely to be incorporated into medical treatment. Although existing RCTs have yielded results consistent overall with those of nonrandomized and uncontrolled studies, suggesting that the positive findings of these latter studies do not simply reflect bias because of poor study design, clearly, additional high quality RCTs are warranted. Specifically, rigorous, well-designed studies are needed to elucidate the effects of standardized yoga programs on IRS-related indices of CVD risk and related clinical endpoints, especially in American and other Western populations that remain underrepresented in the current literature.

We thank Dr. Ada Jacox for very helpful comments on an earlier draft of this article.

References

1. American Heart Association, 2004 Heart and Stroke Statistical Update. American Heart Association, 2004.
2. Bonow RO. Primary prevention of cardiovascular disease: a call to action. *Circulation* 2002;106:3140–1.
3. Lopez-Candales A. Cardiovascular diseases: a review of the Hispanic perspective. Awareness is the first step to action. *J Med* 2002;33:227–45.
4. Watkins LO. Epidemiology and burden of cardiovascular disease. *Clin Cardiol* 2004;27(6 Suppl 3):III2–6.
5. Pramparo P. The epidemiology of hypertension in South America. *J Hum Hypertens* 2002;16(Suppl 1):S3–6.
6. Posner B, Franz M, Quatromoni P. Nutrition and the global risk for chronic diseases: the INTER-HEALTH Nutrition Initiative. *Nutr Rev* 1994;52:210–7.
7. Popkin B. The nutrition transition in low-income countries. *Nutr Rev* 1994;52:285–98.
8. Grundy S. Hypertriglyceridemia, insulin resistance, and the metabolic syndrome. *Am J Cardiol* 1999;83:25F–9F.
9. Ginsberg H. Insulin resistance and cardiovascular disease. *J Clin Invest* 2000;106:453–8.
10. Isomaa B. A major health hazard: the metabolic syndrome. *Life Sci* 2003;73:2395–411.
11. Reilly M, Rader D. The metabolic syndrome: more than the sum of its parts? *Circulation* 2003;108:1546–51.
12. Reaven G, Abbasi F, McLaughlin T. Obesity, insulin resistance, and cardiovascular disease. *Recent Prog Horm Res* 2004;59:207–23.
13. Juhan-Vague I, Morange PE, Alessi MC. The insulin resistance syndrome: implications for thrombosis and cardiovascular disease. *Pathophysiol Haemostasis Thromb* 2002;32:269–73.
14. Ginsberg HN, Huang LS. The insulin resistance syndrome: impact on lipoprotein metabolism and atherothrombosis. *J Cardiovasc Risk* 2000;7:325–31.
15. Natali A, Ferrannini E. Hypertension, insulin resistance, and the metabolic syndrome. *Endocrinol Metab Clin North Am* 2004;33:417–29.
16. Wheatcroft SB, Williams IL, Shah AM, Kearney MT. Pathophysiological implications of insulin resistance on vascular endothelial function. *Diabetic Med* 2003;20:255–68.
17. Hsueh W, Lyon C, Quinones M. Insulin resistance and the endothelium. *Am J Med* 2004;117:109–17.
18. Fernandez-Real JM, Ricart W. Insulin resistance and chronic cardiovascular inflammatory syndrome. *Endo Rev* 2003;24:278–301.
19. Byrne CD. Programming other hormones that affect insulin. *Br Med Bull* 2001;60:153–71.
20. Innes K, Wimsatt J. Pregnancy-induced hypertension and insulin resistance: evidence for a connection. *Acta Obstet Gynecol Scand* 1999;78:263–84.
21. Gadegbeku CA, Dhandayuthapani A, Sadler ZE, Egan BM. Raising lipids acutely reduces baroreflex sensitivity. *American Journal of Hypertension* 2002;15:479–85.
22. Beske SD, Alvarez GE, Ballard TP, Davy KP. Re-

- duced cardiovagal baroreflex gain in visceral obesity: implications for the metabolic syndrome. *Am J Physiol* 2002;282:H630–5.
23. Pikkujamsa SM, Huikuri HV, Airaksinen KE, Rantala AO, Kauma H, Lilja M, et al. Heart rate variability and baroreflex sensitivity in hypertensive subjects with and without metabolic features of insulin resistance syndrome. *Am J Hypertens* 1998; 11:523–31.
24. Bjorntorp P, Holm G, Rosmond R. Hypothalamic arousal, insulin resistance and type 2 diabetes mellitus. *Diabetic Med* 1999;16:373–83.
25. Egan BM. Insulin resistance and the sympathetic nervous system. *Curr Hypertens Rep* 2003;5:247–54.
26. Esler M. The sympathetic system and hypertension. *Am J Hypertens* 2000;13:99S–105S.
27. Palatini P. Sympathetic overactivity in hypertension: a risk factor for cardiovascular disease. *Curr Hypertens Rep* 2001;3(Suppl 1):S3–9.
28. Rozanski A, Blumenthal JA, Kaplan J. Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation* 1999;99:2192–217.
29. Singh RB, Kartik C, Otsuka K, Pella D, Pella J. Brain-heart connection and the risk of heart attack. *Biomed Pharmacother* 2002;56(Suppl 2):257s–65s.
30. Everson SA, Lynch JW, Chesney MA, et al. Interaction of workplace demands and cardiovascular reactivity in progression of carotid atherosclerosis: population based study. *BMJ* 1997;314:553–8.
31. Jennings JR, van der Molen MW, Somsen RJ, Graham R, Gianaros PJ. Vagal function in health and disease: studies in Pittsburgh. *Physiol Behav* 2002; 77:693–8.
32. Bunker SJ, Colquhoun DM, Esler MD, et al. “Stress” and coronary heart disease: psychosocial risk factors. *Med J Aust* 2003;178:272–6.
33. Bjorntorp P, Holm G, Rosmond R, Folkow B. Hypertension and the metabolic syndrome: closely related central origin?. *Blood Pressure* 2000;9:71–82.
34. Hjemdahl P. Stress and the metabolic syndrome: an interesting but enigmatic association. *Circulation* 2002;106:2634–6.
35. Brunner EJ, Hemingway H, Walker BR, et al. Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome: nested case-control study. *Circulation* 2002;106:2659–65.
36. Bernardi L, Sleight P, Bandinelli G, et al. Effect of rosary prayer and yoga mantras on autonomic cardiovascular rhythms: comparative study. *BMJ* 2001; 323:1446–9.
37. Bjorntorp P. Body fat distribution, insulin resistance, and metabolic diseases. *Nutrition* 1997;13: 795–803.
38. Vitaliano PP, Scanlan JM, Zhang J, Savage MV, Hirsch IB, Siegler IC. A path model of chronic stress, the metabolic syndrome, and coronary heart disease. *Psychosom Med* 2002;64:418–35.
39. Weber B, Schweiger U, Deuschle M, Heuser I. Major depression and impaired glucose tolerance. *Exp Clin Endocrinol Diabetes* 2000;108:187–90.
40. Okamura F, Tashiro A, Utumi A, Imai T, Suchi T, Tamura D, et al. Insulin resistance in patients with depression and its changes during the clinical course of depression: minimal model analysis. *Metabolism* 2000;49:1255–60.
41. Raikonen K, Matthews KA, Kuller LH. The relationship between psychological risk attributes and the metabolic syndrome in healthy women: antecedent or consequence? *Metabolism* 2002;51:1573–7.
42. Lucini D, Norbiato G, Clerici M, Pagani M. Hemodynamic and autonomic adjustments to real life stress conditions in humans. *Hypertension* 2002;39: 184–8.
43. Levenstein S, Smith M, Kaplan G. Psychosocial predictors of hypertension in men and women. *Arch Intern Med* 2001;161:1341–6.
44. Pickering T, Clemow L, Davidson K, Gerin W. Behavioral cardiology – has its time finally arrived? *Mt Sinai J Med* 2003;70:101–12.
45. Davidson K, Jonas B, Dixon K, Markovitz J. Do depression symptoms predict early hypertension incidence in young adults in the CARDIA study? *Coronary Artery Risk Development in Young Adults*. *Arch Int Med* 2000;160:1495–500.
46. Ariyo A, Haan M, Tangen C, et al. Depressive symptoms and risks of coronary heart disease and mortality in elderly Americans. *Cardiovascular Health Study Collaborative Research Group*. *Circulation* 2000;102:1773–91.
47. Chen CC, Lu FH, Wu JS, Chang CJ. Correlation between serum lipid concentrations and psychological distress. *Psychiatry Res* 2001;102:153–62.
48. Spieker LE, Hurlimann D, Ruschitzka F, et al. Mental stress induces prolonged endothelial dysfunction via endothelin-A receptors. *Circulation* 2002;105:2817–20.
49. Cannon RO 3rd. Role of nitric oxide in cardiovascular disease: focus on the endothelium. *Clin Chem* 1998;44:1809–19. Erratum in: *Clin Chem* 1998;44: 2070.
50. Sherwood A, Johnson K, Blumenthal JA, Hinderliter AL. Endothelial function and hemodynamic responses during mental stress. *Psychosomatic Medicine* 1999;61:365–70.
51. Eaton W, Armenian H, Gallo J, Pratt L, Ford D. Depression and risk for onset of type II diabetes. A prospective population-based study. *Diabetes Care* 1996;19:1097–102.
52. Black PH, Garbutt LD. Stress, inflammation and cardiovascular disease. *J Psychosom Res* 2002;52:1–23.
53. von Kanel R, Mills PJ, Fainman C, Dimsdale JE. Effects of psychological stress and psychiatric dis-

- orders on blood coagulation and fibrinolysis: a biobehavioral pathway to coronary artery disease? *Psychosom Med* 2001;63:531–44.
54. Goodman E, Whitaker R. A prospective study of the role of depression in the development and persistence of adolescent obesity. *Pediatrics* 2002;110:497–504.
55. Nelson TL, Palmer RF, Pedersen NL, Miles TP. Psychological and behavioral predictors of body fat distribution: age and gender effects. *Obes Res* 1999;7:199–207.
56. Everson SA, Kaplan GA, Goldberg DE, Salonen R, Salonen JT. Hopelessness and 4-year progression of carotid atherosclerosis. The Kuopio Ischemic Heart Disease Risk Factor Study. *Arterioscler Thromb Vasc Biol* 1997;17:1490–5.
57. Greenwood DC, Muir KR, Packham CJ, Madeley RJ. Coronary heart disease: a review of the role of psychosocial stress and social support. *J Public Health Med* 1996;18:221–31.
58. Kop WJ. The integration of cardiovascular behavioral medicine and psychoneuroimmunology: new developments based on converging research fields. *Brain Behav Immun* 2003;17:233–7.
59. Todaro JF, Shen BJ, Niaura R, Spiro A 3rd, Ward KD. Effect of negative emotions on frequency of coronary heart disease (The Normative Aging Study). *Am J Cardiol* 2003;92:901–6.
60. Carney RM, Freedland KE, Miller GE, Jaffe AS. Depression as a risk factor for cardiac mortality and morbidity: a review of potential mechanisms. *J Psychosom Res* 2002;53:897–902.
61. Davidson D. Preventive cardiology. Baltimore, 1991.
62. Carr MC. The emergence of the metabolic syndrome with menopause. *J Clin Endocrinol Metab* 2003;88:2404–11.
63. Ford ES, Giles WH, Mokdad AH. Increasing prevalence of the metabolic syndrome among U.S. adults. *Diabetes Care* 2004;27:2444–9.
64. Seidell JC. Obesity, insulin resistance and diabetes—a worldwide epidemic. *Br J Nutr* 2000;83 Suppl 1:S5–8.
65. Lindsay R, Howard B. Cardiovascular risk associated with the metabolic syndrome. *Curr Diabetes Rep* 2004;4:63–8.
66. Telles S, Naveen K. Yoga for rehabilitation: an overview. *Indian J Med Sci* 1997;51:123–7.
67. Pandya D, Vyas V, Vyas S. Mind-body therapy in the management and prevention of coronary disease. *Compr Ther* 1999;25:283–93.
68. Damodaran A, Malathi A, Patil N, Shah N, Suryavansihi Marathe S. Therapeutic potential of yoga practices in modifying cardiovascular risk profile in middle aged men and women. *J Assoc Physicians India* 2002;50:633–40.
69. Sahay B, Sahay R. Lifestyle modification in management of diabetes mellitus. *J Indian Med Assoc* 2002;100:178–80.
70. Garfinkel M, Schumacher HJ. Yoga. *Rheum Dis Clin N Am* 2000;26:125–32.
71. Chandler K. The emerging field of yoga therapy. *Hawaii Med J* 2001;60:286–7.
72. Raub J. Psychophysiologic effects of Hatha Yoga on musculoskeletal and cardiopulmonary function: a literature review. *J Altern Complementary Med* 2002;8:797–812.
73. Saper R, Eisenberg D, Davis R, Culpepper L, Phillips R. Prevalence and patterns of adult yoga use in the United States: results of a national survey. *Altern Ther Health Med* 2004;10:44–9.
74. Labarthe D, Ayala C. Nondrug interventions in hypertension prevention and control. *Cardiol Clin* 2002;20:249–63.
75. Gimbel M. Yoga, meditation, and imagery: clinical applications. *Nurse Practitioner Forum* 1998;9:243–55.
76. Wood C. Mood change and perceptions of vitality: a comparison of the effects of relaxation, visualization and yoga. *J Roy Soc Med* 1993;86:254–8.
77. Miller J, Fletcher K, Kabat-Zinn J. Three-year follow-up and clinical implications of a mindfulness meditation-based stress reduction intervention in the treatment of anxiety disorders. *Gen Hosp Psychiatry* 1995;17:192–200.
78. Manchanda S, Narang R, Reddy K, et al. Retardation of coronary atherosclerosis with yoga lifestyle intervention. *J Assoc Physicians India* 2000;48:687–94.
79. Patel C, Marmot M, Terry D, Carruthers M, Hunt B, Patel M. Trial of relaxation in reducing coronary risk: four year follow-up. *Br Med J* 1985;290:1103–6.
80. Yogendra J, Yogendra H, Ambardekar S, et al. Beneficial effects of yoga lifestyle on reversibility of ischaemic heart disease: Caring Heart Project of International Board of Yoga. *J Assoc Physicians India* 2004;52:283–9.
81. King MS, Carr T, D'Cruz C. Transcendental meditation, hypertension and heart disease. *Aust Fam Physician* 2002;31:164–8.
82. Kamei T, Toriumi Y, Kimura H, Ohno S, Kumano H, Kimura K. Decrease in serum cortisol during yoga exercise is correlated with alpha wave activation. *Percept Mot Skills* 2000;90:1027–32.
83. Wang C, Collet JP, Lau J. The effect of Tai Chi on health outcomes in patients with chronic conditions: a systematic review. *Arch Intern Med* 2004;164:493–501.
84. Verhagen AP, Immink M, van der Meulen A, Bierma-Zeinstra SM. The efficacy of Tai Chi Chuan in older adults: a systematic review. *Fam Pract* 2004;21:107–13.
85. Singh S, Malhotra V, Singh K, Sharma S. A preliminary report on the role of yoga asanas on oxi-

- dative stress in non-insulin dependent diabetes. *Indian J Clin Biochem* 2001;16:216–20.
86. Malhotra V, Singh S, Singh K, Gupta P, Sharma S, Madhu S, et al. Study of yoga asanas in assessment of pulmonary function in NIDDM patients. *Indian J Physiol Pharmacol* 2002;46:313–20.
87. Fields JZ, Walton KG, Schneider RH, et al. Effect of a multimodality natural medicine program on carotid atherosclerosis in older subjects: a pilot trial of Maharishi Vedic Medicine. *Am J Cardiol* 2002;89:952–8.
88. Naruka J, Mathur R, Mathur A. Effect of pranayama practices on fasting blood glucose and serum cholesterol. *Indian J Med Sci* 1986;40:149–52.
89. Udupa KN, Singh RH. The scientific basis of yoga. *JAMA* 1972;220:1365.
90. Joseph S, Sridharan K, Patil S, et al. Study of some physiological and biochemical parameters in subjects undergoing yogic training. *Indian J Med Res* 1981;74:120–4.
91. Divekar M, Bhat M, Mulla A. Effect of yoga therapy in diabetes and obesity. *J Diab Assoc Ind* 1978;17:75–8.
92. Jain S, Uppal A, Bhatnagar S, Talukdar B. A study of response pattern of non-insulin dependent diabetics to yoga therapy. *Diabetes Res Clin Pract* 1993;19:69–74.
93. Udupa KN, Singh RH, Settiwar RM. Physiological and biochemical studies on the effect of yogic and certain other exercises. *Indian J Med Res* 1975;63:620–4.
94. Sahay BK, Sadasivudu B, Yogi R, et al. Biochemical parameters in normal volunteers before and after yogic practices. *Indian J Med Res* 1982;76(Suppl):144–8.
95. Monro R, Power J, Coumar A, Dandona P. Yoga therapy for NIDDM: a controlled trial. *Complementary Med Res* 1992;6:66–8.
96. Vyas R, Dikshit N. Effect of meditation on respiratory system, cardiovascular system and lipid profile. *Indian J Physiol Pharmacol* 2002;46:487–91.
97. Patel C. Reduction of serum cholesterol and blood pressure in hypertensive patients by behaviour modification. *J Roy Coll Gen Practitioners* 1976;26:211–5.
98. Schmidt T, Wijga A, Von Zur Muhlen A, Brabant G, Wagner T. Changes in cardiovascular risk factors and hormones during a comprehensive residential three month kriya yoga training and vegetarian nutrition. *Acta Physiol Scand Suppl* 1997;640:158–62.
99. Mahajan A, Reddy K, Sachdeva U. Lipid profile of coronary risk subjects following yogic lifestyle intervention. *Indian Heart J* 1999;51:37–40.
100. Jatuporn S, Sangwatanaroj S, Saengsiri A, et al. Short-term effects of an intensive lifestyle modification program on lipid peroxidation and antioxidant systems in patients with coronary artery disease. *Clin Hemorheol Microcirc* 2003;29:429–36.
101. van Montfrans G, Karemaker J, Wieling W, Dunning A. Relaxation therapy and continuous ambulatory blood pressure in mild hypertension: a controlled study. *BMJ* 1990;300:1368–72.
102. Patel C, Marmot MG, Terry DJ. Controlled trial of biofeedback-aided behavioural methods in reducing mild hypertension. *Br Med J (Clin Res Ed)* 1981;282:2005–8.
103. Satyanarayana M, Rajeswari KR, Rani NJ, Krishna CS, Rao PV. Effect of Santhi Kriya on certain psychophysiological parameters: a preliminary study. *Indian J Physiol Pharmacol* 1992;36:88–92.
104. Telles S, Nagarathna R, Nagendra HR, Desiraju T. Physiological changes in sports teachers following 3 months of training in Yoga. *Indian J Med Sci* 1993;47:235–8.
105. Raju PS, Prasad KV, Venkata RY, Murthy KJ, Reddy MV. Influence of intensive yoga training on physiological changes in 6 adult women: a case report. *J Altern Complementary Med* 1997;3:291–5.
106. Bera TK, Rajapurkar MV. Body composition, cardiovascular endurance and anaerobic power of yogic practitioner. *Indian J Physiol Pharmacol* 1993;37:225–8.
107. Murugesan R, Govindarajulu N, Bera T. Effect of selected yogic practices on the management of hypertension. *Indian J Physiol Pharmacol* 2000;44:207–10.
108. Tran MD, Holly RG, Lashbrook J, Amsterdam EA. Effects of Hatha Yoga practice on the health-related aspects of physical fitness. *Prev Cardiol* 2001;4:165–70.
109. Latha Kaliappan K. Yoga, pranayama, thermal biofeedback techniques in the management of stress and high blood pressure. *J Indian Psychol* 1991;9:36–46.
110. Schell F, Allolio B, Schonecke O. Physiological and psychological effects of Hatha-Yoga exercise in healthy women. *Int J Psychosom* 1994;41:46–52.
111. Konar D, Latha R, Bhuvaneshwaran JS. Cardiovascular responses to head-down-body-up postural exercise (Sarvangasana). *Indian J Physiol Pharmacol* 2000;44:392–400.
112. Gopal KS, Bhatnagar OP, Subramanian N, Nishith SD. Effect of yogasanas and pranayamas on blood pressure, pulse rate and some respiratory functions. *Indian J Physiol Pharmacol* 1973;17:273–6.
113. Raju PS, Madhavi S, Prasad KV, et al. Comparison of effects of yoga and physical exercise in athletes. *Indian J Med Res* 1994;100:81–6.
114. Bowman A, Clayton R, Murray A, Reed J, Subhan M, Ford G. Effects of aerobic exercise training and yoga on the baroreflex in healthy elderly persons. *Eur J Clin Invest* 1997;27:443–9.
115. Udupa K, Madanmohan, Bhavanani AB, Vijayalakshmi P, Krishnamurthy N. Effect of pranayam

- training on cardiac function in normal young volunteers. *Indian J Physiol Pharmacol* 2003;47:27–33.
116. Anantharaman R, Kabir R. A study of Yoga. *J Psychol Res* 1984;28:97–101.
117. Mogra A, Singh G. Effect of biofeedback and yogic relaxation exercise on the blood pressure levels of hypertensives: a preliminary study. *Aviation Med* 1986;30:68–75.
118. Lakshmikanthan C, Alagesan R, Thanikachalam S, et al. Long term effects of yoga on hypertension and/or coronary artery disease. *J Assoc Physicians India* 1979;27:1055–8.
119. Sundar S, Agrawal S, Singh V, Bhattacharya S, Udupa K, Vaish S. Role of yoga in management of essential hypertension. *Acta Cardiol* 1984;39:203–8.
120. Selvamurthy W, Sridharan K, Ray U, et al. A new physiological approach to control essential hypertension. *Indian J Physiol Pharmacol* 1998;42:205–13.
121. Patel C. 12-month follow-up of yoga and bio-feedback in the management of hypertension. *Lancet* 1975;1:62–4.
122. Patel C, Datey KK. Relaxation and biofeedback techniques in the management of hypertension. *Angiology* 1976;27:106–13.
123. Chaudhary AK, Bhatnagar HN, Bhatnagar LK, Chaudhary K. Comparative study of the effect of drugs and relaxation exercise (yoga shavasana) in hypertension. *J Assoc Physicians India* 1988;36:721–3.
124. Stone RA, DeLeo J. Psychotherapeutic control of hypertension. *N Engl J Med* 1976;294:80–4.
125. Bagga OP, Gandhi A. A comparative study of the effect of Transcendental Meditation (T.M.) and Shavasana practice on cardiovascular system. *Indian Heart J* 1983;35:39–45.
126. Cusumano JA, Robinson SE. The short-term psychophysiological effects of hatha yoga and progressive relaxation on female Japanese students. *Appl Psychol* 1992;42:77–90.
127. Ray U, Mukhopadhyaya S, Purkayastha S, et al. Effect of yogic exercises on physical and mental health of young fellowship course trainees. *Indian J Physiol Pharmacol* 2001;45:37–53.
128. Harinath K, Malhotra AS, Pal K, et al. Effects of Hatha yoga and Omkar meditation on cardiorespiratory performance, psychologic profile, and melatonin secretion. *J Altern Complementary Med* 2004;10:261–8.
129. Patel C, Marmot M. Can general practitioners use training in relaxation and management of stress to reduce mild hypertension? *Br Med J (Clin Res Ed)* 1988;296:21–4.
130. Patel C, North WR. Randomised controlled trial of yoga and bio-feedback in management of hypertension. *Lancet* 1975;2:93–5.
131. Broota A, Varma R, Singh A. Role of relaxation in hypertension. *J Indian Acad Appl Psychol* 1995;21:29–36.
132. Dhalla N, Temsah R, Netticadan T. Role of oxidative stress in cardiovascular diseases. *J Hypertens* 2000;18:655–73.
133. Lee KU. Oxidative stress markers in Korean subjects with insulin resistance syndrome. *Diabetes Res Clin Pract* 2001;54(Suppl 2):S29–33.
134. Kohler HP. Insulin resistance syndrome: interaction with coagulation and fibrinolysis. *Swiss Medical Weekly* 2002;132:241–52.
135. Maritim A, Sanders R, Watkins JB 3rd. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol* 2003;17:24–38.
136. Chohan IS, Nayar HS, Thomas P, Geetha NS. Influence of yoga on blood coagulation. *Thromb Haemostasis* 1984;51:196–7.
137. Bhattacharya S, Pandey U, Verma N. Improvement in oxidative status with yogic breathing in young healthy males. *Indian J Physiol Pharmacol* 2002;46:349–54.
138. Sharma H, Sen S, Singh A, Bhardwaj NK, Kochupillai V, Singh N. Sudarshan Kriya practitioners exhibit better antioxidant status and lower blood lactate levels. *Biol Psychol* 2003;63:281–91.
139. Joshi L, Joshi V, Gokhale L. Effect of short term 'Pranayam' practice on breathing rate and ventilatory functions of lung. *Indian J Physiol Pharmacol* 1992;36:105–8.
140. Vempati R, Telles S. Baseline occupational stress levels and physiological responses to a two day stress management program. *J Indian Psychol* 2000;18:33–7.
141. Telles S, Narendran S, Raghuraj P, Nagarathna R, Nagendra H. Comparison of changes in autonomic and respiratory parameters of girls after yoga and games at a community home. *Percept Mot Skills* 1997;84:251–7.
142. Harte JL, Eifert GH, Smith R. The effects of running and meditation on beta-endorphin, corticotropin-releasing hormone and cortisol in plasma, and on mood. *Biol Psychol* 1995;40:251–65.
143. Vempati R, Telles S. Yoga-based guided relaxation reduces sympathetic activity judged from baseline levels. *Psychol Rep* 2002;90:487–94.
144. Bera TK, Gore MM, Oak JP. Recovery from stress in two different postures and in Shavasana—a yogic relaxation posture. *Indian J Physiol Pharmacol* 1998;42:473–8.
145. Telles S, Nagarathna R, Nagendra HR. Autonomic changes during "OM" meditation. *Indian J Physiol Pharmacol* 1995;39:418–20.
146. Panjwani U, Gupta HL, Singh SH, Selvamurthy W, Rai UC. Effect of Sahaja yoga practice on stress management in patients of epilepsy. *Indian J Physiol Pharmacol* 1995;39:111–6.
147. Udupa KN, Singh RH, Dwivedi KN, Pandey HP,

- Rai V. Comparative biochemical studies on meditation. *Indian J Med Res* 1975;63:1676–9.
148. Nazzaro P, Triggiani R, Ciancio L, Scarano AM, Pirrelli AM. Insulin resistance in essential hypertension: a psychophysiological approach to the “chicken and egg” question. *Nutr Metab Cardiovasc Dis* 2000;10:275–86.
149. Moan A, Nordby G, Rostrup M, Eide I, Kjeldsen SE. Insulin sensitivity, sympathetic activity, and cardiovascular reactivity in young men. *Am J Hypertens* 1995;8:268–75.
150. Mezzacappa ES, Kelsey RM, Katkin ES, Sloan RP. Vagal rebound and recovery from psychological stress. *Psychosom Med* 2001;63:650–7.
151. Khanam AA, Sachdeva U, Guleria R, Deepak KK. Study of pulmonary and autonomic functions of asthma patients after yoga training. *Indian J Physiol Pharmacol* 1996;40:318–24.
152. Patel C. Yoga and biofeedback in the management of ‘stress’ in hypertensive patients. *Clin Sci Mol Med Suppl* 1975;2:171s–4s.
153. Muralidhara DV, Ranganathan KV. Effect of yoga practice on Cardiac Recovery Index. *Indian J Physiol Pharmacol* 1982;26:279–83.
154. Buchman TG, Stein PK, Goldstein B. Heart rate variability in critical illness and critical care. *Curr Opin Crit Care* 2002;8:311–5.
155. Pumpura J, Howorka K, Groves D, Chester M, Nolan J. Functional assessment of heart rate variability: physiological basis and practical applications. *Int J Cardiol* 2002;84:1–14.
156. Miller AW, Sims JJ, Canavan A, Hsu T, Ujhelyi MR. Impaired vagal reflex activity in insulin-resistant rats. *J Cardiovasc Pharmacol* 1999;33:698–702.
157. Janakiramaiah N, Gangadhar B, Murthy P, et al. Therapeutic efficacy of Sudarshan Kriya Yoga (SKY) in dysthymic disorder. *NIMHANS J* 1998;16:21–8.
158. Carlson LE, Speca M, Patel KD, Goodey E. Mindfulness-based stress reduction in relation to quality of life, mood, symptoms of stress and levels of cortisol, dehydroepiandrosterone sulfate (DHEAS) and melatonin in breast and prostate cancer outpatients. *Psychoneuroendocrinology* 2004;29:448–74.
159. Sahajpal P, Ralte R. Impact of induced yogic relaxation training (IYRT) on stress-level, self-concept and quality of sleep among minority group individuals. *J Indian Psychol* 2000;18:66–73.
160. Walia JJ, Mehra P, Grover P, Earnest C, Verma SK, Sanjeev. Health status of nurses and yoga. IV. Experiment and results. *Nurs J India* 1992;83:27–8.
161. Shannahoff-Khalsa DS, Ray LE, Levine S, Gallen CC, Schwartz BJ, Sidorowich JJ. Randomized controlled trial of yogic meditation techniques for patients with obsessive compulsive disorders. *CNS Spectrums: Int J Neuropsychiat Med* 1999;4:34–46.
162. Berger B, Owen D. Mood alteration with yoga and swimming: aerobic exercise may not be necessary. *Percept Mot Skills* 1992;75:1331–43.
163. Ernst E, Rand J, Stevinson C. Complementary therapies for depression: an overview. *Arch Gen Psychiatry* 1998;55:1026–32.
164. Janakiramaiah N, Gangadhar B, Naga Venkatesha Murthy P, Harish M, Subbakrishna D, Vedamurthachar A. Antidepressant efficacy of Sudarshan Kriya Yoga (SKY) in melancholia: a randomized comparison with electroconvulsive therapy (ECT) and imipramine. *J Affect Disord* 2000;57:255–9.
165. Jorm A, Christensen H, Griffiths K, Rodgers B. Effectiveness of complementary and self-help treatments for depression. *Med J Aust* 2002;176 Suppl: S84–96.
166. Platania-Solazzo A, Field T, Blank J, et al. Relaxation therapy reduces anxiety in child and adolescent psychiatric patients. *Acta Paedopsychiatrica* 1992;55:115–20.
167. Netz Y, Lidor R. Mood alterations in mindful versus aerobic exercise modes. *J Psychol Res* 2003;137: 405–19.
168. Bhushan S, Sinha P. Yoganidra and management of anxiety and hostility. *J Indian Psychol* 2001;19: 44–9.
169. Allen KS, Steinkohl RP. Yoga in a geriatric mental clinic. *Act Adapt Aging* 1987;9:61–8.
170. Joseph CD. Psychological supportive therapy for cancer patients. *Indian J Cancer* 1983;20:268–70.
171. Cohen L, Warneke C, Fouladi RT, Rodriguez MA, Chaoul-Reich A. Psychological adjustment and sleep quality in a randomized trial of the effects of a Tibetan yoga intervention in patients with lymphoma. *Cancer* 2004;100:2253–60.
172. Vickers A, Goyal N, Harland R, Rees R. Do certain countries produce only positive results? A systematic review of controlled trials. *Control Clin Trials* 1998;19:159–66.