Use of Herbal Remedies by Hispanic Patients: Do They Inform Their Physician?

Laura Howell, MD, Komal Kochhar, MBBS, MHA, Robert Saywell, Jr., PhD, MPH, Terrell Zollinger, DrPH, Julie Koehler, PharmD, Christina Mandzuk, BSE, Brittany Sutton, MPH, Javier Sevilla-Martir, MD, and Deborah Allen, MD

Purpose: This study measured the knowledge and use of herbs among Hispanics and assessed their experiences when discussing herb use with their physician.

Methods: Self-administered questionnaires were collected from 620 Hispanic patients seeking treatment in urban health centers.

Results: Most (80.3%) reported using herbs. Herb users were more comfortable speaking Spanish (91.9% vs 80.2%) and had been in the United States less than 5 years (47.0% vs 29.4%). More users considered herbs as drugs (60.5% vs 39.6%). Users were more aware that herbs could harm a baby if taken during pregnancy (56.4% vs 36.0%). The majority did not know the English name for 23 of the 25 herbs. A majority indicated their physician was unaware of their herb use. Few (17.4%) responded that their physicians asked about herb use. Only 41.6% thought their physician would understand their herb use, and 1.8% believed their physician would encourage continued use. There were no significant differences between herb users and nonusers in their perception of patient-physician communication levels.

Conclusion: Primary care physicians need to be aware that most Hispanic patients are likely to use herbs. It is important to initiate and encourage discussion of their patient's interest in and use of these therapies. (J Am Board Fam Med 2006;19:566–78.)

The use of complementary and alternative medical therapies (CAM) has dramatically increased in the United States over the past 2 decades. The percentage of Americans using CAM increased from 33.8% to 42.1% from 1990 to 1997,^{1,2} and it is believed that the percentage has since increased. Most patients consider herbal remedies to be as safe and effective as patented medicines and continue to use various types of CAM.³

The sale of herbal medicines is growing by approximately 20% per year and is the largest growth area in retail pharmacy, far exceeding conventional medicine.⁴ Educated, young to middle-aged, non-

black, and financially well-off women are more likely to use CAM.^{1–3,5–8} Of those regularly taking prescription medicines, nearly 20% concurrently take at least one herb, a high-dose multivitamin, or both.^{2,9} Patients generally report using CAM to augment their conventional medical care rather than as a result of their dissatisfaction with mainstream medicine.^{5,10}

Herbal remedies are generally marketed in the United States as dietary supplements and are regulated as such under the 1994 Dietary Supplement Health and Education Act.⁹ In contrast to conventional drugs, herbal remedies do not undergo rigorous clinical trials and postapproval surveillance to define their effectiveness and relative safety. Consequently, clinical practitioners generally do not have available to them the scientific data needed to weigh the risks and benefits of herbal remedies as they do for conventional pharmaceutical medications.⁴ There is evidence to suggest that herbal remedies may contain ingredients that can worsen medical problems and interact with specific prescription medications.

Studies have estimated that the prevalence of herb use in the United States may be as high as one third^{1,2,5,6,11} and adverse reactions to herbs are

Submitted 20 March 2006; revised 9 June 2006; accepted 12 June 2006.

From the Department of Family Medicine, Indiana University School of Medicine, Indianapolis, IN (KK, RS, TZ, CM, BS, JS-M, DA); Cincinnati Children's Hospital, Cincinnati, OH (LH); and Butler University, Indianapolis, IN (JK).

Funding: This study was funded in part by the Marion County Health Department (Indianapolis, IN).

This study was presented as a poster session at the 2005 American Public Health Association, Annual Meeting and Exposition, Philadelphia, PA, on December 10–14, 2005.

Conflict of interest: none declared.

Corresponding author: Robert M. Saywell, Jr., Department of Family Medicine, Indiana University School of Medicine, 1110 West Michigan Street, Long Hospital 247, Indianapolis, IN 46202-5102 (E-mail: rsaywell@iupui.edu).

probably under-recognized and under-reported.^{4,12} Oral use of herbs constitutes a greater potential for significant health risk than non-ingested treatment modalities because many herbs may be therapeutic at a low dose but toxic at higher doses. In addition, interactions between herbs and drugs may increase or decrease the pharmacological or toxicological effects of either component in addition to the synergistic therapeutic effects that may complicate the dosing of long-term medications.¹³

Effective physician-patient communication is vital to reduce the dangers of herb-drug interactions. Physicians with an open mind to CAM may be more likely to hear from their patients about their herb use as studies show that as many as one half to two thirds of patients do not tell their physicians about their use of herbs.^{1,3} Knowledge and mutual respect are the foundations of negotiating conflicts that arise from approaching an illness from 2 different belief and value systems.¹⁴ Recent literature suggests the importance of providing training in CAM to medical students and residents and encouraging them to communicate these issues with patients.^{4,8,15} One study shows that physicians' exhibit better question-asking skills with non-Hispanic white patients, compared with Hispanic patients, and that patient ethnicity did not influence patient reported use or physician-patient communication about CAM.⁸ It was also observed that resident physicians with less clinical experience were more likely to ask patients one or more questions about their use of CAM.⁸

Even though herb use has increased over the past decade, the number of US pharmacy schools offering courses addressing their use has declined.¹⁵ According to one study, only 9 of 77 pharmacy colleges maintain pharmacognosy as a course in their curricula.^{15,16} As a result, today's pharmacy school graduates may not be adequately equipped to respond to their patients' request for information on herbs.¹⁵ Over one-half (52.0%) of the patients purchased their herbs in drugstores or groceries, where pharmacists would be available, but none of them received any herb information from the pharmacist.¹⁷ Another study found that more practicing pharmacists, compared with pharmacy students, felt that pharmacists should not become practitioners of CAM (26.6% vs 0.0%).¹⁸ This underscores the need for physicians to recognize an even greater responsibility to provide their patients with reliable information about herbs and

to be particularly cognizant of the need to do so for their Hispanic population.

Given the potential for adverse reactions and interactions associated with herbs, it is important to identify the prevalence of herb use in specific populations and whether herb users routinely inform their primary care physician of such use. Several studies have indicated that the Hispanic population, as an ethnic group, exhibits high usage of CAM.^{7,19} One study revealed that Mexican-American women were 3 times more likely to use herbs than the general population.¹⁹ Another study in New Mexico showed that 77% of surveyed Hispanics used herbs, compared with 47% of non-Hispanic whites.⁷ Specific reasons why Hispanics may be more likely to use herbs are not clearly understood.

Ferguson and Candib²⁰ concluded that minority patients, especially those not proficient in English, were less likely to establish rapport with physicians, less likely to receive sufficient medical information, and less likely to be encouraged to participate in medical decision making. Elder et al³ suggested that patients seek CAM because they cannot afford traditional medications. A distrust of physicians or physicians' negative responses to herb use has also been identified as a reason Hispanic patients' turn to herbs.⁵ Studies have also found that some patients believe that herb use, when combined with traditional therapy, may result in more optimal health outcomes.³

Numerous researchers^{14,17,21-28} have examined the unique cultural health beliefs of Hispanic patients, including the use of folk healing or herbs. However, few studies^{17,19,29,30} have actually surveyed Hispanic patients regarding their attitudes about herbs and their comfort level when discussing such remedies with their physicians. No studies have been conducted in Indiana, a Midwestern state with a rapidly growing Hispanic population. The goals of this study were to 1) assess Hispanic patients' knowledge about herbs, 2) explore the underlying reasons why these patients may be more likely to use herbs than the general population, 3) examine patients' comfort levels with discussing their herb use with their physicians, and 4) determine the extent to which physicians inquire about their patients' herb use.

Methods

A patient survey was conducted between October 2003 and February 2004 at 5 urban health centers

in Indianapolis, IN, that treat high numbers of Hispanic patients. The patients were approached in the waiting room of health clinics by bilingual research assistants. The research assistants identified the Hispanic patients visually by their ethnic appearance and asked each of them (if older than 18 years of age) to participate. At no time did the research assistants examine patients' medical records or ask medical staff the ethnicity of patients. It is not known how many patients refused to participate. All self-identified Hispanic patients over the age of 18, who visited the study clinic sites during the data collection period were asked by bilingual research assistants to complete a survey as they waited to see their physician. The survey instrument included questions about the patient's demographic information, familiarity with various specific herbs, attitudes toward and beliefs about herbs, and the patient's comfort level when discussing herb use with their physician. The survey instrument was initially created in English, and then translated into Spanish by an individual with a medical background whose first language was Spanish. It was then given to another medically experienced individual fluent in Spanish who translated it into English to ensure concept equivalency. The Spanish version of the survey instrument was also examined by numerous members of the Indianapolis Hispanic community to ensure that questions were asked appropriately and with cultural sensitivity. Research assistants interviewed patients if they were unable to read the survey instrument. Survey completion was voluntary, no incentives were provided, and the subjects were assured that their responses would be kept confidential. The study received Indiana University Institutional Review Board approval in May 2003.

Completed questionnaires were scanned into an electronic data file, verified by a research assistant and analyzed using SPSS version 12.0.1. Descriptive statistics including mean, median, and frequency distribution were calculated. χ^2 tests were performed to determine significant statistical differences. Values of *P* less than .05 were considered statistically significant.

Results

Demographic and Clinical Characteristics

A total of 620 Hispanic individuals completed the herb use questionnaire. The herbs used in the sur-

vey instrument were chosen by a consensus of numerous Hispanic individuals interviewed in the community as to what were the most common herbs used. Most often, the respondents were female (78.7%), unemployed (56.8%), uninsured (69.7%) and had lived in the United States for less than 5 years (43.9%). One half (58.9%) of the respondents were in their 20s, whereas approximately one tenth (12.2%) were over age 40. More than one half (53.0%) did not have a high school degree, and nearly one half (48.8%) lived with their spouse. More than one fourth (25.4%) had at least one child living in their household with a mean household size of 4.16. More than four fifths (89.2% and 89.9%, respectively) were most comfortable reading and speaking Spanish rather than English. Less than one third (29.8%) had not seen a physician in the past year, whereas more than one third (35.5%) had seen a physician at least 3 times in the past year. A majority (90.1%) had not seen a *curandero* or *hierbero* in the past year.

Use of Herbal Remedies

Four fifths (80.3%) responded that they were currently taking or had taken an oral herbal remedy in the past. Two thirds (59.6%) had taken between 6 and 15 different herbs during their lifetime. One third (30.7%) indicated they obtained their herbs from a Hispanic grocery store. Some indicated that herbs were most often used to treat cough (26.3%), stomach pain (24.9%), sore throat (20.3%), menstrual cramps (19.1%), headache (8.8%), and chest pain (7.2%).

Participants were asked about their use and knowledge of the English names for 25 commonly used herbs for treatment purposes (not as food seasonings) and whether their physicians were aware of their use, as shown in Table 1. More than half of the respondents reported that they had used at least 1 of 10 herbs among the 25 listed. The 10 most commonly used herbs were cinnamon, cloves, cumin, chamomile, garlic, onion, grass syrup, aloe vera, oregano, and lemon. The majority of respondents did not know the English name for 23 of the 25 herbs on the list. The 2 herbs for which the majority knew the English name were spider milkweed and bitter melon. A vast majority of the respondents indicated that their doctors were unaware of their herb use.

Table 2 presents the demographic characteristics of the herb users and non-users. Of the 8

Table 1. Use and Physician Knowledge of My Herbal Use

	Have Used the Herb		Know English Name		Doctor Aware I Use	
	Number	%	Number	%	Number	%
Immortal (Spider Milkweed)	7	1.4	5	71.4	3	42.9
Plumajillo (Pleurisy Root)	16	3.2	6	37.5	5	31.3
Canela (Cinnamon)	415	83.3	81	19.5	22	5.3
Clavo (Cloves)	292	58.6	27	9.2	8	2.7
Comino (Cumin)	295	59.2	27	9.2	14	4.7
Eucalipto (Eucalyptus)	214	43.0	30	14.0	20	9.3
Manzanilla (Chamomile)	421	84.5	54	12.8	45	10.7
Ajo (Garlic)	368	73.9	65	17.7	22	6.0
Jengibre (Ginger)	116	23.3	38	32.8	16	13.8
Cebolla (Onion)	352	70.7	88	25.0	19	5.4
Granada (Pomegranate)	165	33.1	23	13.9	15	9.1
Yerba Buena (Grass Syrup)	353	70.9	38	10.8	24	6.8
Anis Estrella (Star Anise)	131	26.3	22	16.8	17	13.0
Estafiate (Wormwood)	83	16.7	12	14.5	10	12.0
Siete Jarabes (Seven Syrups)	29	5.8	13	44.8	11	37.9
Cundeamore (Bitter Gourd)	14	2.8	11	78.6	6	42.9
Savila (Aloe Vera)	261	52.4	34	13.0	18	6.9
Una de Gato (Cat's Claw)	62	12.4	15	24.2	12	19.4
Pelos de Elote (Corn Silk)	140	28.1	21	15.0	16	11.4
Tlanchalagua (Tea)	31	6.2	15	48.4	11	35.5
Oregano	322	64.7	41	12.7	22	6.8
Limon (Lemon)	378	75.9	82	21.7	22	5.8
Valeriana (Valerian)	66	13.3	19	28.8	10	15.2
Tomillo (Thyme)	137	27.5	12	8.8	9	6.6
Epasote (Wormseed)	220	44.2	8	3.6	6	2.7

* Number using the herb/total number of herb users $[7/498 \cdot 100 = 1.4]$.

** Number knowing the English name/number using the herb $[5/7 \cdot 100 = 71.4]$.

*** Number indicating Doctor is aware/number using the herb $[3/7 \cdot 100 = 42.9]$.

variables included in the analysis, only 2—language and duration of residence in the United States—were significantly different between the 2 groups. Herb users were more comfortable speaking Spanish than English (91.9% vs 80.2%) and more often lived in the United States for less than 5 years (47.0% vs 29.4%). The remaining 6 characteristics: gender, age, education level, employment status, health insurance coverage, and number of physician visits in the past year were not significantly different between the herb users and nonusers. Two multiple logistic regression models were developed to determine the extent to which demographic characteristics explain the use of herbal remedies and whether or not the individual informed their physician of their herb use. None of the findings were found to be statistically significant.

Knowledge and Use of Herbs

Table 3 shows that the overall knowledge about herbs was lacking among the participants. More than half of the respondents felt that herbs were drugs (58.2%) and that some herbs could harm a baby if taken during pregnancy (54.3%). Herb users were significantly more likely than nonusers to know that herbs are drugs (60.5% vs 39.6%) and that herbs could harm a baby if taken during pregnancy (56.4% vs 36.0%). One third believed that some herbs could interact with prescription medications (35.1%) and that herbs were not safe to use during pregnancy, compared with prescription or over-the-counter drugs (35.3%).

More than two thirds (66.0%) indicated that a family member recommended they take herbs. Four in 10 (44.0%) indicated they had given their children or grandchildren herbs. Herb users were

Table 2. Demographic Characteristics of Herb Users and Nonusers

	Yes (N = 498)		No (N = 122)		P Value
	Number	%	Number	%	
Do you currently take or have you ever taken an oral herbal remedy?	498	80.3	122	19.7	
Which language do you feel most comfortable speaking?					.001
Spanish	441	91.9	77	80.2	
English	15	3.1	10	10.4	
Both	24	5	9	9.4	
Missing	18		26		
What is your gender?					.375
Male	99	20.6	26	24.5	
Female	381	79.4	80	75.5	
Missing	18			16.0	
What is your age?					.477
0 to 20	52	10.5	12	10.6	
21 to 24	130	26.4	28	24.8	
25 to 29	113	23.1	22	19.4	
30 to 34	90	18.2	17	15.0	
35 to 39	38	7.6	15	13.4	
≥ 40	70	14.2	19	16.8	
Missing	5		9		
What is the highest level of education you completed?					.269
Less than High School	218	52.7	46	54.8	
High School	153	37.0	34	40.5	
College or above	43	10.4	4	4.8	
Missing	84		38		
Are you currently working?					.343
Yes	203	42.3	47	47.5	
No	277	57.7	52	52.5	
Missing			23		
Do you have health insurance?					.760
Yes	141	30.0	30	31.6	
No	329	70.0	65	68.4	
Missing			27		
How many times have you seen a medical doctor in the past year?					.534
None	128	29.8	23	29.5	
One to two times	145	33.8	31	39.7	
Three or more times	156	36.4	24	30.8	
Missing	69		44		
How long have you lived in the United States?					.000
Less than 5 years	207	47.0	27	29.4	
5 years or more	234	53.0	65	70.6	
Missing	57		30		

significantly more likely than nonusers to reveal that a family member recommended that they take herbs (69.7% vs 39.7%) and that they provide herbs to their children or grandchildren (46.0% vs 23.1%).

Physician Communication and Perceived Attitudes

Approximately three fourths (72.7%) indicated that they were able to communicate easily with their

doctors, as shown in Table 4. However, nearly three fourths (73.7%) of the respondents reported that their doctors did not ask them about their use of herbs, and a majority (90.2%) indicated that their doctor never recommended they take herbs. Majority felt their physician took time to listen to them at each appointment (83.8%); understood their cultural background

Table 3. Knowledge/Behavior about Herbal Remedies

	Users (N = 498)		Non Users (N = 122)		Total (N = 620)		P Value
	Number	%	Number	%	Number	%	
Herbal remedies are drugs							.012
No	65	15.1	14	26.4	79	16.4	
Yes	260	60.5	21	39.6	281	58.2	
Do not know	105	24.4	18	34.0	123	25.5	
Missing	68		69		137		
Some herbal remedies can interact with prescription medications							.327
No	74	17.1	9	17.0	83	17.0	
Yes	157	36.2	14	26.4	171	35.1	
Do not know	203	46.8	30	56.6	233	47.8	
Missing	64		69		133		
Some herbal remedies can harm a baby if taken during pregnancy							.016
No	37	8.4	8	16.0	45	9.2	
Yes	247	56.4	18	36.0	265	54.3	
Do not know	154	35.2	24	48.0	178	36.5	
Missing	60		72		132		
Herbal remedies are safer to use during pregnancy than prescription or over-the-counter drugs							.917
No	154	35.5	17	33.3	171	35.3	
Yes	84	19.4	11	21.6	95	19.6	
Do not know	196	45.2	23	45.1	219	45.2	
Missing	64		71		135		
My family member has recommended an herbal remedy for me to take							.000
No	122	29.6	33	56.9	155	33.0	
Yes	287	69.7	23	39.7	310	66.0	
Do not know	3	0.7	2	3.4	5	1.1	
Missing	86		64		150		
I have given my children (or grandchildren) herbal remedies							.004
No	194	50.5	37	71.2	231	53.0	
Yes	180	46.0	12	23.1	192	44.0	
Do not know	10	2.6	3	5.8	13	3.0	
Missing	114		70		184		

(58.9%); and worked with them to improve their health (74.1%). More than two thirds (71.6%) indicated that if they went to a *curandero* for treatment, they would tell their doctor about it. Three fourths (77.6%) felt that “they can tell their doctor anything.” There were no significant differences noted between the user and nonuser groups.

When asked what they tell their doctors about their use of herbs, one sixth (15.0%) responded that “I would tell my doctor about all the remedies that I use.” When asked what would happen if they told their doctors that they use herbs, only 4 in 10

(41.8%) indicated that their doctor would understand their herb use. More than one-quarter (25.9%) responded that their doctors would not know what the herb was used for and 20.5% thought that their doctor would tell them to stop using all herbs. One in 10 (10.5%) believed that their doctor would encourage them to continue using herbs; few (1.4%) thought their doctors would ridicule them for using herbs.

Some of the more commonly used herbs and their potential drug interactions are presented in Table 5.^{32–34}

Table 4. Physician Communication

	Users (N = 498)		Non Users (N = 122)		Total (N = 620)		P Value
	Number	%	Number	%	Number	%	
I am able to communicate easily with my doctor							.573
No	76	20.2	10	19.2	86	20.1	
Yes	271	72.1	40	76.9	311	72.7	
Do not know	29	7.7	2	3.8	31	7.2	
Missing	122		70		192		
I feel that I can tell my doctor anything							.995
No	52	13.4	7	13.2	59	13.4	
Yes	301	77.6	41	77.4	342	77.6	
Do not know	35	9.0	5	9.4	40	9.1	
Missing	110		69		179		
My doctor asks me whether I am using any herbal remedies							.081
No	310	74.3	35	68.6	345	73.7	
Yes	73	17.5	7	13.7	80	17.1	
Do not know	34	8.2	9	17.6	43	9.2	
Missing	81		71		152		
My doctor has recommended an herbal remedy for me to take							.631
No	362	89.8	54	93.1	416	90.2	
Yes	24	6.0	3	5.2	27	5.9	
Do not know	17	4.2	1	1.7	18	3.9	
Missing	95		64		159		
My doctor takes time to listen to me at each appointment							.823
No	36	9.0	5	9.3	41	9.0	
Yes	338	84.1	44	81.5	382	83.8	
Do not know	28	7.0	5	9.3	33	7.2	
Missing	96		68		164		
My doctor understands my cultural background							.243
No	42	11.7	9	19.1	51	12.6	
Yes	211	58.8	28	59.6	239	58.9	
Do not know	106	29.5	10	21.3	116	28.6	
Missing	139		75		214	52.7	
My doctor works with me (and my family) to improve my health							.525
No	52	14.2	9	19.6	61	14.8	
Yes	275	74.9	31	67.4	306	74.1	
Do not know	40	10.9	6	13.0	46	11.1	
Missing	131		76		207		
If I went to a curandero for treatment, I would tell my doctor about it							.531
No	44	11.6	9	17.0	53	12.2	
Yes	274	72.1	36	67.9	310	71.6	
Do not know	62	16.3	8	15.1	70	16.2	
Missing	118		69		187		
							Number %
Only those who “use” or “ever used” herbal remedies, were asked the following questions:							
What do you tell your doctor about your use of herbal remedies?							
I would tell my doctor about all the remedies that I use							37 15.0
I would tell my doctor about some of the remedies that I use							46 18.7
My doctor never asks whether I use any herbs							163 66.3
Missing							252
What would happen if you told your doctor that you use herbal remedies?							
My doctor would understand the use of these remedies							92 41.8
My doctor would not know what these herbal remedies are used for							57 25.9
My doctor would tell me to stop using all herbal remedies							45 20.5
My doctor would encourage me to keep using the remedies that are safe							23 10.5
My doctor would ridicule me							3 1.4
Missing							278

Table 5. Examples of Selected Herb Use and Drug Interactions

Herb	Use	Potential Interactions with Prescription Medications
Immortal (Spider Milkweed)	Orally, for arthritis, asthma, cough, edema, syphilis, valvular insufficiency, senile heart, to strengthen weak heart muscles following pneumonia, and for diuresis. Topically, for warts.	Digoxin, diuretics
Plumajillo (Pleurisy Root)	Orally, for cough, pleurisy, uterine disorders, shortness of breath, pain, spasms, and to promote sweating; for bronchitis, pneumonitis, and influenza.	Digoxin, diuretics, estrogen
Canella (Cinnamon)	Orally, for colds, poor circulation, and as a bitter tonic. In foods, as a cooking spice.	None known
Clavo (Cloves)	Orally, clove for dyspepsia, as an expectorant, for diarrhea, hernia, halitosis, flatulence, nausea, and vomiting. Topically, for toothache, postextraction alveolitis, pain, a dental anesthetic, mouth and throat inflammation. In combination with other ingredients, topically for premature ejaculation. In foods and beverages, as a flavoring agent. In manufacturing, in toothpaste, soaps, cosmetics, and perfumes.	Antiplatelet and anticoagulant agents
Comino (Cumin)	Orally, as an antispasmodic, stimulant, antispasmodic, diuretic, aphrodisiac, for stimulating menstrual flow, treating diarrhea, colic, and flatulence. In foods, spices, and beverages, as a flavoring agent. In manufacturing, as a cosmetic fragrance.	Antidiabetic agents
Manzanilla (Chamomile)	Orally, for flatulence, travel sickness, nasal mucous membrane inflammation, allergic rhinitis, nervous diarrhea, attention deficit-hyperactivity disorder, restlessness, insomnia, gastrointestinal spasms, inflammatory diseases of the gastrointestinal tract, gastrointestinal ulcers associated with NSAIDs and alcohol consumption, and as an antispasmodic for menstrual cramps. Topically, for hemorrhoids, leg ulcers; for skin, anogenital, and mucous membrane inflammation; for bacterial skin diseases; for treating or preventing chemotherapy- or radiation-induced oral mucositis. Via inhalation, for inflammation and irritation of the respiratory tract. In foods and beverages, as a flavoring agent. In manufacturing, in cosmetics, soaps, and mouthwashes.	Benzodiazepines; CNS depressants (eg, fentanyl, morphine, etc.); contraceptive drugs; cytochrome P450 1a2 (Cyp1a2) substrates (eg, amitriptyline, haloperidol, verapamil, propranolol, theophylline, ondansetron, etc.); cytochrome P450 3a4 (Cyp3a4) substrates (eg, lovastatin, itraconazole, ketoconazole, fexofenadine, triazolam, etc.); estrogens; tamoxifen; warfarin
Ajo (Garlic)	Orally, for hypertension, hyperlipidemia; for prevention of coronary heart disease, age-related vascular changes and atherosclerosis; for reducing reinfarction and mortality rate post-myocardial infarction, earaches, chronic fatigue syndrome, and menstrual disorders; for HIV-drug induced lipid disorders and <i>Helicobacter pylori</i> infection; for prevention of colorectal, gastric, breast, lung, and prostate cancer; for bladder cancer, benign prostatic hyperplasia, diabetes, arthritis, allergies, traveler's diarrhea, colds and flu; for immune system stimulation, prevention of tick bites, and prevention and treatment of bacterial and fungal infections; for diarrhea, amoebic and bacterial dysentery, tuberculosis, bloody urine, diphtheria, whooping cough, scalp ringworm, hypersensitive teeth, and vaginal trichomoniasis; for fever, cough, headache, stomachache, sinus congestion, athlete's foot, gout, rheumatism, hemorrhoids, asthma, bronchitis, shortness of breath, arteriosclerosis, low blood pressure, hypoglycemia, hyperglycemia, and snakebites; as a diuretic, stimulant, cathartic, aphrodisiac; for enhancing circulation, fighting stress and fatigue.	Anticoagulant and antiplatelet agents; contraceptives; cyclosporine; cytochrome P450 2e1 (Cyp2e1) substrates (eg, acetaminophen, ethanol, theophylline, etc.); cytochrome P450 3a4 (Cyp3a4) substrates (eg, lovastatin, itraconazole, ketoconazole, fexofenadine, triazolam, etc.); non-nucleoside reverse transcriptase inhibitors; saquinavir

(Table continues)

Table 5. Continued

Herb	Use	Potential Interactions with Prescription Medications
Jengibre (Ginger)	Orally, for motion sickness, morning sickness, colic, dyspepsia, flatulence, chemotherapy-induced nausea, rheumatoid arthritis, osteoarthritis, loss of appetite, post-operative nausea and vomiting, migraine headache, anorexia, upper respiratory tract infections, cough, bronchitis, and as a diaphoretic, diuretic, and a stimulant. Fresh ginger: Orally, for treating acute bacterial dysentery, baldness, malaria, orchitis, poisonous snake bites, rheumatism, and toothaches. Dried ginger: orally, for chest pain, low back pain, and stomach pain. Topically, for thermal burns and as an analgesic. In foods and beverages, as a flavoring agent. In manufacturing, as a fragrance component in soaps and cosmetics.	Antiplatelet and anticoagulant agents; antidiabetic agents; calcium channel blockers
Granada (Pomegranate)	Orally, for hypertension, heart failure, myocardial ischemia, atherosclerosis, hyperlipidemia, acidosis, hemorrhage, HIV disease, tapeworm infestations, diarrhea, dysentery, and opportunistic intestinal worms; for preventing prostate cancer, and as an astringent and abortifacient; for chronic obstructive pulmonary disease, influenza, stomatitis, periodontal disease, erectile dysfunction, diabetes, and cancer. Topically, as a gargle for sore throat and to treat hemorrhoids.	Antiplatelet and anticoagulant agents; antidiabetic agents; ACE inhibitors; antihypertensive agents; cytochrome P450 3a4 (Cyp3a4) substrates (eg., lovastatin, itraconazole, ketoconazole, fexofenadine, triazolam, etc.); cytochrome P450 2D6 (CYP2D6) substrates (eg, amitriptyline, fluoxetine, codeine, tramadol, ondansetron, etc.)
Anis Estrella (Star Anise)	Orally, for respiratory infections and inflammation, influenza, avian flu, gastrointestinal upset, flatulence, loss of appetite, infant colic, cough, and bronchitis; for increasing milk secretion, promoting menstruation, facilitating childbirth, increasing libido, and treating symptoms of male climacteric. Through inhalation, for respiratory tract congestion. In foods and beverages, as a culinary spice and flavoring agent. In manufacturing, as a fragrance component in soaps, cosmetics, perfumes, and toothpaste, and to mask undesirable odors in drug products.	None known
Estafiate (Wormwood)	Orally, for loss of appetite, indigestion and digestive disorders, biliary dyskinesia, fever, and liver disease; as an anthelmintic, aphrodisiac, tonic, antispasmodic, and to stimulate sweating and the imagination. Topically, for healing wounds and insect bites and as a counterirritant. In foods, wormwood as a flavoring agent. In manufacturing, as a fragrance component in soaps, cosmetics, and perfumes, and as an insecticide.	Anticonvulsants
Cundeamore (Bitter Gourd)	Orally, for diabetes, psoriasis, gastrointestinal upset, ulcers, colitis, constipation, intestinal worms, kidney stones, fever, hepatic disease, and to induce menstruation. Topically, for skin abscesses and wounds, and anorectal herpes lesions.	Antidiabetic agents
Savila (Aloe Vera)	Orally, for osteoarthritis, inflammatory bowel diseases, fever, itching and inflammation, as a general tonic, for gastroduodenal ulcers, diabetes, asthma, and radiation-related mucositis. Topically, for burns, wound healing, psoriasis, sunburn, frostbite, inflammation, osteoarthritis, and cold sores; and as an antiseptic and a moisturizer.	Oral medications (in general); sevoflurane; diuretics; digoxin; antidiabetic agents

(Table continues)

Table 5. Continued

Herb	Use	Potential Interactions with Prescription Medications
Una de Gato (Cat's Claw)	Orally, for diverticulitis, peptic ulcers, colitis, gastritis, hemorrhoids, parasites, Alzheimer's disease, chronic fatigue syndrome, wound healing, arthritis, asthma, allergic rhinitis, cancer (especially of the urinary tract), glioblastoma, gonorrhea, dysentery, birth control, bone pain, "cleansing" the kidneys, and viral infections, including herpes zoster, herpes simplex, and human immunodeficiency virus (HIV).	Antihypertensive agents; immunosuppressants; cytochrome P450 3a4 (Cyp3a4) substrates (eg, lovastatin, itraconazole, ketoconazole, fexofenadine, triazolam, etc.)
Pelos de Elote (Corn Silk)	Orally, for cystitis, urethritis, nocturnal enuresis, prostatitis, inflammation of the urinary tract, diabetes, hypertension, and as a diuretic for congestive heart failure.	Antihypertensive agents; antidiabetic agents; diuretics; corticosteroids; warfarin
Oregano	Orally, for respiratory tract disorders, including cough, asthma, croup, and bronchitis; for gastrointestinal disorders, such as dyspepsia and bloating; for dysmenorrhea, rheumatoid arthritis, urinary tract infections, headaches, heart conditions, intestinal parasites, allergies, sinusitis, arthritis, cold and flu, earaches, and fatigue. Topically, for acne, athlete's foot, dandruff, insect and spider bites, canker sores, gum disease, toothaches, psoriasis, seborrhea, ringworm, rosacea, muscle pain, varicose veins, and warts, and as an insect repellent. In foods and beverages, as a culinary spice and a preservative.	None known
Limon (Lemon)	Orally, as a source of vitamin C in the treatment of scurvy and colds; as a digestive aid, an anti-inflammatory, diuretic, and to improve vascular permeability. In foods, as a food and flavoring agent.	None known
Valeriana (Valerian)	Orally, as a sedative-hypnotic for insomnia and as an anxiolytic for restlessness; for mood disorders such as depression, mild tremors, epilepsy, attention deficit-hyperactivity disorder, and chronic fatigue syndrome; for muscle and joint pain, asthma, hysterical states, excitability, hypochondria, headaches, migraine, stomach upset, menstrual cramps and symptoms associated with menopause, including hot flashes and anxiety. Topically, as a bath additive for restlessness and sleep disorders. In manufacturing, as flavoring in foods and beverages.	Alcohol; benzodiazepines; CNS depressants; cytochrome P450 3a4 (Cyp3a4) substrates (eg, lovastatin, itraconazole, ketoconazole, fexofenadine, triazolam, etc.)
Tomillo (Thyme)	Orally, for bronchitis, pertussis, sore throat, colic, arthritis, dyspepsia, gastritis, diarrhea, enuresis, dyspraxia, flatulence, skin disorders, as a diuretic, urinary disinfectant, anthelmintic, and as an appetite stimulant. Topically, for laryngitis, tonsillitis, stomatitis, and halitosis; as a counterirritant, an antiseptic in mouthwashes and liniments, and for alopecia areata. Orally, as an antibacterial and antifungal ingredient. In foods, as a flavoring agent. In manufacturing, in perfumes, soaps, cosmetics, and toothpastes.	Antiplatelet and anticoagulant agents
Epasote (Wormseed)	Orally, for ascaris and oxyuris infestations.	None known

Discussion

The majority of Hispanics who participated in the survey were herb users and it seems that their physicians were generally unaware of their herb use. The herb users tended to be more comfortable

speaking Spanish rather than English and had been in the United States for less than 5 years indicating that they may have problems communicating with their physician and/or may not have a good understanding of the health care system in the United

States. The fact that many did not know the English term for the herbs they use is further evidence of the existence of a barrier to achieving appropriate communications with their physician. Gordon¹⁴ found that newly arrived Hispanics who were primarily from rural areas were more likely to hold traditional beliefs and values, whereas US-born Hispanics were likely to have more assimilated attitudes, which may explain the higher rate of herb use in recent immigrants.

Elder et al³ pointed out that the main reason given for using herbs, alone or in combination with prescribed medications, was a belief that herbs would be effective. Many of those who communicated with their physician spoke of acceptance and control, but those who did not communicate with their physician mentioned traditional medicine's limitations and the narrow-mindedness of their physicians.³ Patients who chose to reveal details about their herb use did so because they perceived their physician would be respectful, open-minded, and willing to listen to them.³¹ Patients found it easier to discuss their use of CAM when they believed that their physician expected them to be using herbs.³¹ On the other hand, patients who chose not to reveal their herb use with their physician gave several reasons for their decision, such as the impression that their physician would be disinterested in their herb use; the anticipation of a negative response; the conviction that their physician would be unwilling or unable to contribute useful information; and the impression that their physician perceives herbs to be irrelevant in the biomedical treatment course.³¹ In addition, patients may feel intimidated by their physicians and perceive a sense of disapproval with regard to their physician's views on their use of CAM.³⁵ Similar concerns were expressed by participants in this study. The majority of respondents noted that their physician would not know what the herbs were used for, and one fourth felt that their physician would ridicule them for using herbs.

Eisenberg et al¹ noted that 70% of patients do not reveal their herb use to their allopathic practitioners. The current study indicated that for some herbs, this percentage may be much higher, as only 15% indicated that they would tell their doctor about all the herbs they use whereas three fourths replied that they could "tell their doctor anything." Providing herb use information to one's physician was not considered important to the patient; only

one third of the respondents were aware that herbs could interact with prescription medication. Regardless of whether patients plan to inform their provider about herb use or not, the onus is on providers to ask the question of their patients.³⁶

This study presents the findings of a survey and is subject to the limitations of self-reported data. Most of the survey items were value neutral. However, undoubtedly the respondents would want to appear to be knowledgeable and may have misstated their actual herb use. In addition, because this survey was administered in health care clinics, the responses to items relating to physician communication may have over-represented the level of communication between providers and patients.

Approximately one fourth of the participants who answered the first few pages of the survey did not complete the last part. There are at least 2 plausible reasons for the dropoff in response. The respondents may have tired of answering the questions and quit. Secondly, some questions in the last part dealt with their perspective of the physician's attitude about herb use; thus, the respondents may not have wanted to answer these questions for fear that their responses may become known to their physician. Consequently, the proportion who felt that their physician will not support their herb use may actually be higher. Because this study surveyed a convenience sample of Hispanics, the responses of these subjects may not be representative of the general population of Hispanics in Indiana. Those who attend health clinics may be less likely to use herbs, thus the estimates of prevalence in this study may be conservative.

Primary care practitioners need to understand the extent and patterns of herbal use by their multiethnic patients and efforts to elicit information from patients about herbal use maybe warranted.³⁷ It is important to initiate and encourage open, honest discussion about their patients' interest in or use of herbs. Assessments should start with a thorough drug history, which should include inquiry into the use of dietary supplements including herbs. Pharmacists should strive to provide information routinely on the potential intrinsic effects and interactions of herbs with prescribed medications. Therefore, more continuing education programs and drug information resources about herbal medications must be made available to pharmacists as consumer use continues to escalate.³⁸ In addition, therapy guidelines should be clearly defined as in-

creased knowledge of these adverse factors can help design safer pharmacologic and herbal regimens for individual patients, thus minimizing adverse reactions and promoting good health. Lastly, physicians and consumers are encouraged to report suspected adverse effects of herbal products through the Food and Drug Administration Medwatch system (www.fda.gov or 1-800-FDA-1088)^{4,9} and available at <http://www.cfsan.fda.gov/~dms/supplmnt.html>.

It is an obligation of clinicians to elicit information on the use of herbs that may influence their patients' health and to provide information on safe and effective treatment options. It is paramount for clinicians to be aware of known or potential herb-drug interactions to adequately treat their patients. Both physicians and consumers must become more educated about the safe and effective use of herbs. Asking patients about supplement use during an initial history should be made a central component of patient care and medication use monitoring.³⁵

References

- Eisenberg DM, Kessler RC, Foster C, et al. Unconventional medicine in the United States. *N Engl J Med* 1993;328:246–52.
- Eisenberg DM, Davis RB, Ettner SL, et al. Trends in alternative medicine use in the United States, 1990–1997. *JAMA* 1998;280:1569–75.
- Elder NC, Gillcrist A, Minz R. Use of alternative health care by family practice patients. *Arch Fam Med* 1997;6:181–4.
- Winslow LC, Kroll DJ. Herbs as medicines. *Arch Intern Med* 1998;158:2192–9.
- Astin JA. Why patients use alternative medicine: results of a national study. *JAMA* 1998;279:1548–53.
- Ni H, Simile C, Hardy AM. Utilization of complementary and alternative medicine by United States adults. *Med Care* 2001;40:353–8.
- Dole EJ, Rhyne RL, Zeilmann CA, et al. The influence of ethnicity on use of herbal remedies in elderly Hispanics and non-Hispanic whites. *J Am Pharm Assoc* 2000;40:359–65.
- Sleath B, Rubin RH, Campbell W, et al. Ethnicity and physician-older patient communication about alternative therapies. *Altern Complement Med* 2001;7:329–35.
- Boullata JI, Nace AM. Safety issues with herbal medicine. *Pharmacol Ther* 2000;20:257–69.
- Druss BG, Rosenheck RA. Association between use of unconventional therapies and conventional medical services. *JAMA* 1999;282:651–6.
- O'Hara M, Kiefer M, Farrell K, Kemper K. A review of 12 commonly used medicinal herbs. *Arch Fam Med* 1998;7:523–36.
- D'Arcy P. Adverse reactions and interactions with herbal medicines, I: adverse reactions. *Adv Drug React Toxicol Rev* 1991;10:189–208.
- Fugh-Berman A. Herb-drug interactions. *Lancet* 2000;355:134–8.
- Gordon SM. Hispanic cultural health beliefs and folk remedies. *J Holist Nurs* 1994;12:307–22.
- Klepser TB, Klepser ME. Unsafe and potentially safe herbal therapies. *Am J Health-Syst Pharm* 1999;56:125–38.
- Miller LG, Murray WJ. Herbal instruction in United States pharmacy schools. *Am J Pharm Educ* 1997;61:160–2.
- Zeilmann CA, Dole EJ, Skipper BJ, et al. Use of herbal medicine by elderly Hispanic and non-Hispanic white patients. *Pharmacotherapy* 2003;23:526–32.
- Hamilton WR, Monaghan MS, Turner PD. A comparison of pharmacy practitioner and pharmacy student attitudes toward complementary and alternative therapies in a rural state. *Am J Pharm Educ* 2002;66:55–8.
- Zenk SN, Shaver JLF, Peragallo N, et al. Use of herbal therapies among midlife Mexican women. *Health Care for Women Inter* 2001;22:585–97.
- Ferguson WJ, Candib LM. Culture, language, and the doctor-patient relationship. *Fam Med* 2002;34:353–61.
- Murguia A, Zea MC, Reisen CA, Peterson RA. The development of the cultural health attributions questionnaire (CHAQ). *Cultur Divers Ethnic Minor Psychol* 2000;6:268–83.
- Lipton RB, Losey LM, Giachello A, et al. Attitudes and issues in treating Latino patients with type 2 diabetes: views of healthcare providers. *Diabetes Educ* 1998;24:67–71.
- Chesney AP, Thompson BL, Guevara A, Vela A, Schottstaedt MF. Mexican-American folk medicine: implications for the family physician. *J Fam Pract* 1980;11:567–74.
- Larkey LK, Hecht ML, Miller K, Alatorre, C. Hispanic cultural norms for health-seeking behaviors in the face of symptoms. *Health Educ and Behav* 2001;28:65–80.
- Murguia A, Peterson RA, Zea MC. Use and implications of ethnomedical health care approaches among Central American immigrants. *Health and Soc Work* 2003;28:43–51.
- Mendoza M, Peterson MC. New Latino immigration to Tennessee: practicing culturally sensitive health care. *Tenn Med* 2000;10:371–6.
- Borrayo EA, Jenkins SR. Feeling frugal: socioeconomic status, acculturation, and cultural health beliefs among women of Mexican descent. *Cultur Divers Ethnic Minor Psychol* 2003;9:197–206.

28. Mendelson C. Health perceptions of Mexican American women. *J Transcult Nurs* 2002;13:210–17.
29. Davis RE. Understanding women's experiences with pharmacopeia. *Health Care for Women Inter* 2002; 18:210–17.
30. Bharucha DX, Morling BA, Niesenbaum RA. Use and definition of herbal medicines differ by ethnicity. *Ann Pharmacother* 2003;37:1409–13.
31. Adler SR, Fosket JR. Disclosing complementary and alternative medicine use in the medical encounter: a qualitative study in women with breast cancer. *J Fam Pract* 1999;48:453–8.
32. Lee A, Chui PT, Aun CST, Gin T, Lau ASC. Possible interaction between sevoflurane and aloe vera. *Ann Pharmacother* 2004;38:1651–4.
33. Natural Medicines Comprehensive Database. Available at [http://ezproxy.butler.edu:2124/\(bofk055gbk2aybln2fyj0y1\)/home.aspx?li=1&st=2&cs=mlatrial&s=ND](http://ezproxy.butler.edu:2124/(bofk055gbk2aybln2fyj0y1)/home.aspx?li=1&st=2&cs=mlatrial&s=ND).
34. Methodist Medical Group Pharmacies-A Clarian Health Partner. Herbal Products and Supplements-Quick Reference Guide. Indianapolis, IN. July 2005.
35. Cohen KR, Cerone P, Ruggiero R. Complementary/Alternative Medicine Use: Responsibilities and implications for pharmacy services. *P&T* 2002;27: 440–6.
36. Planta M, Gundersen B, Pettitt JC. Prevalence of the use of herbal products in a low-income population. *Fam Med* 2000;32:252–7.
37. Quo GM, Hawley ST, Weiss LD, Balkrishnan R, Volk RJ. Factors associated with herbal use among urban multiethnic primary care patients: A cross-sectional survey. *BMC Complement Alt Med* 2004; 12:4–18.
38. Chang ZG, Kennedy DT, Holdford DA, Small RE. Pharmacists' knowledge and attitude toward herbal medicine. *Ann Pharmacother* 2000;34:710–5.