world, limited resource, high-disparity primary care practice settings” called for in the article.

CBPR and cooperative grants through the National Cancer Institute (NCI) combine principle and practice to yield significant improvements in access to care and standard of care protocols for cancer patients. NCI’s Cancer Disparities Research Partnership (CDRP) (www3.cancer.gov/rrp/CDRP/index.html) links community cancer centers with comprehensive cancer centers to create the triangulation-based care by providers, patient, and community that Rust and Cooper emphasize. The CDRP program was initiated in response to an Institute of Medicine Report in 1999 outlining significant discrepancies between the level of care received in community hospitals versus large academic medical centers.

Current CDRP sites include Rapid City, SD; Wilmington, NC; Inglewood, CA; McKeepsport, PA; and Pascagoula, MS. The program in Rapid City, SD (*Walking Forward*), addresses cancer disparities for nearly 60,000 American Indians (AIs) who suffer from some of the highest cancer death rates in the nation.^{2–4} We are researching methods to improve cancer treatment and outcomes for AIs in western South Dakota. This program consists of patient navigation, clinical trials, surveys to evaluate barriers to access, and a molecular study (ATM [ataxia telangiectasia mutated] gene) to assess a potential molecular reason for increased treatment induced toxicities. To date, more than 1400 AIs have been enrolled in these studies over a period of 3 years. The phase II clinical trials use tomotherapy and brachytherapy to shorten the overall treatment time.^{5,6} In addition, more than 70 clinical trials are open through the cooperative group mechanism.

The *Walking Forward* program works with AIs on 3 reservations and in the Rapid City community. All 4 sites employ Community Research Representatives (CRRs) who serve as a bridge between the cancer center and the communities being served. CBPR is more than a principle to be followed for the CRRs; it is where they live and work. Multiple barriers have been identified, leading to interventions promoting cancer education and screening in hopes of diagnosing patients with earlier stages of cancer.

Because of the overwhelming success of the *Walking Forward* program in navigating patients, our cancer center has implemented a similar program for all patients. One navigator assists breast cancer patients through the continuum of cancer care. A second navigator works with the general population of cancer patients. It is a significant step for responsive patient services to an underserved rural population that reflects disparate access and outcomes to cancer care.

Rust and Cooper challenge us to meet the need of disparities in research in the community setting by providing 12 strategies to move forward. The CDRP model paired with community-based participatory research promotes an investment in health care that is responsive to

community needs and provides significant improvement in access and standard of care.

Mary Reiner, BA
John T. Vucurevich Cancer Care Institute,
Rapid City, SD
mreiner@rcrh.org
Daniel G. Petereit, MD
John T. Vucurevich Cancer Care Institute,
Rapid City, SD
Department of Human Oncology,
University of Wisconsin Paul P. Carbone
Comprehensive Cancer Center, Madison, WI

References

1. Rust G, Cooper LA. How can practice-based research contribute to the elimination of health disparities? *J Am Board Fam Med* 2007;20:105–14.
2. Petereit DG, Rogers D, Govern F, et al. Increasing access to clinical cancer trials and emerging technologies for minority populations: The Native American Project. *J Clin Oncol* 2004;22:4452–5.
3. Petereit DG, Rogers D, Burhansstipanov L, et al. Walking Forward: The South Dakota Native American Project. *J Cancer Educ* 2005;20(Suppl 1):65–70.
4. Rogers D, Petereit DG. Cancer disparities research partnership in Lakota Country: clinical trials, patient services, and community education for the Oglala, Rosebud and Cheyenne River Sioux tribes. *Am J Public Health* 2005; 95(12):2129–32.
5. Mackie TR, Kapatoes J, Ruchala K, et al. Image guidance for precise conformal radiotherapy. *Int J Radiat Oncol Biol Phys* 2003;56(1):89–105.
6. Eastmo E, Petereit DG. Accelerated partial breast irradiation: expanding options for breast preservation. *Adv Imaging Oncol Adm* 2005;91–5.

doi: 10.3122/jabfm.2008.01.070216

The above letter was referred to the authors of the article in question, who offer the following reply.

Building a Bridge Between Community-based Participatory Research (CBPR) and Primary Care Practice-based Research (PBR)

To the Editor: We are pleased to hear of these specific examples of community-based participatory research (CBPR) focused on cancer care and outcomes for American Indian communities in South Dakota. CBPR and primary care practice-based research (PBR) too often operate in separate silos, one focused on community-based, health-promoting interventions outside of clinical health care settings, and the other inherently conducted within the clinical practice. Reiner and Petereit describe a perfect example of how to build a bridge between these 2 important arenas of disparities research toward a common goal of improved health outcomes. The use of community health workers, navigators, *promotoras*, or in this case “Community Research Representatives,” are essential to bridging the culture gap between clinical practitioners and individuals in the communities they serve. As presented here, they also can play a key role in

bridging the gap between research investigators and research subjects.

Much more work needs to be done in this area to further refine the methodology for implementing research that has community oversight and to better capture the insights and the discovery that are generated by the community, rather than merely fostering one-way diffusion of medical innovation from university to community.

We applaud the efforts of Reiner and Petereit, and look forward to seeing the outcomes of their innovative work and to see their methods adapted in many more diverse settings across our nation.

George Rust, MD, MPH
Morehouse School of Medicine, National Center for
Primary Care, Atlanta, GA
grust@msm.edu
Lisa Cooper, MD, MPH
Johns Hopkins University School of Medicine,
Baltimore, MD

doi: 10.3122/jabfm.2008.01.070227

Tar Wars and Smoking Prevention: Premature Claim of Effectiveness

To the Editor: Tar Wars, as evaluated by Cain et al,¹ is part of a large movement of knowledge-based, early-education programs designed to inform elementary school students about the dangers of tobacco use. The authors have attempted to lend support to the claim that "school-based tobacco education is one part of a comprehensive approach to youth tobacco prevention activities that has been shown to decrease youth tobacco initiation." They cite several articles that describe existing programs. Their intervention seems to be quite similar to programs shown to be effective, which raises questions about the authors' claim that Tar Wars contributes anything uniquely effective. We respectfully question the methods presented by the authors and would like to make several suggestions in the hope that future assessments of the Tar Wars program might yield substantive evidence of effectiveness.

One of our concerns is that students in the Cain evaluation study were pre- and posttested after only 1 presentation; moreover, the full Tar Wars program with its various components was not included for testing. Considering that authors of previous studies of tobacco education programs waited 1 to 2 years to test whether information persisted to determine effectiveness, the significance of this short-term test is questionable. The test tapped into whether information would be remembered by students after only 9 to 11 days, and students were aware of the information on the pretest when they heard the presentation. This does not seem to represent an enduring test of knowledge or attitude change. We would suggest that the Tar Wars posttest be conducted after 1 year.

A second concern is that the quantitative measure used in the evaluation included only 14 items, and among them were 3 potentially confusing or irrelevant items about tobacco advertisements in the mass media. If we

eliminated these 3 as too ambiguous, the entire program effectiveness would be based on 11 questions, most of the answers to which were already known by the students. Of 14 questions, students on average answered 8 or 9 correctly on the pretest. Therefore, of the 11 unambiguous questions, there are only 2 previously unknown questions on which to gauge the effectiveness of the entire program. The authors should consider reformatting the questionnaire by adding questions and ensuring that the questions are clear and relevant.

The authors conclude that the students participating in the program indeed learned new information. Results showed an increase in correct responses for all questions. However, because of the methodology, the increase may potentially be explained by a practice effect. If children saw the questionnaire before the presentation and recognized the questions a few days later after the presentation, we fear that the increase in correct answers doesn't say much about the actual effectiveness of the intervention or about attitude change, only that they remembered test items they might have missed. Perhaps this could be remedied if the measures occurred further apart in time.

The authors acknowledge the lack of a control group in this effectiveness study. Students' knowledge after participating in the program was not compared with other students who did not receive the program intervention. However, the authors claim that other agents of change to explain the children's acquisition of knowledge are "unlikely" and insist that "major changes [to the program] are not needed." It would seem that the authors have no research foundation on which to make this claim. The lack of a control group with which to confirm the gain of knowledge combined with the questionable methods used substantially weaken the authors' claim that no improvements are necessary to the program.

Effectiveness cannot be attributed to Tar Wars if much of the program is repetition of previously known information. Although Tar Wars may eventually prove to be a valuable addition to school curricula, at this time it is unlikely that it has added anything unique or novel beyond similar programs.

Tonya Johnson, MA
Department of Psychology, University of Alabama,
Tuscaloosa, AL
Alan Blum, MD
Department of Family Medicine, University of
Alabama School of Medicine, Tuscaloosa, AL
ablum@cchs.ua.edu

Reference

1. Cain JJ, Dickinson WP, Fernald D, Bublitz C, Dickinson LM, West D. Family physicians and youth tobacco-free education: outcomes of the Colorado Tar Wars program. *J Am Board Fam Med* 2006;19(6):579–89.

doi: 10.3122/jabfm.2008.01.070189

The above letter was referred to the authors of the article in question, who offer the following reply.