EST in triage and the lack of effectiveness of EST in screening.

Peter G. Teichman, MD, MPA
FV Hospital
Ho Chi Minh City, Vietnam
keqfap@yahoo.co.nz

References

1. Newman RJ, Darrow M, Cummings DM. Predictive value of exercise stress testing in a family medicine population. *J Am Board Fam Med* 2008;21(6):531–38.

doi: 10.3122/jabfm.2009.02.080242

Dr. Newman declined to respond to this letter.

Re: Predictive Value of Exercise Stress Testing in a Family Medicine Population

We read with great interest the article by Newman et al titled “Predictive Value of Exercise Stress Testing in a Family Medicine Population.”

The results confirm the findings we have reported in our studies a few years ago.^{1,2} In our studies, the predic-

tive negative value of the test was 98% both among working-aged population and among older patients.

Therefore, we would like to point out that the authors’ statement to be the only ones to study predictive values of exercise stress testing for patient-oriented outcomes in family medicine is not entirely correct.

Markku Sumanen, MD, PhD
markku.sumanen@uta.fi

Kari Mattila, MD, PhD
University of Tampere, Medical School
Tampere, Finland

References

1. Sumanen M, Jussila M, Mattila K. Exercise treadmill test may predict clinical outcome among working-aged patients suspected of coronary heart disease in general practice. *Scand J Prim Health Care* 2005;23(1):47–51.
2. Sumanen M, Mattila K. A negative finding in an exercise test is reliable among elderly people: a follow-up study. *Gerontology* 2007;53(3):159–64.

doi: 10.3122/jabfm.2009.02.080243

Dr. Newman declined to respond to this letter.