CLINICAL REVIEW

Foot and Ankle Surgery: Considerations for the Geriatric Patient

Daniel K. Lee, DPM, and Gerit D. Mulder, DPM, MS

The growing number of lower-extremity abnormalities that are seen in inpatient and outpatient settings has paralleled the increased number of elderly in the population. Foot and ankle deformities, disorders, and arthritis, which are not manifested until late in life, have become more common as more individuals attain longer lifespans. Although conservative therapies are a priority when addressing the geriatric population, surgical options may be overlooked secondary to a misunderstanding of their ability to overcome perioperative management. Advanced minimally invasive surgical procedures for the foot and ankle have decreased the complications associated with foot surgery, making surgical intervention a viable option for many of the elderly. The newer procedures do not, however, minimize strict perioperative management, including pharmacological and nutritional assessment, and cardiopulmonary precautions. Outpatient surgical intervention may effectively address many ongoing problems associated with pain, decreased ambulation, and decreased quality of life. Current techniques in joint reconstruction in the forefoot and midfoot allow weight bearing from the day of surgery. Most hindfoot and ankle surgeries now permit minimal bone resection and incision through arthroscopy, resulting in improved muscle and tendon repair and early weight bearing. The changes in surgical approaches for the geriatric foot have permitted more effective and rapid intervention in problems affecting ambulation and quality of life in our aged population. (J Am Board Fam Med 2009;22:316–24.)

Patients over the age of 65 are the fastest growing segment of the population; this population numbers more than 30 million people (>12.5% of the population).^{1–3} Increased life spans have been promoted by advances in medicine, science, and healthy lifestyles. Promotion of physical activity and ongoing exercise among the elderly has resulted in these individuals being stronger and healthier than previous generations. Increased activity and increased lifespan both contribute to the development of extensive lower extremity problems, including degeneration of bone and joints. Ligaments, tendons, and muscles are more easily

damaged or injured, and the lower extremity, foot, and ankle have been specially affected.⁴ Once injuries or physical changes occur and impact quality of life, the question arises as to whether a conservative versus a surgical approach is in the patient's best interest. Numerous factors—including but not limited to medical history, physical and mental condition, and ambulatory status—must be considered because age alone is not a direct contraindication for surgical correction of foot and ankle deformities.

Advances in medicine and surgery are providing more low-risk and sophisticated treatment options that have not been previously available. Latest advances in surgical techniques and a better understanding of biomechanics have allowed us to perform surgery in the geriatric patient with functional results equivalent to those found among younger patients.^{5,6} Figure 1 illustrates successful surgical results of a bony deformity attributed to progressive arthritic changes in the foot. In the absence of any significant contraindications, a forefoot correctional procedure was performed, which decreased patient pain and allowed for improved quality of life.

This article was externally peer reviewed.

Submitted 8 June 2008; revised 20 October 2008; accepted 3 November 2008.

From the Department of Orthopaedic Surgery (DKL) and the Wound Treatment and Research Center and the Department of Surgery, Division of Trauma (GDM), University of California, San Diego.

Funding: none.

Conflict of interest: none declared.

Corresponding author: Daniel K. Lee, DPM, Assistant Clinical Professor, Department of Orthopaedic Surgery, University of California, 350 Dickinson Street, MC8894, San Diego, CA 92103-8894 (E-mail: dklee@ucsd.edu).

Figure 1. Forefoot arthritis. Preoperative (A) and postoperative (B) clinical views after first metatarsophalangeal joint and digital reconstruction.

Surgery in the geriatric patient had been avoided because it was associated with high morbidity and mortality rates related to surgical procedures and general anesthesia. Currently more than 20% of all surgical admissions in the United States are geriatric patients.⁷⁻⁹ The recent medical literature suggests that there were neither increased complications or deaths with the use of general and local anesthesia in the geriatric patient compared with previously known data.^{8,10,11} The medical literature suggests that the severity of the disease or illness is a better indicator or predictor of surgical and anesthesia outcome versus the patient's chronological age.^{12,13} The following is a review of considerations that may assist with referral of the geriatric patient for possible lower extremity surgery.

Surgical Considerations

Nutrition

Foot and ankle deformities, disorders, and arthritis may remain asymptomatic for years before becoming fixed, rigid, and painful among the aged population.¹¹ They are common in the geriatric patient and may lead to loss of lower extremity mobility and function, which in turn may contribute to deterioration of the cardiovascular system, a rapid decline in health, and a reduced life span. The musculoskeletal system undergoes significant change during the aging process as osteoarthritis and osteoporosis develop and progress, especially in women after menopause. Bone loses rigidity and strength and becomes more brittle. Joints and surrounding soft tissue become weak and less flexible

with aging.^{14,15} It is important to consider pharmacological and nutritional management pre- and postoperatively to enhance bone quality and healing.¹⁶⁻¹⁹ Maintaining adequate protein levels as well as appropriate dietary supplementation may assist the healing process in patients considered nutritionally compromised. Patient's protein levels need to be assessed, along with a review of the patient's dietary habits. Special considerations need to be given to obese patients because obesity may be an indicator of poor nutritional status. Obesity also places excessive stress on the lower extremity and may contribute to poor healing and surgical outcomes. Laboratory tests recommended for determining nutritional status include patient's body mass index (BMI; weight in kilograms/height in meters [BMI < 17 kg/m² is associated with protein-energy undernutrition]); serum albumin (<3.5 mg/dL is a simple indicator of possible malnutrition); serum levels of vitamins A, B₁, B₁₂, C, D, E, iron, folic acid, zinc, and magnesium; serum creatinine levels; total urine nitrogen; complete blood count; comprehensive metabolic panel; liver function test; and lipid panel.

Preoperative Medical History

Preexisting medical conditions are of as much concern as nutritional status. Medications affecting cellular function and fibroblast activity are known to delay the closure process. It is not uncommon for elderly patients to be taking high levels of nonsteroidal - medications or possibly even steroidal medications for indications, including various arthritic or vasculitic conditions. A history of a previous deep venous thrombosis is important to determine because the risk of a postoperative deep venous thrombosis is increased if the patient has had one previously. Age, sedentary lifestyle, history of previous lower extremity trauma, hypercoagulability, and even family history of deep venous thrombosis are reasons for anticoagulant prophylaxis. Prophylaxis is a further consideration for patients who are expected to remain in bed for >48 hours or whose ambulation will be limited after discharge. Patients with a short hospital stay (<48 hours) and immediate ambulation still require the use of a compression device during the admission period to decrease risk of clot formation. There have been studies to evaluate these guidelines, but the incidence of venous thromboembolism after foot and ankle surgery has been rare (<1%) and the need for routine

propylaxis postoperatively is not supported by any high level of evidence studies.^{20–22}

Preoperative Examination

Planning for surgery in the geriatric patient involves all the requirements and criteria of the adult patient with special emphasis on preoperative cardiopulmonary precautions, appropriate surgical procedure selection, and proper postoperative management to maximize bone healing.²³ The surgeon should consider obtaining medical clearance for the planned procedure from the patient's primary care physician and cardiologist.

Indications of surgery in the medically cleared patient include the failure of previously attempted conservative management and pain, deformity, and/or functional disability that significantly impairs quality of life. When conservative management provides satisfactory results, surgery should not be encouraged. Postoperative compliance and a suitable postoperative living environment are additional preoperative considerations. Selected patients may require postoperative home care and physical therapy or short-term placement in a rehabilitative or skilled nursing facility. Preoperative considerations include the ability of the planned procedure to address pain, correct deformities, and to restore function, allowing patients to resume their activities of daily living and improve their quality of life.

Preoperatively, patients will require clinical, biomechanical, radiographic, social, and psychological evaluation and preparation. It is imperative to have a thorough discussion of the diagnosis, surgical treatment options, and prognosis with the patient and family involved. Patients may have unrealistic expectations about surgical procedures and may make the assumption that a normal "youthful" lower extremity will result from the intervention. Advantages and disadvantages of the procedure, expected levels of function and pain, level of postoperative care, and anticipated affect on ambulation require clear explanations and realistic expectations before obtaining a signature on an informed consent. Conservative options should have been tried and failed before surgical intervention.

Before scheduling procedures, it is imperative to obtain a medical clearance from the patient's geriatric physician. This is recommended to ascertain the cardiovascular and pulmonary status and risks of perioperative myocardial events, even among

healthy geriatric patients.^{12,24–27} Cardiovascular and pulmonary status are directly related to the rates of morbidity and mortality, and they can pose greater risks and complication when they are not managed properly.^{7,28–30} Functional assessments using activities of daily living scales can predict perioperative rehabilitation requirements based on observations of their functions and multiple customary activity assessment.^{31–33} The American Society of Anesthesiologists criteria are also useful guides in the surgical risk assessment and perioperative management of the geriatric patient.^{8,11,34} When there is doubt concerning the patient's ability to tolerate a procedure, it is wise to choose the conservative option.

Postoperative Considerations

The majority of foot and ankle surgery is performed in an outpatient setting. However, this should not minimize strict perioperative management. Emphasis on mobility and function and aggressive rehabilitation for deep venous thrombosis prophylaxis is important.^{22,35–37} Time for foot and ankle surgery rehabilitation has previously been documented as 1½ to 2 times that of a young patient, with precautions for a short-leg non-weight-bearing cast, contralateral limb injury, and/or potential for over use and falling.¹¹ With the advent of advanced surgical techniques and ambulatory devices, these practices are changing. Early ambulation is recommended to decrease adverse effects of anesthesia after surgery while minimizing the risks of cardiopulmonary complications commonly associated with surgery and anesthesia. Pain management may be accomplished with regional blocks and continuous local infusion systems, which have eliminated or minimized the intake of postoperative pain medications and narcotics.^{12,13} Consideration of all the factors previously discussed allow for selection of the appropriate postoperative setting to assist with rapid recovery and a return to the patient's previous lifestyle.

Surgical Options for Eligible Candidates

Simple surgical procedures with local anesthesia may provide dramatic relief, permitting normal ambulation and resumption of daily activities while prolonging the length and quality of life. Selection of the most appropriate surgical procedure is par-

Table 1. Common Foot and Ankle Surgical Terms

Procedure	Description
AO fixation	Swiss method of application of internal fixation (Arbeitsgemeinschaft Fuer Osteosynthesefragen = Association for the Study of Internal Fixation)
Arthrodesis	Surgical immobilization of a joint by cartilage removal, usually with fixation
Arthroplasty	Formation or restoration of a joint, usually with a metallic or nonmetallic implant
Cheilectomy	Dorsal bone resection, typically of the first metatarsal head
Chondroplasty	Restorative or reshaping surgery of the cartilage
Gastroc recession	Lengthening of gastrocnemius aponeurosis because of gastrocnemius equinus or tightness
Ilizarov method	Russian method of application of external fixation with fine wires and ring fixators
Keller procedure	Resection of the base of proximal phalanx of hallux
Osteotomy	Bone division to correct a deformity, usually stabilized with fixation (Akin for proximal phalanx; Chevron, SCARF, crescentic for first metatarsal; Dwyer for calcaneus)
Synovectomy	Removing part or all of the synovial membrane
Tendon transfer	Shifting a functioning tendon from its original attachment to a new one to restore the action that has been lost
Tendon/Achilles lengthening	Lengthening of Achilles tendon because of gastrocsoleus equinus deformity or tightness, commonly known as TAL

amount to the successful outcome of the surgery. Selection of the specific surgical procedure is based on the patient's health, the location of the problem, the level of severity/deformity, the need for single or multiple staged procedures, the length of rehabilitation, and postoperative course. New advanced procedures allow early weight-bearing and minimal to no use of any non-weight-bearing casts.²² It is not our purpose to describe here the detailed surgical technique of the procedures that will be mentioned; however, a definition of the more common procedures described is given in Table 1.

Osteoarthritis and bony deformity are frequent findings in the forefoot and midfoot of the elderly.³⁸ The etiology can be multifactorial and include muscle imbalance, flat footedness, and trauma. Severity of deformities may be more pronounced with rheumatoid arthritis and other systemic arthritic

Figure 2. Hallux valgus. Note the severe contractures and deviations in the forefoot.

conditions. Patients typically complain of pain and disability causing difficulty in ambulation, primarily with forefoot loading and propulsion phases of gait, as well as pain with shoe wear at the forefoot or distally from the midfoot. Corns and callouses are findings that may reflect an underlying osseous deformity.

Forefoot Procedures

The most common forefoot deforming arthritic conditions include hammertoes, hallux valgus ("medial bunion"), hallux rigidus ("dorsal bunion"), metatarsophalangeal joint derangement, metatarsalgia, first ray hypermobility and instability, tailor's bunion (fifth metatarsal bunion), and posttraumatic arthritis and deformity (Lisfranc injury, fractures of the phalanx, metatarsal, cuneiforms, cuboid, navicular, and Charcot foot). Figures 2, 3, and 4 These deformities are primarily addressed with less complicated osteotomies and minor bone excision.

These procedures are usually performed under intravenous sedation (monitored anesthesia care) with regional local blocks (marcaine/lidocaine) in an outpatient setting^{39,40} (Figure 5). Most patients are treated with a postoperative weight-bearing shoe or a temporary non-weight-bearing splint, which is eventually switched to a walking boot. Patients with gait instability may prefer a walker or Roll-A-Bout device (Roll-A-Bout Corporation, Frederica, DE; Figure 6) because they provide

Figure 3. Tailor's bunionette. Note the angular osseous malalignment of the 5th metatarsal.

3-point walking stability over the traditional cane or crutches.

Digital procedures including arthroplasties or arthrodeses correct multiplanar deformities of the proximal and distal interphalangeal joints. These corrections allow proper alignment of digits and the removal of painful joint surfaces for ease of shoe wear and prevention of arthritic ulcerations.

Midfoot Procedures

Procedures at the first metatarsophalangeal joints are divided into cheilectomy, osteotomy, implant arthroplasty, and arthrodesis.⁴¹ They are either joint-sparing or joint-replacing procedures. Joint-sparing procedures (cheilectomy, osteotomy) have an excellent outcome in the presence of end-stage arthritis. Increased deformities have had better outcomes with joint-replacing procedures (implant arthroplasty and arthrodesis). Metatarsophalangeal joint resections and Keller-type procedures are usually reserved for end-stage conditions in which ambulation and flexibility are not a concern. Instability and posttraumatic arthritis in the tarsometatarsal joints require bone resection, which is the simplest approach, or arthrodesis to eliminate the source of pain and provide stability. Although bone resection does not require the use of fixation devices, arthrodesis requires joint preparation and fixation. These particular joints are not essential for gait. Their range of motion is minimal compared

Figure 4. Diabetic Charcot deformity. Note the osseous destruction and collapse in the midfoot preoperatively (A) and the postoperative view of the reconstruction (B).

with the essential joints of the ankle, subtalar, midtarsal, and first metatarsophalangeal joints.

The fusion of tarsometatarsal joints provides significant pain relief and stability to the midfoot in stance and gait. With the introduction of external fixation they may now be used in combination with internal fixation for further added stability of these bone segments, allowing the patient to perform protected partial to full ambulation postoperatively, which previously required 4 to 8 weeks of non-weight-bearing immobilization.

Hindfoot Procedures

At the hindfoot and ankle levels, arthritis, deformity, and muscle imbalance can be common in the geriatric patient. Similar to the forefoot and midfoot, the causes can also be multifactorial and result from osteoarthritis or stroke. The arthritic events affecting the forefoot and midfoot can also affect the hindfoot and ankle. The ankle, subtalar, and midtarsal (talonavicular and calcaneocuboid) joints can be affected in isolation or combination. These

Figure 5. Regional blocks: (A) ankle and (B) popliteal.

joints are very complex and multiplanar in range of motion. Their 3-dimensional joint motion leads to a combination of arthritic events with joint crepitus at multiple levels. Neuromuscular conditions can affect the distal extrinsic muscles in the lower extremity leading to muscle imbalance, weakness, spasticity, and contractures.

The most common conditions include ankle arthritis, ankle valgum/varum deformity, drop

foot, equinus, Achilles partial/full rupture, posterior tibial tendon dysfunction/insufficiency, subtalar arthritis, subtalar valgum/varum deformity, midtarsal joint arthritis, collapsed hindfoot complex, posttraumatic joints, and nonunion/malunion conditions. Typically patients with these conditions complain of pain around the heel and/or ankle. Disability is associated with difficulty standing and/or walking for long distances. It is not uncommon for muscle weakness and imbalance go unnoticed by the patient (Figure 7) During examination the clinician can determine the level of arthritis, misalignment, and deformity through muscle testing and evaluation of range of motion and gait.

The goals for geriatric hindfoot and ankle surgery are focused on achieving a plantigrade foot, allowing full ground contact, ambulation with a brace, and elimination of the need for a brace.³⁸ Most of these procedures can now be performed under intravenous sedation (monitored anesthesia care) with regional popliteal, sciatic, or femoral local blocks in an outpatient setting. Unlike forefoot and midfoot procedures, most patients are protected with a temporary non-weight-bearing splint, which is then switched to a short-leg partial-walking cast after 2 to 4 weeks, and then to a full-walking cast during the following 3 to 4 weeks. In selected patients, an Ilizarov external fixation may be applied; this can allow postoperative weight bearing beginning in the first 1 to 2 weeks with a walking aid. Patients undergoing an Ilizarov procedure must be selected with special caution because strict compliance is needed (Figure 8). The daily postoperative care for these more complicated

Figure 6. Non-weight bearing assistance devices as alternatives to traditional crutches and walkers.

Figure 7. Chronic Achilles tendon rupture. Note (A) the clinical view of the interrupted integrity of the Achilles tendon; (B) T2-weighted images of the ruptured Achilles tendon; (C) intraoperative view of the ruptured Achilles tendon; and (D) Achilles tendon reconstruction with graft.

procedures are best addressed in a rehabilitative or skilled nursing facility.

Arthrodesis of the ankle and subtalar joints is still the gold standard in the treatment of end-stage arthritis (Figure 9). Although joint replacements that provide increased range of motion and flexibility are treatment options, the intermediate and short-term results are not as satisfactory and have

Figure 8. Severe ankle osteoarthritis. Note the Ilizarov external fixation for earlier postoperative weight-bearing tolerance.

Figure 9. Subtalar joint arthritis. A postoperative view after isolated subtalar joint arthrodesis.

higher complication rates compared with knee and hip joint replacements (Figure 10). Until this technology improves, extra-articular arthrodesis and joint resection with synovectomy and debridement are better options for geriatric patients. Isolated midtarsal joints arthrodesis reduces pain and disability as well as total range of motion of the subtalar joint. These procedures can proceed with early weight bearing compared with ankle and subtalar joint arthrodesis. There are currently no replacements available for these joints. Osteotomies in the ankle and hindfoot are viable extra-articular procedures, which preserve joints and provide realignment of the structures.

Tendinopathies associated the Achilles and posterior tibial tendon are the most frequently performed procedures. Although rupture repairs of the Achilles tendon is normally performed in isolation, repairs of the posterior tibial tendon in isolation without bone correction or realignment do not provide enough stability to hold the correction. Because most posterior tibial tendon conditions occur with progressive pes plano valgus and flatfoot deformity, the correction of bone pathology has

Figure 10. Ankle arthritis. A postoperative view of total ankle replacement. Note the complete bipolar components for the tibiotalar joint.

priority over the tendon repair. Drop foot requires evaluation of in-phase and out-phase muscles because tendon transfer techniques can prevent ankle arthrodesis.^{42,43}

Well-designed and well-controlled clinical trials among the geriatric population are still needed to validate the improvement in reported clinical outcomes and to assist the primary care provider in giving a more accurate description of outcome expectations to the patient. Advantages, disadvantages, benefits, risks, and time to recovery need to be clearly covered by the surgeon, although it is helpful for the primary care physician to inform the patient of options for which an educational visit and consult with the surgeon may be made. We strongly believe that the geriatric patient who is asymptomatic, able to ambulate without significant difficulty, and who is not in a limb threatening situation should not undergo a surgical procedure simply for cosmetic purposes. The primary care physician must carefully screen and the patient's current medical situation and medical and social history to determine whether surgical intervention should be considered. Patients, regardless of age, commonly anticipate that a surgical procedure will create an "anatomically normal," fully functional, and completely pain-free foot. Surgical procedures are meant to address problematic foot problems, improve ambulation, and to decrease pain. Patients may need to be educated postoperatively about ongoing, albeit decreased, pain, the need for special shoes, and limitations to daily activities. Patients will appreciate an honest and straightforward explanation of what to expect.

Currently, well-powered and controlled trials are not available to provide evidence-based and statistically significant data about the new and advanced surgeries that result in less pain, decreased recovery time, and improved function scores. The aforementioned lack of studies highlights the need for determining the correlation between perioperative medical management and treatment outcomes, as well as any relationship between newer techniques and surgeries on the elderly and common postoperative complications such as the incidence of deep venous thrombosis/pulmonary embolism (PE), postoperative pneumonia, or postoperative injuries (falls). The rise in numbers of senior patients and the parallel increase in surgeries performed on this population justify the need for such studies to be performed in the near future.

Summary

The increase in the aged population and their continued active lifestyles and a desire for a better quality of life make it important for the health care provider to have a basic understanding of conservative versus surgical options for treatment of the geriatric lower extremity. Surgical versus conservative approaches have been emphasized here; however, potential problems from operative intervention as well as postoperative care need to be discussed in detail with patients and family members. Advanced foot and ankle surgery is not immune from common postoperative orthopedic and cardiovascular complications. Conservative options need to be considered before surgery is recommended. Candidates that do qualify for surgery may now benefit from the advanced and improved treatments that have been developed to address lower extremity problems among the aging population.

References

1. US Bureau of the Census, Population Division. United States population estimates, by age, sex, race, and Hispanic origin, 1990–1997. Release PPL-91. Washington, DC: US Bureau of the Census; 1998.
2. Johnson JC. The medical evaluation and management of the elderly surgical patient. *J Am Geriatr Soc* 1983;33:621–5.
3. Mohr DN. Estimation of surgical risk in the elderly: a correlative review. *J Am Geriatr Soc* 1983;33:99.
4. Sobel E, Giorgini RJ. Surgical considerations in the geriatric patient. *Clin Podiatr Med Surg* 2003;20:607–26.
5. Shah MR, Aharonoff GB, Wolinsky P, et al. Outcome after hip fracture in individuals ninety years of age and older. *J Ortho Trauma* 2001;15:34–9.
6. Berger DH, Roslyn JJ. Cancer surgery in the elderly. *Clin Geriatr Med* 1997;13:119–41.
7. Hirsch CH. When your patient needs surgery: weighing risks versus benefits. *Geriatrics* 1995;50:26–31.
8. Hackford AW. Surgical principles for the aged. In: Reichel W, editor. *Care of the elderly clinical aspects of aging*. 4th ed. Baltimore: Williams & Wilkins; 1995:408–15.
9. Djokovic JL, Hedley-Whyte J. Prediction of outcome of surgery and anesthesia in patients over 80. *JAMA* 1979;242:2301–6.
10. Kaufman JL. Local anesthesia for surgery on the foot: efficiency in the ischemic or diabetic extremity. *Ann Vasc Surg* 1991;5:354–8.
11. Hoskings MP, Warner MA, Lobdell CM, et al. Out-

- comes of surgery in patients 90 years of age and older. *JAMA* 1989;261:1909–15.
12. Dunlop WE, Rosenblood L, Lawrason L, et al. Effects of age and severity of illness on outcome and length of stay in geriatric patients. *Am J Surg* 1993;165:577–80.
13. Adkins RB Jr, Scott HW Jr. Surgical procedures in patients 90 years and older. *South Med J* 1984;77:1357–64.
14. Gajdosik RL, Vander Linden DW, Williams AK. Influence of age on concentric isokinetic torque and passive extensibility variables of the calf muscles of women. *Eur J Appl Physiol Occup Physiol* 1996;74:279–86.
15. Alnaqeeb MA, Alzaid NS, Goldspink G. Connective tissue changes and physical properties of developing and aging skeletal muscle. *J Anat* 1984;139:677–89.
16. Howard KW. The geriatric patient. In: McGlamry DB, editor. *Fundamentals of foot surgery*. Baltimore: Williams & Wilkins; 1987:423–35.
17. Bernard RL, Giorgini RJ, Giorgini TL. Podiatric surgical considerations in the older patient. *Clin Podiatr Med Surg* 1993;10:129–36.
18. Delmi M, Rapin CH, Bengoa JM, et al. Dietary supplementation in elderly patients with fractured neck of the femur. *Lancet* 1990;335:1013–6.
19. Morley JE, Silver AJ, Fiatarone M, Mooradian AD. Geriatric grand rounds: nutrition and the elderly. *J Am Geriatr Soc* 1986;34:823–32.
20. Wukich DK, Waters DH. Thromboembolism following foot and ankle surgery: a case series and literature review. *J Foot Ankle Surg* 2008;47:243–9.
21. Gadgil A, Thomas RH. Current trends in thromboprophylaxis in surgery of the foot and ankle. *Foot Ankle Int* 2007;28:1069–73.
22. Wolf JM, DiGiovanni CW. A survey of orthopedic surgeons regarding DVT prophylaxis in foot and ankle trauma surgery. *Orthopedics* 2004;27:504–8.
23. Hirsch CH. When your patient needs surgery: how planning can avoid complications. *Geriatrics* 1995;50:39–44.
24. Miller DL. Perioperative care of the elderly patient: special considerations. *Cleve Clin J Med* 1995;62:383–90.
25. Meneilly GS, Rowe JW, Minaker KL. Anesthesia and surgery in the elderly. In: Rowe JW, Besdine RW, editors. *Geriatric medicine*. 2nd ed. Boston: Little, Brown; 1988:122–35.
26. Goldman L, Caldera DL, Nussbaum SR, et al. Multifactorial index of cardiac risk in noncardiac surgical procedures. *N Engl J Med* 1977;297:845–50.
27. Foster ED, Davis KB, Carpenter JA, et al. Risk of noncardiac operation in patients with defined coronary disease: The Coronary Artery Surgery Study (CASS) registry experience. *Ann Thorac Surg* 1986;41:42–50.
28. Evers BM, Townsend CM Jr, Thompson JC. Organ physiology of aging. *Surg Clin North Am* 1994;74:23–39.
29. Seymour DG, Pringle R. Postoperative complications in the elderly surgical patient. *Gerontology* 1983;29:262–70.
30. Vaz FG, Seymour DG. A prospective study of elderly general surgical patients I: preoperative medical problems. *Age Aging* 1989;18:309–15.
31. Rozanski A, Bairey CN, Krantz DS, et al. Mental stress and the induction of silent myocardial ischemia in patients with coronary artery disease. *N Engl J Med* 1988;318:1005–12.
32. Katz S, Ford A, Moskowitz RW, et al. Studies of illness in the aged. The index ADL: a standardized measure of biological psychosocial function. *JAMA* 1963;185:914–9.
33. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969;9:179–86.
34. Cohen SJ. Elective podiatric surgery in the geriatric patient. *J Am Podiatr Med Assoc* 1976;66:886–90.
35. Wolf JM, DiGiovanni CW. A survey of orthopedic surgeons regarding DVT prophylaxis in foot and ankle trauma surgery. *Orthopedics* 2004;27:504–8.
36. Slaybaugh RS, Beasley BD, Massa EG. Deep venous thrombosis risk assessment, incidence, and prophylaxis in foot and ankle surgery. *Clin Podiatr Med Surg* 2003;20:269–89.
37. Jegathesan S, Whitehill JL. Surgery in the elderly. *Postgrad Med* 1971;49:211.
38. Kuhns JG. The foot in chronic arthritis. *Clin Orthop* 1960;16:141.
39. Hansen E, Eshelman MR, Cracchiolo A. Popliteal fossa neural blockade as the sole anesthetic technique for outpatient foot and ankle surgery. *Foot Ankle Int* 2000;21:38–44.
40. Myerson MS, Ruland CM, Allon SM. Regional anesthesia for foot and ankle surgery. *Foot Ankle* 1992;13:282–8.
41. McDonough MW. Geriatric hallux valgus deformity. In: McGlamry ED, Banks AS, Downey MS, editors. 2nd ed. *Comprehensive textbook of foot surgery*. Volume 1. Baltimore: Williams & Wilkins; 2000:578–86.
42. Yamamoto H, Okumura S, Morita S, et al. Surgical correction of foot deformities after stroke. *Clin Orthop* 1992;282:213–8.
43. Waters RL, Botte MJ, Jordan C, et al. Symposium: rehabilitation of stroke patients—the role of the orthopaedic surgeon. *Contemp Orthop* 1990;20:311–46.