SPECIAL COMMUNICATION

Birth History Is Forever: Implications for Family Medicine

Casey Crump, MD, PhD

The rapidly growing number of adult survivors of preterm birth has necessitated and made possible for the first time large-scale investigations of long-term outcomes of preterm birth. Large epidemiologic studies have shown that the long-term sequelae are wide-ranging, including metabolic disorders, cardiovascular and respiratory disease, psychiatric disorders, and increased mortality risk. Clinicians should now recognize preterm birth as a long-term, multidisease risk factor in adults. These research findings contribute to a growing body of evidence of early life programming for chronic disease, which in turn supports a “life course” paradigm for patient care. Family medicine is an ideally conceived discipline for this paradigm because of its unique role in caring for patients across the entire life span. As our understanding of early life influences on long-term health continues to advance, family physicians are ideally positioned to incorporate this knowledge into clinical practice. (J Am Board Fam Med 2015; 28:121–123.)

Keywords: Birth, Chronic Disease, Epidemiology, Longitudinal Studies

Preterm birth (<37 weeks of gestation completed) has become more prevalent over the past four decades and now accounts for nearly 12% of all US births (~500,000 births annually).¹ It is the leading cause of perinatal morbidity and mortality in the United States and other developed countries.¹ Its health outcomes during infancy and early childhood, including respiratory complications and increased mortality, have been widely studied.^{2,3} Until recently, the available evidence suggested that these health risks wane during and after childhood

and thereafter begin to approach similar levels as the general population.^{2,4} Data on longer-term outcomes have been more scarce. However, starting with neonatal care advances in the 1970s and 1980s, early survival among preterm infants improved dramatically, and the first generation to benefit from those advances has now reached young adulthood. The rapidly growing number of adult survivors of preterm birth now necessitates and has made possible, for the first time, large-scale investigations of longer-term outcomes into adulthood.

Large epidemiologic studies subsequently found that preterm birth has wide-ranging sequelae that persist into adulthood,³ including increased risks of diabetes,^{5,6} hypertension,^{7,8} venous thromboembolism,⁹ asthma,¹⁰ infections,¹¹ thyroid disorders,¹² epilepsy,¹³ and psychiatric disorders.¹⁴ These health sequelae also carry increased mortality risks in adulthood. A large national cohort study in Sweden found that among all people who survived to young adulthood, those who were born preterm had a ~40% increased risk of dying during young adulthood (ages 18–36 years) compared with those who were born at full term (adjusted hazard ratio for all-cause mortality, 1.4; 95% confidence interval, 1.2–1.6).^{15,16} This risk increased by earlier gesta-

This article was externally peer reviewed.

Submitted 10 December 2013; revised 7 August 2014; accepted 11 August 2014.

From the Department of Medicine, Stanford University, Palo Alto, CA.

Funding: This work was supported by grants from the National Institute of Child Health and Human Development (R01HD052848), the Swedish Research Council (2008–3110 and 2008–2638), the Swedish Council for Working Life and Social Research (2006–0386, 2007–1754, and 2007–1962), and ALF project grant, Lund, Sweden.

Conflict of interest: none declared.

Disclaimer: The funding agencies had no role in the design and conduct of the study; in the collection, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

Corresponding author: Casey Crump, MD, PhD, Department of Medicine, Stanford University, 211 Quarry Road, Suite 405, MC 5985, Palo Alto, CA 94304-1426 (E-mail: kccrump@stanford.edu).

tional ages and was ~2-fold among young adults who were born extremely preterm (<28 weeks).¹⁵ This increased mortality had multiple causes, including diabetes, cardiovascular disease, and respiratory disease, and was not explained by sociodemographic differences or birth defects that are more common with preterm birth. Although longer follow-up into later adulthood is needed, other studies of diabetes and cardiovascular disease have suggested that the risks associated with preterm birth persist at older ages. The continued high prevalence of preterm birth means that the long-term health effects will have a growing clinical and public health impact, reinforcing the need for better prevention of preterm birth.

These research discoveries contribute to a growing body of evidence of the origins of chronic disease in early life. According to the “developmental origins hypothesis,” variations in fetal or postnatal nutrition may permanently alter gene expression, resulting in early life programming for the onset of chronic disease in later life.^{17,18} The underlying mechanisms involve a complex interplay of hormonal and immunologic processes that are still being delineated and may be commonly associated with preterm birth. The cumulative evidence from this research shows that factors in early life can have important long-term health effects throughout the life span.^{3,17,18}

These findings have important implications for clinical practice and family medicine in particular. First, preterm birth should now be recognized as a long-term multidisease risk factor in adults.¹⁹ Despite the substantial research evidence of long-term sequelae, clinical standards of care do not yet reflect this knowledge. For example, relatively few physicians ascertain birth history in patients beyond childhood. A UK survey of respiratory specialists found that only a small minority (<25%, mostly hospital pediatricians) asked “most respiratory patients” about birth history such as preterm birth, birth weight, or perinatal complications, and a large proportion did not ask patients about birth history at all.²⁰ In primary care settings, ascertainment of birth history among adult patients is not well studied but is likely to be even less common, despite the fact that the reported mortality risks associated with preterm birth^{15,6} are higher than those of other factors commonly elicited by medical history, such as physical inactivity, poor nutrition, or heavy alcohol use.^{21,22} Although birth history (unlike life-

style factors) is not modifiable, better awareness of the associated risks can help motivate healthy behavior change to reduce other common risk factors.

Ascertaining a history of preterm birth among adults is now supported by the substantial evidence of long-term health risks. The medical history of adults, such as that taken during a periodic health evaluation, should encompass birth history including gestational age (or preterm birth [yes/no]), birth weight, and perinatal complications, when available.¹⁹ *International Classification of Diseases* (ICD) codes can be used to denote preterm birth (765 in ICD-9, P07 in ICD-10) or small for gestational age (764 in ICD-9, P05 in ICD-10) in the medical record, and further subclassifications can indicate birth weight. Such information can enable physicians and patients to be better aware of the potential health risks (such as metabolic disorders and cardiovascular disease) throughout life, consider more intensive screening for such risks, motivate healthy behavior change to reduce other modifiable risk factors, and better understand the context of chronic diseases when they occur.¹⁹ Additional research is needed to determine the predictive significance of preterm birth for specific clinical profiles. Nonetheless, birth history can provide important context for understanding a patient's health.

Systematic tracking of birth history information should be developed in health care systems in which it does not currently exist to facilitate better long-term patient care, as well as additional research on health outcomes. When birth records are unavailable, maternal recall of gestational age (or birth weight) has been reported to be sufficiently accurate for clinical use,²³ but self-reported birth history among adult patients has not been well studied. The accuracy of self-reported birth information would, however, be expected to improve as public and media awareness of its long-term relevance also improves.

These research findings have further implications for family medicine in particular. The substantial and growing evidence for early life origins of chronic disease supports a “life course” paradigm for patient care. In such a paradigm, health and illness are conceived not primarily in terms of episodic events marked by health care encounters, but within the longitudinal trajectory of an entire life span.²⁴ This perspective is consistent with family medicine's traditional emphasis on a highly contex-

tual approach to patient care, seeking to understand individual health in the context of relationships and function within family and community. However, the life course paradigm also adds to this contextual background a temporal dimension that is critically important for integrating knowledge about health risks that often originate before birth and continue to develop into adulthood. Family medicine is an ideally conceived discipline for this approach because of its unique role in caring for patients across the entire life span. As our understanding of early life influences on long-term health continues to advance, family physicians are ideally positioned to incorporate this new knowledge into clinical practice.

References

1. Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet* 2008;371:75–84.
2. Baraldi E, Filippone M. Chronic lung disease after premature birth. *N Engl J Med* 2007;357:1946–55.
3. Saigal S, Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet* 2008;371:261–9.
4. Swamy GK, Ostbye T, Skjaerven R. Association of preterm birth with long-term survival, reproduction, and next-generation preterm birth. *JAMA* 2008;299:1429–36.
5. Crump C, Winkleby MA, Sundquist K, Sundquist J. Risk of diabetes among young adults born preterm in Sweden. *Diabetes Care* 2011;34:1109–13.
6. Hofman PL, Regan F, Jackson WE, et al. Premature birth and later insulin resistance. *N Engl J Med* 2004;351:2179–86.
7. Crump C, Winkleby MA, Sundquist K, Sundquist J. Risk of hypertension among young adults who were born preterm: a Swedish national study of 636,000 births. *Am J Epidemiol* 2011;173:797–803.
8. de Jong F, Monuteaux MC, van Elburg RM, Gillman MW, Belfort MB. Systematic review and meta-analysis of preterm birth and later systolic blood pressure. *Hypertension* 2012;59:226–34.
9. Zöller B, Li X, Sundquist J, Sundquist K, Crump C. Gestational age and risk of venous thromboembolism from birth through young adulthood. *Pediatrics* 2014;134:e473–80.
10. Crump C, Winkleby MA, Sundquist J, Sundquist K. Risk of asthma in young adults who were born preterm: a Swedish national cohort study. *Pediatrics* 2011;127:e913–20.
11. Crump C, Sundquist K, Sundquist J. Prematurity and mortality in childhood and early adulthood (reply). *JAMA* 2012;307:32–3.
12. Crump C, Winkleby MA, Sundquist J, Sundquist K. Preterm birth and risk of medically treated hypothyroidism in young adulthood. *Clin Endocrinol (Oxf)* 2011;75:255–60.
13. Crump C, Sundquist K, Winkleby MA, Sundquist J. Preterm birth and risk of epilepsy in Swedish adults. *Neurology* 2011;77:1376–82.
14. Crump C, Winkleby MA, Sundquist K, Sundquist J. Preterm birth and psychiatric medication prescription in young adulthood: a Swedish national cohort study. *Int J Epidemiol* 2010;39:1522–30.
15. Crump C, Sundquist K, Sundquist J, Winkleby MA. Gestational age at birth and mortality in young adulthood. *JAMA* 2011;306:1233–40.
16. Crump C, Sundquist K, Winkleby MA, Sundquist J. Early-term birth (37–38 weeks) and mortality in young adulthood. *Epidemiology* 2013;24:270–6.
17. Barker DJ. The origins of the developmental origins theory. *J Intern Med* 2007;261:412–7.
18. Gluckman PD, Hanson MA. Living with the past: evolution, development, and patterns of disease. *Science* 2004;305:1733–6.
19. Crump C. Medical history taking in adults should include questions about preterm birth. *BMJ* 2014;349:g4860.
20. Bolton CE, Bush A, Hurst JR, et al. Are early life factors considered when managing respiratory disease? A British Thoracic Society survey of current practice. *Thorax* 2012;67:1110.
21. Khaw KT, Wareham N, Bingham S, Welch A, Luben R, Day N. Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. *PLoS Med* 2008;5:e12.
22. Kvaavik E, Batty GD, Ursin G, Huxley R, Gale CR. Influence of individual and combined health behaviors on total and cause-specific mortality in men and women: the United Kingdom health and lifestyle survey. *Arch Intern Med* 2010;170:711–8.
23. Adegboye AR, Heitmann B. Accuracy and correlates of maternal recall of birthweight and gestational age. *BJOG* 2008;115:886–93.
24. Daaleman TP, Elder GH Jr. Family medicine and the life course paradigm. *J Am Board Fam Med* 2007;20:85–92.