

## Correspondence

---

### Response: Re: Clinical Diagnosis of Bordetella Pertussis Infection: A Systematic Review

*To the Editor:* We thank Dr. Poddighi for his thoughtful commentary on our article regarding the clinical diagnosis of pertussis.<sup>1</sup> We fully agree that the overall clinical impression, sometimes called “clinical gestalt,” is a vital diagnostic tool for many conditions. In fact, we are in the process of performing a systematic review of clinical gestalt for the diagnosis of a range of respiratory infections; we hope that study will shed more light on its accuracy.

Well-validated clinical decision rules (CDRs) can be very useful to support a physician’s clinical gestalt. Of course, it is important to remember that CDRs are clinical decision *support* tools, not decision *replacement* tools. They must always be applied in the full context of what is known about a patient. For example, a Centor Score in the range of low or moderate probability will be interpreted differently if the clinician knows that a household member had been diagnosed with streptococcal pharyngitis the previous week. CDRs can also help physicians, especially those in training, learn what is important in a history and physical examination and what is not, even if they only integrate these signs and symptoms implicitly (ie, clinical gestalt) rather than explicitly using a CDR.

Finally, we fully agree with Dr. Poddighi’s advice to always consider pertussis in the differential diagnosis of a patient with prolonged cough or typical signs and symptoms, such as the characteristic whoop, paroxysmal cough, or posttussive retching or vomiting. A previous systematic review by our research group showed that nearly 1 in 5 children with prolonged cough have pertussis.<sup>2</sup>

Mark H. Ebell, MD, MS  
Department of Epidemiology and Biostatistics  
University of Georgia, Athens, GA  
ebell@uga.edu

### References

1. Ebell MH, Marchello C, Callahan M. Clinical diagnosis of bordetella pertussis infection: a systematic review. *J Am Board Fam Med* 2017;30:308–19.
2. Marchello C, Perry A, Thai T, Han DS, Ebell MH. Prevalence of atypical pathogens in patients with cough and community-acquired pneumonia: a meta-analysis. *Ann Fam Med* 2016;14:552–66.

doi: 10.3122/jabfm.2017.05.170318

The above letter was referred to the author of the article in question, who offers the following reply.