

CLINICAL REVIEW

Practical Guidelines for the Recognition and Diagnosis of Dementia

James E. Galvin, MD, MPH, and Carl H. Sadowsky, MD

To date, user-friendly, practical guidelines for dementia have not been available for busy family physicians. However, the growing number of patients with dementia means that primary care physicians will have an increasingly important role in the diagnosis and subsequent management of dementia. This article provides practical guidance for the recognition and diagnosis of dementia and is aimed at family physicians, who are usually the first clinicians to whom patients present with dementia symptoms. Because Alzheimer disease (AD) is the most common form of dementia, this condition is the main focus of this article. We review the pathophysiology of AD and discuss recommended diagnostic protocols and the importance of early diagnosis. An AD diagnostic algorithm is provided, with clearly defined steps for screening and diagnosing AD and assessing daily functioning, behavioral symptoms, and caregiver status. (J Am Board Fam Med 2012;25:367–382.)

Keywords: Dementia, Diagnosis, Diagnostic Techniques and Procedures, Guidelines

Dementia: an Overview

Alzheimer disease (AD) is the most common form of dementia, accounting for approximately 60% of all cases,^{1,2} and it is therefore the main focus of this article. AD destroys brain cells, causing problems with memory, thinking, and behavior severe enough to affect work, family, and social relationships and, eventually, the most basic activities of daily living (ADLs). AD and related disorders are a growing public health problem in the United States, with a prevalence ranging from 3% to 11%

among people aged 65 years and older and from 25% to 47% among those aged older than 85 years.^{3,4} There are an estimated 5.3 million cases of dementia in the United States,⁵ and this number is expected to increase to 18.5 million by 2050.⁶ The number of those afflicted is increasing annually as a result of the aging population. Dementia leads to a high burden of suffering for patients, families, and society, with an annual estimated cost of \$172 billion.⁵

People with dementia usually present first to their family physician, although an estimated 39% present to specialist clinics (neurologists, psychiatrists and geriatricians).⁷ The primary care physician (PCP) is often the first physician to observe patients with possible dementia and often the only physician involved in making the diagnosis.⁸ Because of the key role that PCPs play in the long-term management of elderly individuals with chronic disease, the growing number of patients with dementia will have a significant impact on these health professionals.⁹

The rising number of patients with dementia means that family physicians will have an increasingly important role in recognizing early signs and symptoms of disease, ordering appropriate tests, formally diagnosing, and, finally, treating these patients. However, at present, diagnosing AD can be

This article was externally peer reviewed.

Submitted 4 August 2010; revised 17 October 2011; accepted 19 October 2011.

From the Center of Excellence for Brain Aging, New York University Langone School of Medicine, New York, NY (JEG); Division of Neurology, Nova Southeastern University, Fort Lauderdale, FL, and Premiere Research Institute, Palm Beach Neurology, West Palm Beach, FL (CHS).

Funding: none.

Conflict of interest: Dr Sadowsky has served as a consultant to Novartis Pharmaceuticals Corporation, has served as a speaker for Novartis Pharmaceuticals Corporation and Forest Pharmaceuticals and Accera, and has received honoraria from these companies. James E Galvin is a paid consultant for Pfizer, Eisai, Novartis Pharmaceuticals Corporation, Ortho-McNeil and Forest. Neither Dr Sadowsky nor Dr Galvin received financial support for this manuscript.

Corresponding author: James E. Galvin, MD, MPH, Center of Excellence for Brain Aging, New York University Langone School of Medicine, 145 East 32nd Street, 2nd Floor, New York, NY 10016 (E-mail: James.Galvin@nyumc.org).

challenging. In the early stages, AD can be difficult to distinguish from the decline in cognitive abilities due to normal aging and the mild cognitive impairment (MCI) that often precedes AD.^{10,11} Furthermore, there are insufficient numbers of neurologists to care for all the patients with MCI and dementia, and there are a limited number of dementia specialists available for consultation. It is, therefore, imperative that PCPs learn how to assess their patients for dementia; specialty clinics cannot deal with the numbers of patients—a problem that will only worsen as the population continues to age.

To date, user-friendly, practical guidelines have not been available for busy family physicians. Indeed, studies suggest that family physicians may have a relatively limited knowledge of dementia.^{12,13} Thus, there is a pressing need to facilitate diagnosis, which can be simplified by adherence to clearly defined guidelines. This article provides practical guidance for the recognition and diagnosis of dementia and is aimed at PCPs, who are usually the first clinicians to whom patients present with dementia symptoms.

Pathophysiology of AD

AD is a progressive neurodegenerative disorder that represents the most common form of dementia. The most prominent clinical feature of AD is an early impairment of episodic memory,² which manifests as memory impairment of recent events, unusual repeated omissions, and difficulty learning new information. For example, an individual with AD may ask the same question repeatedly throughout the day, such as what they are going to eat for dinner, or frequently misplace items (glasses, keys, or wallet). The loss of memory, especially for learning and retaining new information, reflects impaired function in the hippocampus and other medial temporal lobe structures, which are sites of early pathologic change.² As the disease progresses, the symptoms often manifest in more persistent language disturbance and difficulties completing more complex tasks of daily living. Patients progress from loss of higher level ADLs, such as the ability to perform financial transactions and drive a car or use public transportation, to abnormalities in the more basic ADLs (eg, personal hygiene, toileting). Behavioral problems frequently develop and include depression, apathy, anxiety, agitation, psychosis (delusions and hallucinations), wandering, and aggression.²

Guidelines for Diagnosis

The diagnosis of AD is usually based on the National Institute of Neurological, Communicative Disorders and Stroke–Alzheimer Disease and Related Disorders Association (NINCDS-ADRDA).¹⁴ According to these guidelines, the diagnosis is classified as definite (clinical diagnosis with histologic confirmation), probable (typical clinical syndrome without histologic confirmation), or possible (atypical clinical features but no alternative diagnosis apparent; no histologic confirmation). The currently accepted criteria support a probabilistic diagnosis of AD within a clinical context where there is no definitive diagnostic biomarker. A definite diagnosis of AD is only made according to the NINCDS-ADRDA criteria when there is histopathologic confirmation of the clinical diagnosis.¹⁵ Autopsy results support the “probable” clinical diagnosis in 86% to 90% of cases.¹⁶

Although the NINCDS-ADRDA criteria for AD are the prevailing diagnostic standards in research, these have now fallen behind the unprecedented growth in the understanding of the disease process.¹⁵ The clinical phenotype of AD is no longer described in exclusionary terms, but it can be characterized more definitively on a phenotypic basis. Distinctive markers of the disease are now recognized, including structural brain changes visible on magnetic resonance imaging with early and extensive involvement of the medial temporal lobe, molecular neuroimaging changes seen with positron emission tomography (PET), and changes in cerebrospinal fluid (CSF) biomarkers.¹⁵ There recently has been intense research interest in characterizing the earliest stages of disease that predate the crossing of the dementia threshold, defined by functional disability. Prodromal AD is the symptomatic predementia phase of AD, generally included in the MCI category; this phase is characterized by symptoms not severe enough to meet currently accepted diagnostic criteria for AD.¹⁵ The presence of at least one biologic footprint of AD should improve the specificity for diagnosis. This concept forms the foundation of the new diagnostic criteria proposed in 2007, which were developed to allow an earlier and more specific AD diagnosis.¹⁵

The proposed criteria move away from the traditional 2-step approach of first identifying dementia according to degree of functional disability and

then specifying its cause.¹⁵ Instead, they aim to define the clinical, biochemical, structural, and metabolic presence of AD at the earliest stages before full-blown dementia. These new criteria are centered on a clinical core of early and significant episodic memory impairment. They stipulate that there also must be at least one or more abnormal biomarkers among structural neuroimaging with magnetic resonance imaging, molecular neuroimaging with PET, and CSF analysis of β -amyloid or τ proteins.¹⁵ For example, a decrease in the β -amyloid (1–42) peptide and an increase in the τ and phospho- τ proteins may be the earliest signs of AD.^{2,10,17} These criteria represent a cultural shift requiring a more biologically focused workup than previous approaches, and their timeliness is highlighted by the development of drugs that are directed at altering pathogenesis, particularly at the production and clearance of β -amyloid.^{2,15}

Although the focus of research recently has shifted to the development of new tools that increase the specificity of the prodromal AD diagnosis,² there are no readily available biomarkers, laboratory tests, or brain imaging techniques for routine use to aid diagnosis.^{18,19} As such, their use has limited application for family physicians at this time. Likewise, there are no genetic markers currently recommended for routine use in the diagnosis of AD, including testing for the apolipoprotein E4 gene or for mutations in amyloid, presenilin, or τ that have been linked to rare forms of familial dementia.¹⁸ It therefore remains necessary to screen patients by taking a diligent approach to assessing cognition and function.

Importance and Challenges of Early Diagnosis

Making a diagnosis of dementia in the early stages can be a clinical challenge. The insidious and variable emergence of dementia symptoms makes recognition of the syndrome problematic, particularly in the primary care setting,²⁰ with the often limited time available for consultation. In addition, physicians need to be wary of patients' ability to hide their symptoms. In the early stages of dementia, accommodation to or denial of changes in cognition, functional ability, mood, or behavior are common coping strategies.²¹ As the person's denial strengthens, the concerns of the family become more pressing, with the physician often caught in between and faced with apparently irreconcilable needs.²¹ Most specialty associations, including the

American Academy of Neurology, the American Geriatrics Society, the American Medical Association, and the American Association of Family Practitioners, encourage physicians to be alert to cognitive problems in older adults and to take steps to identify cognitive disorders at the earliest possible stage. However, clinicians in primary care often fail to screen older adults for AD on a routine basis because of insufficient time, inadequate reimbursement for services, and uncertainty about resources available to facilitate diagnosis.^{22,23} Furthermore, given the wide range of services needed, PCPs often fail to make referrals because of insufficient knowledge about resources.^{23,24}

Although it is recognized that current pharmacologic and behavioral interventions do not prevent eventual disease progression, there is good evidence to show that their use can lead to disease stabilization and delayed progression of cognitive, functional, and behavioral outcomes, and these outcomes may provide improvement in quality of life for the patient and their family.²² Nevertheless, family physicians should counsel patients with AD and their families about realistic expectations of treatment outcomes with these agents, which are likely to be modest,²⁵ and the potential for mild to moderate adverse effects (mostly nausea, vomiting, diarrhea, or all three). Because damage caused by the pathophysiologic mechanisms associated with AD is presumed to be irreversible, early detection of AD offers better prospects for patients with AD and their families and friends. This allows both the patient and family to participate in their care plan and to prepare better for future challenges as a result of the neurodegenerative process^{26,27} because currently available medications, and any future disease-modifying agents, will have the greatest opportunity for providing benefit. In addition, recent evidence suggests that early treatment provides economic benefits both to the patient and caregiver as well as society as a whole.^{28–30} Hence, the early establishment of a diagnosis and subsequent initiation of an appropriate management program can optimize the prognosis for patients with AD.

Despite the benefits of early intervention, dementia remains underdiagnosed; an estimated 50% of primary care patients aged older than 65 years have not been diagnosed by their PCPs. A primary reason cited for the delay in AD diagnosis has been the difficulty in identifying early signs of AD by

both PCPs and the general public.³¹ Given the challenges and benefits of timely and accurate diagnosis of neurodegenerative disorders, physicians and patients alike desire tools that aid in dementia diagnosis as early as possible.³²

AD Diagnosis Algorithm

Diagnosis of dementia is a stepwise process that involves examination of patient history and early warning signs, as well as performance screening, assessment of daily functioning, behavioral problems, and caregiver status, with possible referral to specialist clinics for more thorough assessment (Table 1).

Step 1: Prediagnostic Tests and Early Warning Signs

Before any diagnostic tests are performed, patient history, physical examination, and laboratory findings as well as input from family should be considered because these factors are paramount to the diagnosis of dementia (Table 1).³³ In particular, family physicians should take into account any risk factors for AD that may be present, including older age, lower education, female sex, and family history of AD (Table 1).

There are several conditions that mimic dementia (eg, depression, hypothyroidism) that can be missed if not evaluated and, if detected, are potentially reversible. To diagnose AD, other medical, neurologic, or psychiatric disorders that could account for the impairment in memory and related symptoms must be investigated¹ (Table 1). Furthermore, family physicians need to be aware of the potential for vision and hearing problems that could be contributing in a significant manner to an apparent decline in cognitive function. It also should be noted that patients with AD frequently have comorbid medical conditions (Table 1), the presence of which can further impair patient function; their appropriate treatment can optimize outcomes and minimize excess disability.¹² Laboratory tests are necessary to identify or rule out secondary causes of dementia and coexisting disorders that are common among elderly people.¹ Recommended tests are shown in Table 1.

Even before the onset of cognitive problems, there are several warning signs that can predict those individuals at a higher risk of developing dementia, including increased frequency of patient visits to their family physician before diagnosis.^{34,35} There are also somatic changes that precede the

onset of dementia that may provide important clues, including accelerated weight loss,³⁶ gait disturbances, physical frailty,³⁷ and slowed psychomotor speed³⁶ (Table 1).

Ten Key Warning Signs for AD The Alzheimer's Association lists 10 key warning signs of AD¹⁰:

- Memory loss
- Difficulty performing familiar tasks
- Problems with language
- Disorientation to time and place
- Poor or decreased judgment
- Problems with abstract thought
- Misplacing things
- Changes in mood or behavior
- Changes in personality
- Loss of initiative

In many patients, the most clinically prominent feature of AD is the decline in cognitive function with an early impairment of episodic memory, for example, what the patient cannot recall what they had for breakfast, even with the provision of cues.¹ Memory complaints are thus the most reliable and easiest to test for signs of early AD.

Step 2: Screening Tools Employed by Family Physicians

It is crucial that physicians screen for AD under appropriate conditions with validated screening tools that are suitable for a busy practice. Several screening tools are available for use in primary care and are continually being updated and re-evaluated to provide family physicians with brief, easy to administer, and effective diagnostic tools. All screening tools are readily available to family physicians and can be obtained without incurring costs. Many family physicians may find it helpful to select a few screening tools and become familiar and proficient with them. It should be noted, however, that these screening tools are not specifically diagnostic for AD. Rather, they provide evidence that an impairment exists and a rationale to move to a more formal evaluation for confirmation.

There are 2 basic ways to assess the patient for dementia. One approach is a performance measure in which a test is administered and scored. The score then is compared with a published norm. This method is useful for determining how the patient performs in comparison with other age-

Table 1. Stepwise Approach to Diagnosing and Assessing Alzheimer Disease in Primary Care

Stage	Purpose	Tools to Use/Information to Obtain
Step 1: Prediagnostic tests	Differential diagnosis and determination of coexisting disorders	<p>Risk factors, including age, female sex, apolipoprotein E4 gene, prior head injury, low education, and family history of AD; stroke, obesity, hypertension, hyperlipidemia, hyperhomocysteinemia, diabetes, hyperinsulinemia, and smoking^{10,65,66}</p> <p>Medical history:</p> <ul style="list-style-type: none"> • Other ailments that mimic dementia include normal age-associated memory changes, depression, delirium, drug reactions, vision and hearing problems¹ <p>Key questions to ask:</p> <ul style="list-style-type: none"> • Has the patient had any recent illnesses? • Has the patient used any new prescription or over-the-counter medications that could cause memory loss, such as benzodiazepines, anticholinergic drugs for urinary incontinence • Has the patient used or been exposed to illicit drugs? • Has there been any exposure to environmental toxins, eg, fuels or solvents? • Has the patient had any head injuries recently? • Is there any history of epilepsy? <p>Laboratory/medical tests:</p> <ul style="list-style-type: none"> • Complete blood cell count (to ascertain presence of anemia/infection), glucose and thyroid function tests, serum electrolytes, serum B₁₂ levels (to identify vitamin deficiencies), liver function tests, renal function tests, and urinalysis, if appropriate¹ • Patients with AD frequently have comorbid medical conditions, eg, cardiovascular disease, infection, pulmonary, renal insufficiency, and arthritis.¹³ <p>Early warning signs of preclinical dementia:</p> <ul style="list-style-type: none"> • Increased frequency of patient visits to the PCP prior to diagnosis, over a period up to 5 years prior to the diagnosis^{34,35} • Accelerated weight loss,³⁶ late-life depression, gait disturbances, and physical frailty³⁷
Step 2: Assess performance	Cognitive assessments that help screen for/diagnose AD	<p>Cognitive tests:*</p> <ul style="list-style-type: none"> • MMSE^{12,13} • Mini-Cog^{39,40,42,67} • MoCA⁶⁸ <p>Informant-rated tool:*</p> <ul style="list-style-type: none"> • AD8⁶³
Step 3: Assess daily functioning	Determine level of independence and degree of disability	<p>Daily function assessment tool:*</p> <ul style="list-style-type: none"> • IADL^{48,49}
Step 4: Assess behavioral symptoms	Determine presence and degree of behavioral symptoms	<p>Behavioral assessment tool:*</p> <ul style="list-style-type: none"> • NPI-Q⁵⁷ <p>Assess the patient for drug toxicity and medical psychiatric, psychosocial, or environmental problems that may underlie behavioral changes.¹³</p>
Step 5: Identify caregiver and assess needs	Identify the primary caregiver and assess adequacy of family and other support systems	<p>Identify primary carers and establish collaboration:</p> <ul style="list-style-type: none"> • Family caregivers are central to the PCP's assessment and care of the patient.⁵⁹ • Establish and maintain collaboration with caregivers.^{59,60} • Routinely incorporate caregivers' reports of patients' changes in daily routine, mood, behavior, and sleeping patterns. <p>Assess health of primary caregiver:</p> <ul style="list-style-type: none"> • Regularly monitor the physical and emotional health of the primary caregiver as well as that of the patient.¹³ • The PCP should assess the caregiver themselves or refer them to a psychologist, social worker, or other member of the health care delivery team.

Continued

Table 1. Continued

Stage	Purpose	Tools to Use/Information to Obtain
Special considerations	Identify culture, language, and literacy of patient and caregiver	<p>Culture:</p> <ul style="list-style-type: none"> • Recognize the caregiving patterns of ethnic minority groups, eg, African American and Hispanic families distribute care among several family members, rather than one primary carer. • Ethnic minority groups may place different interpretations on memory and behavioral problems. <p>Language:</p> <ul style="list-style-type: none"> • Be aware of the preferred language of the patient and family.⁶⁹ <p>Literacy:</p> <ul style="list-style-type: none"> • Recognize that paper-and-pencil tests and forms may not work well with diverse patient populations if basic literacy is not present, even when such forms are in the person's native language. • Some experts suggest that patients be tested only on what they reasonably may be expected to know,⁷⁰ eg, a person with little schooling may not know how to do the serial sevens on the MMSE, but may be competent at applying simple math, such as subtraction, when handling monetary transactions.

The initial assessment of cognitive and functional abilities is important to determine a baseline to which future deficits may be compared. Reassessment of the patient every 6 months is recommended, and more often in cases of sudden changes in behavior or increased rate of decline.¹³

*Refer to Table 2 for further details.

AD, Alzheimer disease; AD8, 8-item Ascertain Dementia tool; IADL, instrumental (or intermediate) activities of daily living; MMSE, Mini-Mental State Examination; Mini-Cog, Mini Cognitive Assessment Instrument; MoCA, Montreal Cognitive Assessment; NPI-Q, Neuropsychiatric Inventory Questionnaire; PCP, primary care physician.

matched individuals, and scores can be followed over time; however, unless previous baseline testing was performed, the score is not able to give the clinician information about whether the patient's cognitive abilities have changed, nor whether the impairment interferes with everyday activities. In addition, brief performance measures may be biased by age, race, sex, education, and socioeconomic status. A second approach is an informant interview. An observant informant can provide information about how the patient's cognitive abilities have changed and whether the change interferes with everyday activities. Because each patient serves as his or her own control, these assessments suffer less from biases. However, the limitation may be in finding an informant for older adults who live by themselves or in skilled nursing facilities. A combination approach with an informant and performance measure may improve the ability to detect dementia at the earliest stages.³⁸

Performance-Based Screening Tools

In terms of assessing cognitive function, the Mini-Mental State Examination (MMSE) test has been used frequently for initial assessment of AD, and its

sensitivity increases if a decline of the score over time is taken into account^{7,17} (Table 2). Although the MMSE is quick and easy to administer and can track the overall progression of cognitive decline, it is not considered to be a good test for definitive AD diagnosis.¹ In addition, there are several issues associated with the MMSE, including bias according to age, race, education, and socioeconomic status (Table 2).³⁹

Several screening tools are now available for use in primary care as alternatives to the MMSE^{7,17}; these are being updated continually to provide PCPs with brief, easy to administer, and effective diagnostic tools. The Mini-Cognitive Assessment Instrument (Mini-Cog) has sensitivity and specificity for dementia similar to that of the MMSE (Table 2).⁴⁰ However, the Mini-Cog's brevity may be an advantage when trying to improve recognition of cognitive impairment in primary care.^{40,41} In addition, the Mini-Cog is not associated with the same language or education bias as the MMSE.^{39,42} An example of the Mini-Cog test is provided in Appendix 1.

Newer instruments, such as the Montreal Cognitive Assessment, a screening tool developed to

Table 2. Key Screening and Assessment Tools for Alzheimer Disease Used in Primary Care

	Key Features Relevant to Clinical Practice	Number of Items	Time Required	Max Score	Cutoff	Sensitivity/Specificity (%)
Performance testing/screening tools						
MMSE	<ul style="list-style-type: none"> Covers 6 areas: (1) orientation, (2) registration, (3) attention and calculation, (4) recall, (5) language, and (6) ability to copy a figure^{12,13} Quick and easy to administer Can track the overall progression of cognitive decline, but not a good test for definitive/early AD diagnosis¹ Results biased according to age, race, education, and socioeconomic status³⁹ 	30 items	10 minutes	30	23–26	79/88
Mini-Cog	<ul style="list-style-type: none"> Combines an uncued 3-item recall test with a CDT that serves as a recall distractor Easy to administer and requires no special equipment⁴⁰ Powerful predictor of impaired activities of daily living⁶⁷ Good performance in ethnolinguistically diverse populations⁴² Easier to administer to non-English speakers³⁹ Less biased by low educational status and literacy level than MMSE³⁹ 	2 items	3 minutes	Maximum, 5		76/89
MoCA	<ul style="list-style-type: none"> Cognitive screening tool for detection of MCI in primary care⁶⁸ Addresses frontal/executive functioning Low susceptibility to cultural and educational biases⁴³ High sensitivity and specificity for detecting MCI in those patients who perform within the normal range of the MMSE^{68,71} 	12 items	10 minutes	30	26	Sensitivity of 90% for MCI 100% for mild AD ⁶⁸

Continued

Table 2. Continued

	Key Features Relevant to Clinical Practice	Number of Items	Time Required	Max Score	Cutoff	Sensitivity/ Specificity (%)
AD8	<ul style="list-style-type: none"> • Informant-rated change • Screening interview: brief, sensitive measure of memory, orientation, judgment, and function • Can also be directly administered to the patient as a self-rating tool⁴⁴ • Use of the AD8 with a brief patient assessment, e.g. word list, could improve detection of dementia in the primary setting to 97% for dementia and 91% for MCI⁶³ • More sensitive to early stages of dementia: strong correlation with biological markers of AD, e.g. CSF⁴⁵ 	8 Yes/No	2 to 3 minutes	<ul style="list-style-type: none"> • Scores of 0 to 1 = normal cognition • Scores of ≥ 2 = probable cognitive impairment 		74–80/80–86
Daily functioning assessment						
IADL	<ul style="list-style-type: none"> • Evaluates patient's ability to perform more complex activities that are necessary for optimal independent functioning^{48,49} • Data collected from the patient or informant (e.g. family member, caregiver) 	7	10–15 minutes	<p>Three choices of answer per question:</p> <p>I = Patient is completely independent A = Assistance required D = Dependent on help for each activity</p>		N/A
Behavioral assessment						
NPI-Q	<ul style="list-style-type: none"> • Rates frequency and severity of behavioral symptoms commonly seen in dementia, and caregiver distress⁵⁷ • Useful tool for family physicians; assesses both severity of symptoms in patient and distress caused to caregiver 	12 symptoms/questions	5 minutes	<p>Each question is scored as either present or absent</p> <p>Present behaviors are scored for severity and distress:</p> <ul style="list-style-type: none"> • Severity: 1 = mild to 3 = severe • Distress: 0 = no distress; 5 = extremely distressing 		N/A

AD, Alzheimer disease; AD8, 8-item Acertain Dementia tool; CDT, clock-drawing test; CSF, cerebrospinal fluid; IADL, instrumental (or intermediate) activities of daily living; MCI, mild cognitive impairment; Mimi-Cog, Mini-Cognitive Assessment Instrument; MMSE, Mini-Mental State Examination; MoCA, Montreal Cognitive Assessment; NPI-Q, Neuropsychiatric Inventory Questionnaire; N/A, not applicable; PCP, primary care physician.

assist PCPs in detecting MCI, are gaining credibility because of improvements in sensitivity and decreasing susceptibility to cultural and educational biases (Table 2).⁴³ Although more complex than the MMSE and Mini-Cog, the Montreal Cognitive Assessment offers the advantage of testing multiple cognitive domains with an easy scoring system and is free for clinical use (www.mocatest.org). There are many other short and simple memory tests available that can be used as first-line screening tools for use in primary care.¹⁷ Each has its pros and cons; the important point is that an objective measurement can provide an accurate “snapshot” of the patient’s cognitive abilities and permits a quantitative measurement to follow for evidence of a treatment response.

Informant-Based Screening Tools

Another key test used in primary care is the 8-item Ascertain Dementia (AD8) screening interview, which is a brief, sensitive measure that reliably differentiates between individuals with and without dementia by querying memory, orientation, judgment, and function (Table 2). The AD8 can be completed by the informant in the waiting room before the office visit. In the absence of an informant, the AD8 also can be completed by the patient as a self-rating tool.⁴⁴ Because the diagnosis of dementia requires a cognitive deficit that represents a change from premorbid abilities and interferes with social and occupational functioning, an informant assessment provides information that a performance measure cannot. Informant interviews such as the AD8 may be more sensitive to early stages of dementia and have strong correlation with biological markers of AD such as CSF and PET studies.⁴⁵ An example of the AD8 test is provided in Appendix 2.

Of the tests described above, the Mini-Cog and AD8 are recommended as key diagnostic tools to use in primary care because they are brief, valid, and reliable instruments that are easy to administer, clinically acceptable, and effective. Both have psychometric properties superior to the MMSE; as such, they are clinically and psychometrically robust and, it has been argued, are more appropriate for routine use in primary care.^{27,46}

Step 3: Assessment of Daily Functioning

An assessment of daily function is vital to determine the extent of the patient’s disability and de-

pendence on the caregiver, the results of which help to enable planning to maximize patients’ independence.⁴⁷ Basic ADLs, such as feeding and toileting, can be assessed with an interview or by using a tool such as the ADL Scale. Assessment of instrumental ADLs (IADLs) addresses more advanced activities, such as shopping, cooking, and managing finances. The IADL scale is used most frequently and measures 7 areas of more complex activities required for optimal independent functioning^{48,49} (Table 2; Appendix 3). The cognitive changes commonly associated with AD first impact the instrumental and, eventually, the basic ADLs.^{50–52} Another commonly used scale is the Functional Assessment Questionnaire (FAQ),⁵³ which includes 10 items and has been developed from the IADL scale. It assesses shopping, handling finances, preparing a meal, and traveling (which are also in the IADL scale); remembering appointments; and paying attention to, understanding, and discussing television, a book, or a magazine. The total score ranges from 0 (independent) to 30 (dependent).⁵³ An example of the FAQ is provided in Appendix 4.

Step 4: Assessment of Behavioral Symptoms, Psychotic Symptoms, and Depression

More than 80% of patients with AD experience some form of behavioral symptoms such as anxiety, agitation, depression, and apathy during the course of the disease,^{54–56} and patients should, therefore, be assessed periodically. Although these symptoms may be observed by the family physician, they are more often reported by the primary caregiver.¹³

Standardized tools can be used by PCPs or clinic staff to gather information about behavioral symptoms from the caregiver and evaluate effectiveness of interventions over time. The Neuropsychiatric Inventory Questionnaire is a quickly administered instrument that provides reliable assessment of behaviors commonly observed in patients with dementia⁵⁷ (Table 2; Appendix 5). In addition, the Behavioral Pathology in Alzheimer’s Disease (BEHAVE-AD) Rating Scale was designed particularly to be useful in prospective studies of behavioral symptoms and in pharmacologic trials to document behavioral symptoms in patients with AD.⁵⁸ The BEHAVE-AD Rating Scale has 2 parts: the first concentrates on symptomatology and the second requires a global rating of the symptoms on a 4-point scale of severity. The domains covered are paranoid and delusional ideation, hallucinations,

activity disturbances, aggression, diurnal variation, mood and anxieties, and phobias.⁵⁸

Step 5: Caregiver Needs and Status/Support system

Family caregivers are central to the PCP's assessment and care of the patient with AD,⁵⁹ and establishing and maintaining collaboration with caregivers is critical for care of the AD patient.^{59,60} Indeed, major physician organizations, such as the American Academy of Neurology⁶¹ and the American Association for Geriatric Psychiatry,⁶² have emphasized the importance of family caregivers by encouraging family physicians to form partnerships with families who care for dementia patients (Table 1).

In addition, the physical and emotional health of the primary caregiver is crucial in obtaining optimal care for the AD patient. Caregivers suffer from increased rates of depression and physical illness,¹³ and family physicians need to monitor regularly the health of the primary caregiver as well as that of the patient with AD (Table 1). Assessing caregiver status can lead to the implementation of measures that minimize patient-caregiver stress and delay institutionalization of the patient.

Additional Considerations

The family physician should be aware of the need for cultural, language, and literacy assessment within the clinical evaluation process for AD. Identifying the patient's and family's culture, values, primary language, literacy level, and decision-making processes will enable optimal assessment and management of AD patients and their families (Table 1).

Referral: Role of the Specialist and Further Investigations

Difficulty making a diagnosis may necessitate referral to a specialist, such as a geriatrician, neurologist, or psychiatrist, or require the request for further investigations to be completed by a neuropsychologist. For those with mild or questionable impairment, more comprehensive standardized cognitive assessments can be useful to assist with establishing a firm diagnosis; these are normally undertaken as part of a specialist assessment after referral.⁷

Conclusion

The growing number of patients with dementia means that PCPs will have an increasingly impor-

tant role in the diagnosis and subsequent management of disease. There has been unprecedented growth of scientific knowledge about AD and a subsequent move toward its earlier diagnosis. However, in the absence of biomarkers and brain imaging that can be used routinely in primary care, the emphasis remains on the family physician's ability to recognize and diagnose AD using performance screening tools. Of the newly available screening tools for use by PCPs, the Mini-Cog and AD8 are particularly useful as complementary, brief, easy to administer, and effective diagnostic assessments that can be used in everyday clinical practice. Alongside cognitive and daily functioning assessments, a thorough evaluation of behavioral symptoms and caregiver status is required to ensure that both the patient and the patient's family receive optimal care.

The authors would like to thank Frances Gambling for her editorial assistance with the manuscript. Administrative, editorial, and technical assistance was funded by Novartis Pharmaceuticals Corporation.

References

- deSouza L, Sarazin M, Goetz C, Dubois B. Clinical investigations in primary care. *Front Neurol Neurosci* 2009;24:1-11.
- Dubois B, Picard G, Sarazin M. Early detection of Alzheimer's disease: new diagnostic criteria. *Dialogues Clin Neurosci* 2009;11:135-9.
- Boustani M, Callahan CM, Unverzagt FW, et al. Implementing a screening and diagnosis program for dementia in primary care. *J Gen Intern Med* 2005; 20:572-7.
- Boustani M, Peterson B, Hanson L, Harris R, Lohr KN. Screening for dementia in primary care: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2003;138:927-37.
- Alzheimer's Association. 2010 Alzheimer disease facts and figures. *Alzheimer's & Dementia*. Vol. 6. Available from: http://www.alz.org/documents_custom/report_alzfactsfigures2010.pdf. Accessed 14 March 2012.
- Sloane PD, Zimmerman S, Suchindran C, et al. The public health impact of Alzheimer's disease, 2000-2050: potential implication of treatment advances. *Annu Rev Public Health* 2002;23:213-31.
- National Institutes of Health and Clinical Excellence. Dementia: supporting people with dementia and their carers in health and social care. Available from: <http://www.nice.org.uk/guidance/cg42>. Accessed 27 March 2012.
- van Hout HP, Vernooij-Dassen MJ, Stalman WA. Diagnosing dementia with confidence by GPs. *Fam Pract* 2007;24:616-21.

9. Langa KM, Chernew ME, Kabeto MU, et al. National estimates of the quantity and cost of informal caregiving for the elderly with dementia. *J Gen Intern Med* 2001;16:770–8.
10. Salloway S, Correia S. Alzheimer disease: time to improve its diagnosis and treatment. *Cleve Clin J Med* 2009;76:49–58.
11. Morris JC, Storandt M, Miller JP, et al. Mild cognitive impairment represents early-stage Alzheimer disease. *Arch Neurol* 2001;58:397–405.
12. Cummings JL, Frank JC, Cherry D, et al. Guidelines for managing Alzheimer's disease: part II. Treatment. *Am Fam Physician* 2002;65:2525–34.
13. Cummings JL, Frank JC, Cherry D, et al. Guidelines for managing Alzheimer's disease: part I. Assessment. *Am Fam Physician* 2002;65:2263–72.
14. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;34:939–44.
15. Dubois B, Feldman HH, Jacova C, et al. Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol* 2007;6:734–46.
16. Galasko D, Hansen LA, Katzman R, et al. Clinical-neuropathological correlations in Alzheimer's disease and related dementias. *Arch Neurol* 1994;51:888–95.
17. Waldemar G, Dubois B, Emre M, et al. Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *Eur J Neurol* 2007;14:e1–26.
18. Knopman DS, DeKosky ST, Cummings JL, et al. Practice parameter: diagnosis of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001;56:1143–53.
19. Hye A, Lynham S, Thambisetty M, et al. Proteome-based plasma biomarkers for Alzheimer's disease. *Brain* 2006;129:3042–50.
20. Iliffe S, Robinson L, Brayne C, et al. Primary care and dementia: 1. diagnosis, screening and disclosure. *Int J Geriatr Psychiatry* 2009;24:895–901.
21. Woods RT, Moniz-Cook E, Iliffe S, et al. Dementia: issues in early recognition and intervention in primary care. *J R Soc Med* 2003;96:320–4.
22. Cotter VT. Alzheimer's disease: issues and challenges in primary care. *Nurs Clin North Am* 2006;41:83–93, vi.
23. Hinton L, Franz CE, Reddy G, Flores Y, Kravitz RL, Barker JC. Practice constraints, behavioral problems, and dementia care: primary care physicians' perspectives. *J Gen Intern Med* 2007;22:1487–92.
24. Reuben DB, Roth C, Kamberg C, Wenger NS. Restructuring primary care practices to manage geriatric syndromes: the ACOVE-2 intervention. *J Am Geriatr Soc* 2003;51:1787–93.
25. Kaduszkiewicz H, Zimmermann T, Beck-Bornholdt HP, van den Bussche H. Cholinesterase inhibitors for patients with Alzheimer's disease: systematic review of randomised clinical trials. *BMJ* 2005;331:321–7.
26. Vestegaard M, Kerman K, Tamiya E. The study of Alzheimer's disease biomarkers: current role and future prospects of nanosensor technology. *Nanobiotechnology* 2006;2:5–16.
27. Milne A, Culverwell A, Guss R, Tuppen J, Whelton R. Screening for dementia in primary care: a review of the use, efficacy and quality of measures. *Int Psychogeriatr* 2008;20:911–26.
28. Winblad B, Wimo A. Assessing the societal impact of acetylcholinesterase inhibitor therapies. *Alzheimer Dis Assoc Disord* 1999;13(Suppl 2):S9–19.
29. Wimo A. Cost effectiveness of cholinesterase inhibitors in the treatment of Alzheimer's disease: a review with methodological considerations. *Drugs Aging* 2004;21:279–95.
30. Mesterton J, Wimo A, By A, Langworth S, Winblad B, Jonsson L. Cross-sectional observational study on the societal costs of Alzheimer's disease. *Curr Alzheimer Res* 2010;7:358–67.
31. Wilkinson D, Sganga A, Stave C, O'Connell B. Implications of the Facing Dementia Survey for health care professionals across Europe. *Int J Clin Pract Suppl* 2005;27–31.
32. Weaver J, Espinoza R, Weintraub N. The utility of PET brain imaging in the initial evaluation of dementia. *J Am Med Dir Assoc* 2007;8:150–7.
33. Christensen DD, Lin P. Practical treatment strategies for patients with Alzheimer's disease. *J Fam Pract* 2007;56:S17–23.
34. Bamford C, Eccles M, Steen N, Robinson L. Can primary care record review facilitate earlier diagnosis of dementia? *Fam Pract* 2007;24:108–16.
35. Ramakers IH, Visser PJ, Aalten P, et al. Symptoms of preclinical dementia in general practice up to five years before dementia diagnosis. *Dement Geriatr Cogn Disord* 2007;24:300–6.
36. Johnson DK, Wilkins CH, Morris JC. Accelerated weight loss may precede diagnosis in Alzheimer disease. *Arch Neurol* 2006;63:1312–7.
37. Buchman AS, Schneider JA, Leurgans S, Bennett DA. Physical frailty in older persons is associated with Alzheimer disease pathology. *Neurology* 2008;71:499–504.
38. Galvin JE, Powlishta KK, Wilkins K, et al. Predictors of preclinical Alzheimer disease and dementia: a clinicopathologic study. *Arch Neurol* 2005;62:758–65.
39. Borson S, Scanlan JM, Watanabe J, Tu SP, Lessig M. Simplifying detection of cognitive impairment: comparison of the Mini-Cog and Mini-Mental State

- Examination in a multiethnic sample. *J Am Geriatr Soc* 2005;53:871–4.
40. Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The mini-cog: a cognitive ‘vital signs’ measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry* 2000;15:1021–7.
 41. Borson S, Scanlan JM, Watanabe J, Tu SP, Lessig M. Improving identification of cognitive impairment in primary care. *Int J Geriatr Psychiatry* 2006;21:349–55.
 42. Borson S, Scanlan JM, Chen P, Ganguli M. The Mini-Cog as a screen for dementia: validation in a population-based sample. *J Am Geriatr Soc* 2003;51:1451–4.
 43. Ismail Z, Rajji TK, Shulman KI. Brief cognitive screening instruments: an update. *Int J Geriatr Psychiatry* 2010;25:111–20.
 44. Galvin JE, Roe CM, Coats MA, Morris JC. Patient’s rating of cognitive ability: using the AD8, a brief informant interview, as a self-rating tool to detect dementia. *Arch Neurol* 2007;64:725–30.
 45. Galvin JE, Fagan AM, Holtzman DM, Mintun MA, Morris JC. Relationship of dementia screening tests with biomarkers of Alzheimer’s disease. *Brain* 2010;133:3290–300.
 46. Borson S, Scanlan J, Hummel J, Gibbs K, Lessig M, Zuhr E. Implementing routine cognitive screening of older adults in primary care: process and impact on physician behavior. *J Gen Intern Med* 2007;22:811–7.
 47. Holmes SB, Adler D. Dementia care: critical interactions among primary care physicians, patients and caregivers. *Prim Care* 2005;32:671–82, vi.
 48. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969;9:179–86.
 49. Barberger-Gateau P, Commenges D, Gagnon M, Letenneur L, Sauvel C, Dartigues JF. Instrumental activities of daily living as a screening tool for cognitive impairment and dementia in elderly community dwellers. *J Am Geriatr Soc* 1992;40:1129–34.
 50. Fitz AG, Teri L. Depression, cognition, and functional ability in patients with Alzheimer’s disease. *J Am Geriatr Soc* 1994;42:186–91.
 51. Monllau A, Pena-Casanova J, Blesa R, et al. [Diagnostic value and functional correlations of the ADAS-Cog scale in Alzheimer’s disease: data on NORMACODEM project]. *Neurologia* 2007;22:493–501.
 52. Park KW, Pavlik VN, Rountree SD, Darby EJ, Doody RS. Is functional decline necessary for a diagnosis of Alzheimer’s disease? *Dement Geriatr Cogn Disord* 2007;24:375–9.
 53. Juva K, Makela M, Erkinjuntti T, et al. Functional assessment scales in detecting dementia. *Age Ageing* 1997;26:393–400.
 54. Craig D, Mirakhor A, Hart DJ, McIlroy SP, Passmore AP. A cross-sectional study of neuropsychiatric symptoms in 435 patients with Alzheimer’s disease. *Am J Geriatr Psychiatry* 2005;13:460–8.
 55. Steffens DC, Maytan M, Helms MJ, Plassman BL. Prevalence and clinical correlates of neuropsychiatric symptoms in dementia. *Am J Alzheimers Dis Other Demen* 2005;20:367–73.
 56. Lyketsos CG, Lee HB. Diagnosis and treatment of depression in Alzheimer’s disease. A practical update for the clinician. *Dement Geriatr Cogn Disord* 2004;17:55–64.
 57. Kaufer DI, Cummings JL, Ketchel P, et al. Validation of the NPI-Q, a brief clinical form of the Neuropsychiatric Inventory. *J Neuropsychiatry Clin Neurosci* 2000;12:233–9.
 58. Reisberg B, Borenstein J, Salob SP, Ferris SH, Franssen E, Georgotas A. Behavioral symptoms in Alzheimer’s disease: phenomenology and treatment. *J Clin Psychiatry* 1987;48(Suppl):9–15.
 59. Family Caregiver Alliance. Caregiver assessment: principles, guidelines and strategies for change. Report from a National Consensus Development Conference. Vol. 1. San Francisco: Family Caregiver Alliance; 2006.
 60. Bultman DC, Svarstad BL. Effects of physician communication style on client medication beliefs and adherence with antidepressant treatment. *Patient Educ Couns* 2000;40:173–85.
 61. Doody RS, Stevens JC, Beck C, et al. Practice parameter: management of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001;56:1154–66.
 62. Lyketsos CG, Colenda CC, Beck C, et al. Position statement of the American Association for Geriatric Psychiatry regarding principles of care for patients with dementia resulting from Alzheimer disease. *Am J Geriatr Psychiatry* 2006;14:561–72.
 63. Galvin JE, Roe CM, Powlishta KK, et al. The AD8: a brief informant interview to detect dementia. *Neurology* 2005;65:559–64.
 64. Pfeffer RI, Kurosaki TT, Harrah CH Jr, Chance JM, Filos S. Measurement of functional activities in older adults in the community. *J Gerontol* 1982;37:323–9.
 65. Luchsinger JA, Mayeux R. Cardiovascular risk factors and Alzheimer’s disease. *Curr Atheroscler Rep* 2004;6:261–6.
 66. Dunn N, Mullee M, Perry VH, Holmes C. Association between dementia and infectious disease: evidence from a case-control study. *Alzheimer Dis Assoc Disord* 2005;19:91–4.
 67. Scanlan JM, Binkin N, Michieletto F, Lessig M, Zuhr E, Borson S. Cognitive impairment, chronic disease burden, and functional disability: a population study of older Italians. *Am J Geriatr Psychiatry* 2007;15:716–24.

68. Nasreddine ZS, Phillips NA, Bedirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005;53:695–9.

69. Folsom DP, Gilmer T, Barrio C, et al. A longitudinal study of the use of mental health services by persons with serious mental illness: do Spanish-speaking Latinos differ from English-speaking Latinos and Caucasians? *Am J Psychiatry* 2007;164:1173–80.

70. Teng EL, Manly JJ. Neuropsychological testing: helpful or harmful? *Alzheimer Dis Assoc Disord* 2005;19:267–71.

71. Luis CA, Keegan AP, Mullan M. Cross validation of the Montreal Cognitive Assessment in community dwelling older adults residing in the Southeastern US. *Int J Geriatr Psychiatry* 2009;24:197–201.

- Apple
- Watch
- Penny

2. Administer the CDT: ask the patient to draw a traditional clock face showing a time that is potentially confusing for someone with AD, such as “10 minutes past 11.”
3. Ask the patient to repeat the 3 words given previously.

Scoring (number of correct items recalled):
 3 = Negative screen
 1 or 2 plus normal CDT = Negative screen
 1 or 2 plus abnormal CDT = Positive for cognitive impairment

Appendix 1

The Mini-Cog Test

Scoring: 1 point for each recalled word (Figure 1).

Score the clock drawing test (CDT) as *normal* (the patient places the correct time and the clock appears grossly normal) or *abnormal*.


CDT Instructions

A pen/pencil and a blank sheet of paper are required.

1. Instruct the patient to listen carefully and repeat the following:

Figure 1. The Mini-Cog Scoring Algorithm. 0 = Positive for cognitive impairment; 1 or 2 plus an abnormal clock drawing test (CDT) = positive for cognitive impairment; 1 or 2 plus a normal CDT = negative for cognitive impairment; 3 = Negative screen for dementia (no need to score CDT).

Reproduced with permission from Borson S et al. *Int J Geriatr Psychiatry* 2000;15:1021–1027. © 2000 John Wiley & Sons³⁹



Appendix 2

The AD8 Administration and Scoring Guidelines

A spontaneous self-correction is allowed for all responses without counting as an error.

The questions are given to the respondent on a clipboard for self-administration or can be read aloud to the respondent either in person or over the phone (Table 3). It is preferable to administer the AD8 to an informant, if available. If an informant is not available, the AD8 may be administered to the patient.

- When administered to an informant, specifically ask the respondent to rate change in the patient
- When administered to the patient, specifically ask the patient to rate changes in his or her ability for each of the items *without* attributing causality

If read aloud to the respondent, it is important for the clinician to read the phrase carefully as worded and give emphasis to note changes caused by cognitive problems (not physical problems). There should be a 1-second delay between individual items.

- No time frame for change is required
- The final score is a sum of the number items marked “yes, a change”

Interpretation of the AD8⁶³

A screening test in itself is insufficient to diagnose a dementing disorder. The AD8 is, however, quite sensitive to detecting early cognitive changes asso-

Table 3. AD8 Dementia Screening Interview

	Yes, a Change*	No, No Change	N/A, Don't Know
1. Problems with judgment (eg, problems making decisions, bad financial decisions, problems with thinking)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Less interest in hobbies/activities	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Repeats the same things over and over (questions, stories, or statements)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Trouble learning how to use a tool, appliance, or gadget (eg, VCR, computer, microwave, remote control)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Forgets correct month or year	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Trouble handling complicated financial affairs (eg, balancing checkbook, income taxes, paying bills)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. Trouble remembering appointments	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. Daily problems with thinking, memory, or both	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
TOTAL AD8 SCORE	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

*“Yes, a change” indicates that there has been a change in the last several years caused by cognitive (thinking and memory) problems. Scores of 0 to 1 indicate normal cognition; scores of ≥ 2 indicate that cognitive impairment is likely to be present.

AD8, 8-item Ascertain Dementia screening.

Adapted with permission from Galvin J et al. *Neurology* 2006;67:1942–8. © 2006 AAN Enterprises, Inc.⁶³

ciated with many common dementing illnesses including AD, vascular dementia, Lewy body dementia, and frontotemporal dementia.

- Scores in the impaired range (see below) indicate a need for further assessment.
- Scores in the “normal” range suggest that a dementing disorder is unlikely, but an early disease process cannot be ruled out.

More advanced assessment may be warranted in cases where other objective evidence of impairment exists.

The following cut points are used to diagnose dementia:

- 0 or 1: Normal cognition
- ≥ 2 : Cognitive impairment is likely to be present

Appendix 3

Instrumental Activities of Daily Living¹³

This tool evaluates the patient’s ability to perform the more complex activities that are neces-

sary for optimal independent functioning (Table 4). The scoring indicates whether the patient is completely independent (I), requires assistance (A), or is dependent (D) for the performance of each activity. The IADL instrument may be repeated periodically to determine the need for more support.

Appendix 4

Functional Assessment Questionnaire (FAQ)⁶⁴

The FAQ includes 10 items and has been developed from the IADL scale (Table 5). The total score ranges from 0 (independent) to 30 (dependent).

Appendix 5

Neuropsychiatric Inventory Questionnaire¹³

The Neuropsychiatric Inventory Questionnaire provides a reliable assessment of behaviors frequently observed in patients with dementia. (Figure 2).

Table 4. Instrumental Activities of Daily Living Scale

Name of patient		Date.....	
This form may help you assess the functional capabilities of your older patients. The data can be collected by a nurse from the patient or from an informant such as a family member or other caregiver (I = independent; A = assistance required; D = dependent)			
Obtained from		Activity	Guidelines for assessment
Patient	Informant		
I A D	I A D	Using telephone	I = Able to look up numbers, dial telephone, and receive and make calls without help A = Able to answer telephone or dial operator in an emergency, but needs special telephone or help in getting numbers and/or dialing D = Unable to use telephone
I A D	I A D	Traveling	I = Able to drive own car or travel alone on buses or in taxis A = Able to travel, but needs someone to travel with D = Unable to travel
I A D	I A D	Shopping	I = Able to take care of all food and all clothes shopping with transportation provided A = Able to shop, but needs someone to shop with D = Unable to shop
I A D	I A D	Preparing meals	I = Able to plan and cook full meals A = Able to prepare light foods, but unable to cook full meals alone D = Unable to prepare any meals
I A D	I A D	Housework	I = Able to do heavy housework (i.e. scrub floors) A = Able to do light housework, but needs help with heavy tasks D = Unable to do any housework
I A D	I A D	Taking medicine	I = Able to prepare and take medications in the right dose at the right time A = Able to take medications, but needs reminding or someone to prepare them D = Unable to take medications
I A D	I A D	Managing money	I = Able to manage buying needs (i.e. write checks, pay bills) A = Able to manage daily buying needs, but needs help managing checkbook and/or paying bills D = Unable to handle money

Adapted with permission from: Lawton MP, Brody EM. *Gerontologist* 1969;9:179–86. 1969 © The Gerontological Society of America.

Table 5. Functional Assessment Questionnaire

In the Past 4 Weeks, Did the Subject Have Any Difficulty or Need Help With:	Not Applicable	Normal	Has Difficulty, But Does by Self	Requires Assistance	Dependent
1. Writing checks, paying bills or balancing a checkbook	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Assembling tax records, business affairs, or other papers	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Shopping alone for clothes, household necessities, or groceries	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Playing a game of skill such as bridge or chess, working on a hobby	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Heating water, making a cup of coffee, turning off the stove	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Preparing a balanced meal	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. Keeping track of current events	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. Paying attention to and understanding a TV program, book, or magazine	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Remembering appointments, family occasions, holidays, medications	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. Traveling out of the neighborhood, driving, or arranging to take public transportation	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Adapted from Pfeffer RI, Kurosaki TT, Harrah CH Jr, Chance JM, Filos S. Measurement of functional activities in older adults in the community. *J Gerontol* 1982;37:323–9. 1982 © The Gerontological Society of America.

Figure 2. Neuropsychiatric Inventory Questionnaire¹³

Name of patient: _____ Date: _____

Informant: Spouse: _____ Child: _____ Other: _____

Please answer the following questions based on *changes* that have occurred since the patient first began to experience memory problems. Circle "yes" only if the symptom has been present in the *past month*. Otherwise, circle "no".

For each item marked "yes":

Rate the *severity* of the symptom (how it affects the patient):

- 1 = Mild (noticeable, but not a significant change)
- 2 = Moderate (significant, but not a dramatic change)
- 3 = Severe (very marked or prominent; a dramatic change)

Rate the *distress* you experience because of that symptom (how it affects you):

- 0 = Not distressing at all
- 1 = Minimal (slightly distressing, not a problem to cope with)
- 2 = Mild (not very distressing, generally easy to cope with)
- 3 = Moderate (fairly distressing, not always easy to cope with)
- 4 = Severe (very distressing, difficult to cope with)
- 5 = Extreme or very severe (extremely distressing, unable to cope with)

Please answer each question honestly and carefully. Ask for assistance if you are not sure how to answer any question.

Delusions	Does the patient believe that others are stealing from him or her, or planning to harm him or her in some way?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Hallucinations	Does the patient act as if he or she hears voices? Does he or she talk to people who are not there?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Agitation or aggression	Is the patient stubborn and resistive to help from others?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Depression or dysphoria	Does the patient act as if he or she is sad or in low spirits? Does he or she cry?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Anxiety	Does the patient become upset when separated from you? Does he or she have any other signs of nervousness, such as shortness of breath, sighing, being unable to relax, or feeling excessively tense?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Elation or euphoria	Does the patient appear to feel too good or act excessively happy?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Apathy or indifference	Does the patient seem less interested in his or her usual activities and in the activities and plans of others?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Disinhibition	Does the patient seem to act impulsively? For example, does the patient talk to strangers as if he or she knows them, or does the patient say things that may hurt people's feelings?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Irritability or lability	Is the patient impatient and cranky? Does he or she have difficulty coping with delays or waiting for planned activities?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Motor disturbance	Does the patient engage in repetitive activities, such as pacing around the house, handling buttons, wrapping string, or doing other things repeatedly?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Nighttime behaviors	Does the patient awaken you during the night, rise too early in the morning, or take excessive naps during the day?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			
Appetite and eating	Has the patient lost or gained weight, or had a change in the food he or she likes?	Severity: 1 2 3	Distress: 0 1 2 3 4 5
Yes No			

Adapted with permission from: Kaufer DI, Cummings JL, Ketchel P et al. J Neurosychiatry Clin Neurosci 2000; 12:233-9. 2000 © American Psychiatric Press, Inc.⁵⁷