

# Subject Index to Volume 29, 2016

- Academic medical centers**, patient portals and blood pressure control, 29(4):452-459
- Access to health care**  
association with lower obesity rates, 29(2):182-190  
hearing loss health care for older adults, 29(3):394-403  
school-based health centers, 29(3):339-347  
underuse of primary care in China, 29(2):240-247
- Accountable care organizations**, telehealth  
in primary care, 29(4):432-433  
as useful tool, 29(4):430-431
- Acne**, treatment, in pregnancy, 29(2):254-262
- Admission data**, prediction of hospital readmission risk, 29(1):50-59
- Adolescent health**, school-based health centers, 29(3):339-347
- Adult survivors of adverse events**, screening for adverse childhood experiences, 29(3):303-307
- Aging**, reducing functional decline in frail elderly, 29(2):209-217
- Albuminuria**, clinical decision support for chronic kidney disease, 29(5):604-612
- Alcohol-related disorders**, self-report of alcohol treatment referrals, 29(6):682-687
- Alcoholism**, pretreatment use and drinking goal in alcohol treatment, 29(1):37-49
- Algorithms**  
communicating delays in diagnostic evaluation, 29(4):469-473  
triage amalgamated dermoscopic algorithm, 29(6):694-701
- Ambulatory care facilities**  
communicating delays in diagnostic evaluation, 29(4):469-473  
electronic portal use among vulnerable patients, 29(5):592-603  
kiosk technology for blood pressure, 29(5):620-629  
problem drug-related behavior, 29(6):718-726
- American Board of Family Medicine (ABFM)**, election of new officers and board members, 29(5):637-638
- Analgesics**  
PCMH recognition and chronic pain management, 29(4):474-481  
problem drug-related behavior, 29(6):718-726
- Analysis of variance**, academic detailing intervention for cancer screening, 29(5):533-542
- Anemia**, multiple myeloma and diagnostic delay, 29(6):702-709
- Angiotensin-converting enzyme inhibitors**, clinical decision support for chronic kidney disease, 29(5):604-612
- Angiotensin receptor antagonists**, clinical decision support for chronic kidney disease, 29(5):604-612
- Antibacterial agents**  
antibiotic prophylaxis and total joint replacement, 29(4):500-507  
levofloxacin in males with a UTI, 29(6):654-662
- Area under curve**, almond "appetizer" effect on GTT results, 29(6):759-766
- Arthralgia**, PCMH recognition and chronic pain management, 29(4):474-481
- Arthritis**, infectious, antibiotic prophylaxis and total joint replacement, 29(4):500-507
- Asthma**, same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688-693
- Attitudes**, physician, toward diabetes prevention, 29(6):663-671
- Auditory brainstem implants**, pediatric, for habilitation, 29(2):286-288
- Awards and prizes**  
federal research funding for family medicine, 29(5):531-532  
limited NIH funding for family medicine research, 29(5):528-530
- Back pain**, multiple myeloma and diagnostic delay, 29(6):702-709
- Bacterial infections**, nonbacterial causes of lymphangitis, 29(6):808-812
- Basal cell carcinoma**, diagnostic maneuver to increase early detection, 29(3):404-407
- Behavior therapy**  
childhood obesity intervention, 29(4):434-443  
physician attitudes toward diabetes prevention, 29(6):663-671
- Behavioral medicine**  
integrating behavioral health using lean workflow analysis, 29(3):385-393  
provider diversity on the family medicine team, 29(1):8-9
- Biomarkers**, fertility apps marketed to avoid pregnancy, 29(4):508-511
- Biopsy**, triage amalgamated dermoscopic algorithm, 29(6):694-701
- Birth**, doula support during pregnancy, 29(3):308-317
- Birth rate**, fertility apps marketed to avoid pregnancy, 29(4):508-511
- Bisexual persons**, improving patient-centered "LGBTQ" primary care, 29(1):156-160
- Bisphosphonates**, insufficiency fractures and bisphosphonate therapy, 29(3):404-407
- Blood glucose**, almond "appetizer" effect on GTT results, 29(6):759-766
- Blood pressure**  
childhood obesity intervention, 29(4):434-443  
control, patient portals and, 29(4):452-459  
kiosk technology, 29(5):620-629  
patient portals and blood pressure control, 29(4):452-459
- Blue Cross Blue Shield insurance plans**, medical home implementation in small practices, 29(6):767-774
- Body mass index**  
childhood obesity intervention, 29(4):434-443  
participation in weight loss programs, 29(5):572-580
- Bone fractures**  
distal radius fracture treatment, 29(2):218-225  
insufficiency fractures and bisphosphonate therapy, 29(3):404-407
- Breast cancer**, academic detailing intervention for cancer screening, 29(5):533-542
- Breastfeeding**, diagnosis and management of breast milk oversupply, 29(1):139-142
- C-reactive protein**, level, as marker in febrile children, 29(4):460-468
- California**, FIT for colon cancer screening, 29(6):672-681
- Cancer prevention and control**, academic detailing intervention for cancer screening, 29(5):533-542
- Cancer screening**  
academic detailing intervention, 29(5):533-542  
basal cell blanché, 29(3):408-410  
FIT for colon cancer screening, 29(6):672-681  
lung cancer screening guidelines, 29(1):152-155  
PCMH increased colorectal cancer screening, 29(2):191-200
- Carcinoma**, basal cell blanché, 29(3):408-410
- Cardiovascular diseases**, PCMH impact on chronically ill patients' utilization, 29(4):482-495
- Care coordination**, in PCMHs, 29(1):90-101
- Career choice**  
family medicine residents and educational debt, 29(2):180-181  
impact of debt on young family physicians, 29(2):177-179  
provider diversity on the family medicine team, 29(1):8-9
- Case reports**  
insufficiency fractures and bisphosphonate therapy, 29(3):404-407  
vanished twins and misdiagnosed sex, 29(3):411-413
- Centers for Disease Control and Prevention (U.S.)**, federal re-

- search funding for family medicine, 29(5):531–532
- Certification**, care coordinator prevalence in practices and PCMHs, 29(6):652–653
- Cervical cancer**, academic detailing intervention for cancer screening, 29(5):533–542
- Child abuse**, screening for adverse childhood experiences, 29(3):303–307
- Childbirth**, doula support during pregnancy, 29(3):308–317
- Children**, see Pediatrics
- China**, underuse of primary care, 29(2):240–247
- Choice behavior**, physician use of low-value clinical services, 29(6):785–792
- Chronic disease**  
 clinical decision support for chronic kidney disease, 29(5):604–612  
 local learning collaboratives for chronic kidney disease, 29(5):543–552  
 patient portals and blood pressure control, 29(4):452–459  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495  
 PCMH recognition and chronic pain management, 29(4):474–481  
 same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Chronic pain**, management, PCMH recognition and, 29(4):474–481
- Ciprofloxacin**, levofloxacin in males with a UTI, 29(6):654–662
- Clinical coding**, simulation of ICD-9 to ICD-10-CM transition, 29(1):29–36
- Clinical decision support systems**, for chronic kidney disease, 29(5):604–612
- Clinical medicine**  
 hearing loss health care for older adults, 29(3):394–403  
 social determinants of health in a clinic setting, 29(3):414–418
- Cochlear implants**, pediatric auditory brainstem implant for habilitation, 29(2):286–288
- Cohort studies**  
 C-reactive protein level as marker in febrile children, 29(4):460–468  
 FIT for colon cancer screening, 29(6):672–681
- Collaborative care management**, in primary care, time to remission for depression with, 29(1):10–17
- Colon cancer**, PCMH increased colorectal cancer screening, 29(2):191–200
- Colorado**, childhood obesity intervention, 29(4):434–443
- Colorectal cancer**  
 academic detailing intervention for cancer screening, 29(5):533–542  
 FIT for colon cancer screening, 29(6):672–681  
 PCMH increased colorectal cancer screening, 29(2):191–200
- Commentary**  
 Are We Learning More about Patient-centered Medical Homes (PCMHs), or Learning More about Primary Care?, 29(1):4–7  
 Care Coordination for Primary Care Practice, 29(6):649–651  
 Evidence, Engagement, and Technology: Themes of and the State of Primary Care Practice-based Network Research, 29(5):521–524  
 The Impact of Debt on Young Family Physicians: Unanswered Questions with Critical Implications, 29(2):177–179  
 The Need to Systematically Evaluate Clinical Practice Guidelines, 29(6):644–648  
 Social Determinants of Health and Primary Care: Intentionality Is Key to the Data We Collect and the Interventions We Pursue, 29(3):297–300  
 Social Justice as the Moral Core of Family Medicine: A Perspective from the Keystone IV Conference, 29(4):S69–S71  
 Telehealth: A Very Useful Tool That Enables and Improves Patient Access, 29(4):430–431  
 Where You Stand Is What You See: We See a Need for More Primary Care Research Funding, 29(5):525–527
- Communicable diseases**, preexposure prophylaxis for HIV prevention, 29(1):143–151
- Communication**  
 communicating delays in diagnostic evaluation, 29(4):469–473  
 electronic messaging and number of incoming telephone calls, 29(5):613–619  
 watchful waiting and diagnostic testing, 29(6):710–717
- Community-based participatory research**, using patient navigators to improve diabetes outcome, 29(1):78–89
- Community health centers**  
 patients with diabetes served by safety-net centers, 29(3):356–370  
 primary care continuity improves diabetic health, 29(3):318–324
- Community medicine**, school-based health centers, 29(3):339–347
- Comorbidity**  
 admission data predict hospital readmission risk, 29(1):50–59  
 patient portals and blood pressure control, 29(4):452–459  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495
- Comparative effectiveness research**, stakeholder engagement in patient-reported outcomes measure implementation, 29(1):102–115
- Comprehensive health care**, primary care panel size, 29(4):496–499
- Consensus**, evaluating Choosing Wisely™ using SORT, 29(4):512–515
- Continuity of patient care**, continuity of care and reducing emergency department visits, 29(2):201–208
- Contraception**  
 fertility apps marketed to avoid pregnancy, 29(4):508–511  
 IUD use based on who initiated discussion of placement, 29(1):24–28
- Control groups**, communicating delays in diagnostic evaluation, 29(4):469–473
- Cooperative behavior**, local learning collaboratives for chronic kidney disease, 29(5):543–552
- Correspondence**  
 clinical decisions made in primary care clinics before and after Choosing Wisely™, 29(1):167–168  
 the diversity of providers on the family medicine team, 29(3):424  
 electronic health record challenges, workarounds, and solutions observed in practices integrating behavioral health and primary care, 29(2):289–290  
 over half of graduating family medicine residents report more than \$150,000 in educational debt, 29(4):516–517, 29(6):814  
 patient preferences for receiving reports of test results, 29(2):289–290  
 real life is full of choices, 29(5):636  
 reporting and using near-miss events to improve patient safety in diverse primary care practices: a collaborative approach to learning from our mistakes, 29(1):165–167  
 social justice as the moral core of family medicine: a perspective from the Keystone IV Conference, 29(6):813  
 the use of medical scribes in health care settings: a systematic review and future direction, 29(3):423–424
- Cost control**, cost for level 3 medical home recognition, 29(1):69–77
- Cost-effectiveness**  
 distal radius fracture treatment, 29(2):218–225  
 hearing loss health care for older adults, 29(3):394–403
- Cotton fever**, self-diagnosis by IV drug users, 29(2):276–279
- Counseling**, watchful waiting and diagnostic testing, 29(6):710–717
- Creatinine**, multiple myeloma and diagnostic delay, 29(6):702–709
- Cross-sectional studies**  
 participation in weight loss programs, 29(5):572–580  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495  
 triage amalgamated dermoscopic algorithm, 29(6):694–701
- Cultural diversity**, doula support during pregnancy, 29(3):308–317

- Deafness**, pediatric auditory brainstem implant for habilitation, 29(2): 286–288
- Decision making**  
 decision-to-implement worksheet, 29(5):553–562  
 lung cancer screening guidelines, 29(1):152–155  
 parents' and pediatricians' views of partnership, 29(5):563–571
- Decision support techniques**, technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Delivery of health care**  
 care coordination in PCMHs, 29(1): 90–101  
 care coordinator prevalence in practices and PCMHs, 29(6):652–653  
 clinical risk tools for predicting osteoporosis, 29(2):233–239  
 integrating behavioral health using lean workflow analysis, 29(3): 385–393  
 reducing functional decline in frail elderly, 29(2):209–217  
 school-based health centers, 29(3): 339–347  
 strategies for stakeholders to improve EHRs, 29(1):126–134
- Demography**, self-report of alcohol treatment referrals, 29(6):682–687
- Depression**  
 community engagement in impoverished communities, 29(3):325–338  
 family medicine: bridge to life, 29(1): 161–164  
 same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693  
 time to remission with CCM in primary care, 29(1):10–17
- Dermatology**  
 basal cell blanché, 29(3):408–410  
 treatment of acne in pregnancy, 29(2):254–262  
 triage amalgamated dermoscopic algorithm, 29(6):694–701
- Dermoscopy**, triage amalgamated dermoscopic algorithm, 29(6):694–701
- Diabetes mellitus**  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495  
 prediabetes diagnosis and treatment, 29(2):283–285  
 prevention, physician attitudes toward, 29(6):663–671  
 primary care continuity improves diabetic health, 29(3):318–324  
 primary care panel size, 29(4):444–451  
 referrals for diabetes education in a medical home, 29(3):377–384  
 same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693  
 using patient navigators to improve outcome, 29(1):78–89
- Diagnosis**, differential, nonbacterial causes of lymphangitis, 29(6): 808–812
- Diagnostic errors**, nonbacterial causes of lymphangitis, 29(6):808–812
- Diagnostic tests**, watchful waiting and diagnostic testing, 29(6):710–717
- Disclosure**, same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Disease management**  
 patient portals and blood pressure control, 29(4):452–459  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495
- Doctor-patient relations**  
 family medicine: bridge to life, 29(1): 161–164  
 IUD use based on who initiated discussion of placement, 29(1): 24–28
- Doctoring**  
 complexity of “place” in primary care, 29(4):S24–S27  
 countercultural heritage of family medicine, 29(4):S45–S48  
 family medicine as a disruptive innovation, 29(4):S60–S63  
 Generation III family physicians, 29(4):S49–S53  
 holding on and letting go, 29(4):S32–S39  
 impact of time and timing, 29(4): S28–S31  
 people are primary, 29(4):S40–S44  
 personal doctoring manifesto, 29(4): S64–S68  
 recharging family medicine, 29(4): S15–S18  
 reimagining relationships with patients, 29(4):S1–S11  
 relationships in tomorrow's health system, 29(4):S54–S59  
 seeking relationships in primary care, 29(4):S19–S23  
 social justice and family medicine, 29(4):S69–S71  
 tribute to G. Gayle Stephens, 29(4): S12–S14
- Double-blind method**, levofloxacin in males with a UTI, 29(6):654–662
- Early detection of cancer**  
 basal cell blanché, 29(3):408–410  
 FIT for colon cancer screening, 29(6):672–681  
 multiple myeloma and diagnostic delay, 29(6):702–709
- Eating**, almond “appetizer” effect on GTT results, 29(6):759–766
- Editorial Office News and Notes**, Peer Reviewers for the Journal of the American Board of Family Medicine in 2015, 29(2):173–176
- Editorials**, clinical practice guidelines—is “regulation” the answer?, 29(6):642–643
- Editor's Notes**  
 Bread and Butter of Family Medicine: Guidelines, Population Screening, Diagnostic Evaluations, and Practice Models, 29(6):639–641
- Family Medicine Research That Provides Compelling, Urgent Data to Improve Patient Care, 29(1): 1–3
- Outcomes of Health System Structures, Highly Pertinent Clinical Information, Idea Stimulators, Clinical Reviews, and Prediction Tools: JABFM Exemplified, 29(2):171–172
- Research in Family Medicine by Family Physicians for the Practice of Family Medicine, 29(4): 427–429
- Social Determinants of Health and Beyond: Information to Help Family Physicians Improve Patient Care, 29(3):295–296
- Education**  
 family medicine residents and educational debt, 29(2):180–181  
 impact of debt on young family physicians, 29(2):177–179  
 referrals for diabetes education in a medical home, 29(3):377–384
- Elderly**  
 exercising, avoiding myalgia with statins, 29(6):727–740  
 frail, reducing functional decline, 29(2):209–217
- Electronic cigarettes**, use, physician advice for, 29(6):741–747
- Electronic health records (EHRs)**  
 communicating delays in diagnostic evaluation, 29(4):469–473  
 electronic messaging and number of incoming telephone calls, 29(5): 613–619  
 electronic portal use among vulnerable patients, 29(5):592–603  
 multiple myeloma and diagnostic delay, 29(6):702–709  
 patient portals and blood pressure control, 29(4):452–459  
 simulation of ICD-9 to ICD-10-CM transition, 29(1):29–36  
 strategies for stakeholders to improve EHRs, 29(1):126–134  
 technology-based patient engagement strategies within PBRNs, 29(5): 581–591
- Electronic mail**, communicating delays in diagnostic evaluation, 29(4): 469–473
- Emergency departments**  
 C-reactive protein level as marker in febrile children, 29(4):460–468  
 continuity of care and reducing emergency department visits, 29(2):201–208  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495
- Empathy**, PCMH recognition and chronic pain management, 29(4): 474–481
- Energy intake**, almond “appetizer” effect on GTT results, 29(6): 759–766



- Epidemiology**  
reducing health inequities using sociodemographic data, 29(3):348–355  
social determinants of health and primary care, 29(3):297–300
- Erythema**, nonbacterial causes of lymphangitis, 29(6):808–812
- Ethnic groups**, participation in weight loss programs, 29(5):572–580
- Evidence-based medicine**  
decision-to-implement worksheet, 29(5):553–562  
distal radius fracture treatment, 29(2):218–225  
participation in weight loss programs, 29(5):572–580
- Exercise**, avoiding myalgia with statins, 29(6):727–740
- Family medicine**  
care coordinator prevalence in practices and PCMHs, 29(6):652–653  
complexity of “place” in primary care, 29(4):S24–S27  
countercultural heritage of, 29(4):S45–S48  
as a disruptive innovation, 29(4):S60–S63  
federal research funding, 29(5):531–532  
holding on and letting go, 29(4):S32–S39  
impact of time and timing, 29(4):S28–S31  
people are primary, 29(4):S40–S44  
provider diversity on the family medicine team, 29(1):8–9  
recharging, 29(4):S15–S18  
reflections: bridge to life, 29(1):161–164  
relationships in tomorrow’s health system, 29(4):S54–S59  
research, limited NIH funding for, 29(5):528–530  
residents and educational debt, 29(2):180–181  
seeking relationships in primary care, 29(4):S19–S23  
social justice and, 29(4):S69–S71  
team, provider diversity on, 29(1):8–9  
tribute to G. Gayle Stephens, 29(4):S12–S14
- Family physicians**  
complexity of “place” in primary care, 29(4):S24–S27  
Generation III, 29(4):S49–S53  
holding on and letting go, 29(4):S32–S39  
impact of time and timing, 29(4):S28–S31  
institutions with fewer residencies produce more physicians, 29(3):301–302  
people are primary, 29(4):S40–S44  
personal doctoring manifesto, 29(4):S64–S68  
physician attitudes toward diabetes prevention, 29(6):663–671
- Pisacano scholars’ reflections from the Starfield Summit, 29(6):793–804  
primary care panel size, 29(4):444–451  
reimagining relationships with patients, 29(4):S1–S11  
relationships in tomorrow’s health system, 29(4):S54–S59  
seeking relationships in primary care, 29(4):S19–S23  
telehealth as useful tool, 29(4):430–431  
triage amalgamated dermoscopic algorithm, 29(6):694–701  
use of low-value clinical services, 29(6):785–792  
use of point-of-care tests, 29(3):371–376  
use of telehealth in primary care, 29(4):432–433  
young, impact of debt on, 29(2):177–179
- Family planning**  
fertility apps marketed to avoid pregnancy, 29(4):508–511  
IUD use based on who initiated discussion of placement, 29(1):24–28
- Fasting**, almond “appetizer” effect on GTT results, 29(6):759–766
- Fecal immunochemical test (FIT)**, for colon cancer screening, 29(6):672–681
- Fertility**, apps marketed to avoid pregnancy, 29(4):508–511
- Fever**  
C-reactive protein level as marker in febrile children, 29(4):460–468  
self-diagnosis of cotton fever by IV drug users, 29(2):276–279
- Focus groups**  
academic detailing intervention for cancer screening, 29(5):533–542  
kiosk technology for blood pressure, 29(5):620–629  
medical home implementation in small practices, 29(6):767–774
- Follow-up studies**  
C-reactive protein level as marker in febrile children, 29(4):460–468  
childhood obesity intervention, 29(4):434–443  
communicating delays in diagnostic evaluation, 29(4):469–473  
patient portals and blood pressure control, 29(4):452–459  
watchful waiting and diagnostic testing, 29(6):710–717
- Food habits**, almond “appetizer” effect on GTT results, 29(6):759–766
- Food supply**, childhood obesity intervention, 29(4):434–443
- Forecasting**, Pisacano scholars’ reflections from the Starfield Summit, 29(6):793–804
- Frail elderly**, reducing functional decline, 29(2):209–217
- Gay persons**, improving patient-centered “LGBTQ” primary care, 29(1):156–160
- Genetics**, vanished twins and misdiagnosed sex, 29(3):411–413
- Geriatrics**, reducing functional decline in frail elderly, 29(2):209–217
- Glucose tolerance test**, results, almond “appetizer” effect on, 29(6):759–766
- Glycated hemoglobins**, primary care continuity improves diabetic health, 29(3):318–324
- Grants**  
federal research funding for family medicine, 29(5):531–532  
limited NIH funding for family medicine research, 29(5):528–530
- Grip strength**, low, prediabetes and, 29(2):280–282
- Guideline adherence**, physician use of low-value clinical services, 29(6):785–792
- Hand strength**, low grip strength and prediabetes, 29(2):280–282
- Health care delivery**  
complexity of “place” in primary care, 29(4):S24–S27  
countercultural heritage of family medicine, 29(4):S45–S48  
family medicine as a disruptive innovation, 29(4):S60–S63  
Generation III family physicians, 29(4):S49–S53  
holding on and letting go, 29(4):S32–S39  
impact of time and timing, 29(4):S28–S31  
people are primary, 29(4):S40–S44  
personal doctoring manifesto, 29(4):S64–S68  
recharging family medicine, 29(4):S15–S18  
reimagining relationships with patients, 29(4):S1–S11  
relationships in tomorrow’s health system, 29(4):S54–S59  
seeking relationships in primary care, 29(4):S19–S23  
social justice and family medicine, 29(4):S69–S71  
tribute to G. Gayle Stephens, 29(4):S12–S14
- Health care disparities**  
community engagement in impoverished communities, 29(3):325–338  
doula support during pregnancy, 29(3):308–317  
reducing health inequities using sociodemographic data, 29(3):348–355  
school-based health centers, 29(3):339–347  
social determinants of health and primary care, 29(3):297–300
- Health care economics and organizations**, effects of PCMH transformation on child patient experience, 29(1):60–68
- Health care reform**, underuse of primary care in China, 29(2):240–247

- Health care systems**, choosing words wisely in the PCMH, 29(2):248–253
- Health care team**, teamlets in primary care, 29(1):135–138
- Health literacy**, toolkit, to improve medication review, 29(1):18–23
- Health maintenance organizations** multiple myeloma and diagnostic delay, 29(6):702–709  
telehealth as useful tool, 29(4):430–431  
use of telehealth in primary care, 29(4):432–433
- Health personnel**  
electronic messaging and number of incoming telephone calls, 29(5):613–619  
self-report of alcohol treatment referrals, 29(6):682–687
- Health planning guidelines**, improving patient-centered “LGBTQ” primary care, 29(1):156–160
- Health policy**  
bridging communities, funders, and policymakers for PBRNs, 29(5):630–635  
family medicine residents and educational debt, 29(2):180–181  
impact of debt on young family physicians, 29(2):177–179  
institutions with fewer residencies produce more physicians, 29(3):301–302  
provider diversity on the family medicine team, 29(1):8–9
- Health resources**, providers’ perspectives on palliative care, 29(6):748–758
- Health services**  
access to care associated with lower obesity rates, 29(2):182–190  
care coordinator prevalence in practices and PCMHs, 29(6):652–653
- Health services research**, technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Hearing loss**  
hearing loss health care for older adults, 29(3):394–403  
pediatric auditory brainstem implant for habilitation, 29(2):286–288
- Hemoglobins**  
clinical decision support for chronic kidney disease, 29(5):604–612  
FIT for colon cancer screening, 29(6):672–681  
primary care continuity improves diabetic health, 29(3):318–324
- Hispanic Americans**, electronic portal use among vulnerable patients, 29(5):592–603
- HIV infections**, same- vs. opposite-sex-partnered patients’ medical diagnoses, 29(6):688–693
- Homelessness**, community engagement in impoverished communities, 29(3):325–338
- Homosexuality**  
improving patient-centered “LGBTQ” primary care, 29(1):156–160  
same- vs. opposite-sex-partnered patients’ medical diagnoses, 29(6):688–693
- Hospital readmission**, risk, prediction by admission data, 29(1):50–59
- Hospital records**, C-reactive protein level as marker in febrile children, 29(4):460–468
- Human immunodeficiency virus (HIV)**, prevention, preexposure prophylaxis for, 29(1):143–151
- Humans**  
C-reactive protein level as marker in febrile children, 29(4):460–468  
communicating delays in diagnostic evaluation, 29(4):469–473  
PCMH recognition and chronic pain management, 29(4):474–481
- Hydroxymethylglutaryl-CoA reductase inhibitors**, avoiding myalgia with statins in exercising adults, 29(6):727–740
- Hyperglycemia**, almond “appetizer” effect on GTT results, 29(6):759–766
- Hyperlactation**, diagnosis and management, 29(1):139–142
- Hypertension**  
clinical decision support for chronic kidney disease, 29(5):604–612  
kiosk technology for blood pressure, 29(5):620–629  
patient portals and blood pressure control, 29(4):452–459  
stakeholder engagement in patient-reported outcomes measure implementation, 29(1):102–115
- Hysterectomy**, physician use of low-value clinical services, 29(6):785–792
- Iatrogenic disease**, nonbacterial causes of lymphangitis, 29(6):808–812
- Industry**, researchers’ experience with data sharing, 29(6):805–807
- Infectious arthritis**, antibiotic prophylaxis and total joint replacement, 29(4):500–507
- Infectious diseases**, preexposure prophylaxis for HIV prevention, 29(1):143–151
- Information services**, electronic portal use among vulnerable patients, 29(5):592–603
- Information systems**  
electronic messaging and number of incoming telephone calls, 29(5):613–619  
strategies for stakeholders to improve EHRs, 29(1):126–134
- Inpatient care**, PCMH impact on chronically ill patients’ utilization, 29(4):482–495
- Insects**, nonbacterial causes of lymphangitis, 29(6):808–812
- Insurance**, patients with diabetes served by safety-net centers, 29(3):356–370
- Insurance coverage**  
participation in weight loss programs, 29(5):572–580  
PCMH impact on chronically ill patients’ utilization, 29(4):482–495
- Interdisciplinary health team**, reducing functional decline in frail elderly, 29(2):209–217
- Internet**  
care coordination in PCMHs, 29(1):90–101  
electronic messaging and number of incoming telephone calls, 29(5):613–619  
patient portals and blood pressure control, 29(4):452–459
- Intrauterine devices**, use based on who initiated discussion of placement, 29(1):24–28
- Intravenous drug abuse**, cotton fever self-diagnosis, 29(2):276–279
- Journal of the American Board of Family Medicine (JABFM)*, peer reviewers for, 29(2):173–176
- Kidney disease**, chronic  
clinical decision support, 29(5):604–612  
local learning collaboratives, 29(5):543–552
- Lactation disorders**, breast milk oversupply, 29(1):139–142
- Leadership**, Pisacano scholars’ reflections from the Starfield Summit, 29(6):793–804
- Learning**, collaboratives, local, for chronic kidney disease, 29(5):543–552
- Lesbians**, improving patient-centered “LGBTQ” primary care, 29(1):156–160
- Levofloxacin**, in males with a UTI, 29(6):654–662
- Life style**  
childhood obesity intervention, 29(4):434–443  
physician attitudes toward diabetes prevention, 29(6):663–671
- Loneliness**, same- vs. opposite-sex-partnered patients’ medical diagnoses, 29(6):688–693
- Long-term care**, continuity of care and reducing emergency department visits, 29(2):201–208
- Lung cancer**, screening guidelines, 29(1):152–155
- Lymphangitis**, nonbacterial causes, 29(6):808–812
- Mass screening**, FIT for colon cancer screening, 29(6):672–681
- Maternal health**, treatment of acne in pregnancy, 29(2):254–262
- Medical assistance**, electronic portal use among vulnerable patients, 29(5):592–603

- Medical errors**, using a health literacy toolkit to improve medication review, 29(1):18–23
- Medical home**  
 case mix severity impact on quality improvement in a PCMH, 29(1):116–125  
 cost for level 3 medical home recognition, 29(1):69–77  
 PCMH increased colorectal cancer screening, 29(2):191–200  
 referrals for diabetes education in a medical home, 29(3):377–384  
 school-based health centers, 29(3):339–347  
 time to remission for depression with CCM, 29(1):10–17
- Medical informatics**, simulation of ICD-9 to ICD-10-CM transition, 29(1):29–36
- Medical records**, same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Medical societies**, evaluating Choosingly Wisely™ using SORT, 29(4):512–515
- Medically uninsured**, electronic portal use among vulnerable patients, 29(5):592–603
- Medication**  
 pretreatment use and drinking goal in alcohol treatment, 29(1):37–49  
 review, improving, using a health literacy toolkit, 29(1):18–23
- Medication adherence**, stakeholder engagement in patient-reported outcomes measure implementation, 29(1):102–115
- Medicine**, researchers' experience with data sharing, 29(6):805–807
- MEDLINE**, nonbacterial causes of lymphangitis, 29(6):808–812
- Menstrual cycle**, fertility apps marketed to avoid pregnancy, 29(4):508–511
- Mental health**  
 family medicine: bridge to life, 29(1):161–164  
 patient portals and blood pressure control, 29(4):452–459  
 self-report of alcohol treatment referrals, 29(6):682–687
- Metformin**, physician attitudes toward diabetes prevention, 29(6):663–671
- Minnesota**, PCMH impact on chronically ill patients' utilization, 29(4):482–495
- Minority health**  
 improving patient-centered “LG-BTQ” primary care, 29(1):156–160  
 vitamin D deficiency by body weight and race/ethnicity, 29(2):226–232
- Motivation**  
 electronic messaging and number of incoming telephone calls, 29(5):613–619  
 physician attitudes toward diabetes prevention, 29(6):663–671
- Multicultural practices**, improving patient-centered “LGBTQ” primary care, 29(1):156–160
- Multiple myeloma**, diagnostic delay and, 29(6):702–709
- Muscles**, avoiding myalgia with statins in exercising adults, 29(6):727–740
- Muscular diseases**, avoiding myalgia with statins in exercising adults, 29(6):727–740
- Musculoskeletal pain**, PCMH recognition and chronic pain management, 29(4):474–481
- Myalgia**, avoiding, with statins, in exercising adults, 29(6):727–740
- Mycoses**, nonbacterial causes of lymphangitis, 29(6):808–812
- Natural family planning**, fertility apps marketed to avoid pregnancy, 29(4):508–511
- Neoplasms**, communicating delays in diagnostic evaluation, 29(4):469–473
- Nurse practitioners**, medical home implementation in small practices, 29(6):767–774
- Nutritional sciences**, vitamin D deficiency by body weight and race/ethnicity, 29(2):226–232
- Obesity**  
 childhood obesity intervention, 29(4):434–443  
 participation in weight loss programs, 29(5):572–580  
 patient portals and blood pressure control, 29(4):452–459  
 rates, lower, association with access to care, 29(2):182–190  
 same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Obstetrics**  
 doula support during pregnancy, 29(3):308–317  
 vanished twins and misdiagnosed sex, 29(3):411–413
- Obstructive sleep apnea**, perioperative care, 29(2):263–275
- Office visits**, participation in weight loss programs, 29(5):572–580
- Opioid analgesics**  
 PCMH recognition and chronic pain management, 29(4):474–481  
 problem drug-related behavior, 29(6):718–726
- Opioid-related disorders**, problem drug-related behavior, 29(6):718–726
- Opioids**, perioperative care of obstructive sleep apnea, 29(2):263–275
- Oregon**, electronic portal use among vulnerable patients, 29(5):592–603
- Orthopedics**, insufficiency fractures and bisphosphonate therapy, 29(3):404–407
- Osteoporosis**  
 clinical risk tools for predicting, 29(2):233–239  
 physician use of low-value clinical services, 29(6):785–792
- Otolaryngology**, hearing loss health care for older adults, 29(3):394–403
- Outcome assessment**  
 C-reactive protein level as marker in febrile children, 29(4):460–468  
 childhood obesity intervention, 29(4):434–443
- Outpatients**  
 levofloxacin in males with a UTI, 29(6):654–662  
 PCMH impact on chronically ill patients' utilization, 29(4):482–495
- Palliative care**, providers' perspectives on, 29(6):748–758
- Palpation**, triage amalgamated dermoscopic algorithm, 29(6):694–701
- Papanicolaou test**, physician use of low-value clinical services, 29(6):785–792
- Parents**, opinions about concepts of partnership, 29(5):563–571
- Patient advisory councils**, patient-centered care and, 29(6):775–784
- Patient care**  
 patient portals and blood pressure control, 29(4):452–459  
 primary care panel size, 29(4):444–451
- Patient-centered care**  
 advisory councils and, 29(6):775–784  
 cost for level 3 medical home recognition, 29(1):69–77  
 improving patient-centered “LG-BTQ” primary care, 29(1):156–160  
 lung cancer screening guidelines, 29(1):152–155  
 medical home implementation in small practices, 29(6):767–774  
 parents' and pediatricians' views of partnership, 29(5):563–571  
 provider diversity on the family medicine team, 29(1):8–9  
 school-based health centers, 29(3):339–347  
 telehealth as useful tool, 29(4):430–431  
 use of telehealth in primary care, 29(4):432–433
- Patient-centered medical homes (PCMHs)**  
 care coordination in, 29(1):90–101  
 care coordinator prevalence, 29(6):652–653  
 case mix severity impact on quality improvement, 29(1):116–125  
 choosing words wisely in, 29(2):248–253  
 impact on chronically ill patients' utilization, 29(4):482–495  
 implementation in small practices, 29(6):767–774  
 increased colorectal cancer screening, 29(2):191–200  
 recognition and chronic pain management, 29(4):474–481  
 transformation, effects on child patient experience, 29(1):60–68



- vs primary care, learning more about, 29(1):4–7
- Patient-centered outcomes research**  
 stakeholder engagement in a patient-reported outcomes measure implementation, 29(1):102–115  
 stakeholder engagement in patient-reported outcomes measure implementation, 29(1):102–115  
 time to remission for depression with CCM, 29(1):10–17
- Patient compliance**, FIT for colon cancer screening, 29(6):672–681
- Patient education**  
 referrals for diabetes education in a medical home, 29(3):377–384  
 using a health literacy toolkit to improve medication review, 29(1):18–23
- Patient engagement**, advisory councils and patient-centered care, 29(6):775–784
- Patient navigators**, to improve diabetes outcome, 29(1):78–89
- Patient outcome assessment**, stakeholder engagement in patient-reported outcomes measure implementation, 29(1):102–115
- Patient participation**  
 kiosk technology for blood pressure, 29(5):620–629  
 technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Patient satisfaction**, primary care panel size, 29(4):444–451
- Pediatrics**  
 C-reactive protein level as marker in febrile children, 29(4):460–468  
 childhood obesity intervention, 29(4):434–443  
 effects of PCMH transformation on child patient experience, 29(1):60–68  
 pediatric auditory brainstem implant for habilitation, 29(2):286–288  
 pediatricians' opinions about concepts of partnership, 29(5):563–571  
 perioperative care of obstructive sleep apnea, 29(2):263–275
- Perception**, providers' perspectives on palliative care, 29(6):748–758
- Personal health records**, patient portals and blood pressure control, 29(4):452–459
- Personal physicians**  
 complexity of "place" in primary care, 29(4):S24–S27  
 countercultural heritage of family medicine, 29(4):S45–S48  
 family medicine as a disruptive innovation, 29(4):S60–S63  
 Generation III family physicians, 29(4):S49–S53  
 holding on and letting go, 29(4):S32–S39  
 impact of time and timing, 29(4):S28–S31  
 people are primary, 29(4):S40–S44  
 personal doctoring manifesto, 29(4):S64–S68
- recharging family medicine, 29(4):S15–S18  
 reimagining relationships with patients, 29(4):S1–S11  
 relationships in tomorrow's health system, 29(4):S54–S59  
 seeking relationships in primary care, 29(4):S19–S23  
 social justice and family medicine, 29(4):S69–S71  
 tribute to G. Gayle Stephens, 29(4):S12–S14
- Personal satisfaction**, Pisacano scholars' reflections from the Starfield Summit, 29(6):793–804
- Pharmacotherapy**, preexposure prophylaxis for HIV prevention, 29(1):143–151
- Physical examination**, watchful waiting and diagnostic testing, 29(6):710–717
- Physician-patient relations**  
 family medicine: bridge to life, 29(1):161–164  
 IUD use based on who initiated discussion of placement, 29(1):24–28
- Physicians**  
 advice for e-cigarette use, 29(6):741–747  
 institutions with fewer residencies produce more physicians, 29(3):301–302  
 pediatricians' views of partnership, 29(5):563–571  
 physician attitudes toward diabetes prevention, 29(6):663–671  
 triage amalgamated dermoscopic algorithm, 29(6):694–701  
 use of point-of-care tests, 29(3):371–376
- Pilot projects**, childhood obesity intervention, 29(4):434–443
- Point-of-care systems**  
 C-reactive protein level as marker in febrile children, 29(4):460–468  
 use of point-of-care tests by primary care physicians, 29(3):371–376
- Polypharmacy**  
 admission data predict hospital readmission risk, 29(1):50–59  
 using a health literacy toolkit to improve medication review, 29(1):18–23
- Population characteristics**, taking the community pulse while caring for patients, 29(3):419–422
- Population health**  
 complexity of "place" in primary care, 29(4):S24–S27  
 countercultural heritage of family medicine, 29(4):S45–S48  
 family medicine as a disruptive innovation, 29(4):S60–S63  
 Generation III family physicians, 29(4):S49–S53  
 holding on and letting go, 29(4):S32–S39  
 impact of time and timing, 29(4):S28–S31  
 people are primary, 29(4):S40–S44  
 personal doctoring manifesto, 29(4):S64–S68
- recharging family medicine, 29(4):S15–S18  
 reimagining relationships with patients, 29(4):S1–S11  
 relationships in tomorrow's health system, 29(4):S54–S59  
 seeking relationships in primary care, 29(4):S19–S23  
 social justice and family medicine, 29(4):S69–S71  
 tribute to G. Gayle Stephens, 29(4):S12–S14
- Postal service**, FIT for colon cancer screening, 29(6):672–681
- Poverty**, electronic portal use among vulnerable patients, 29(5):592–603
- Practice-based research**  
 bridging communities, funders, and policymakers for PBRNs, 29(5):630–635  
 case mix severity impact on quality improvement in a PCMH, 29(1):116–125  
 clinical decision support for chronic kidney disease, 29(5):604–612  
 decision-to-implement worksheet, 29(5):553–562  
 electronic portal use among vulnerable patients, 29(5):592–603  
 kiosk technology for blood pressure, 29(5):620–629  
 local learning collaboratives for chronic kidney disease, 29(5):543–552  
 participation in weight loss programs, 29(5):572–580  
 stakeholder engagement in patient-reported outcomes measure implementation, 29(1):102–115  
 technology-based patient engagement strategies within PBRNs, 29(5):581–591  
 using patient navigators to improve diabetes outcome, 29(1):78–89
- Practice improvement**, advisory councils and patient-centered care, 29(6):775–784
- Practice management**, cost for level 3 medical home recognition, 29(1):69–77
- Prediabetes**  
 almond "appetizer" effect on GTT results, 29(6):759–766  
 low grip strength and prediabetes, 29(2):280–282  
 physician attitudes toward diabetes prevention, 29(6):663–671  
 prediabetes diagnosis and treatment, 29(2):283–285
- Pregnancy**  
 doula support during, 29(3):308–317  
 fertility apps marketed to avoid, 29(4):508–511  
 treatment of acne during, 29(2):254–262
- Prenatal care**, vanished twins and misdiagnosed sex, 29(3):411–413
- Prevalence**  
 care coordinator prevalence in practices and PCMHs, 29(6):652–653

- problem drug-related behavior, 29(6):718–726
- same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Prevention and control**, preexposure prophylaxis for HIV prevention, 29(1):143–151
- Preventive medicine**, vitamin D deficiency by body weight and race/ethnicity, 29(2):226–232
- Primary care physicians**  
 communicating delays in diagnostic evaluation, 29(4):469–473  
 medical home implementation in small practices, 29(6):767–774  
 panel size, 29(4):496–499  
 use of low-value clinical services, 29(6):785–792  
 watchful waiting and diagnostic testing, 29(6):710–717
- Primary health care**  
 academic detailing intervention for cancer screening, 29(5):533–542  
 access to care associated with lower obesity rates, 29(2):182–190  
 bridging communities, funders, and policymakers for PBRNs, 29(5):630–635  
 care coordination in PCMHs, 29(1):90–101  
 childhood obesity intervention, 29(4):434–443  
 choosing words wisely in the PCMH, 29(2):248–253  
 clinical decision support for chronic kidney disease, 29(5):604–612  
 complexity of “place” in, 29(4):S24–S27  
 continuity of care and reducing emergency department visits, 29(2):201–208  
 countercultural heritage of family medicine, 29(4):S45–S48  
 decision-to-implement worksheet, 29(5):553–562  
 evaluating Choosing Wisely™ using SORT, 29(4):512–515  
 family medicine as a disruptive innovation, 29(4):S60–S63  
 federal research funding for family medicine, 29(5):531–532  
 Generation III family physicians, 29(4):S49–S53  
 holding on and letting go, 29(4):S32–S39  
 impact of time and timing, 29(4):S28–S31  
 improving patient-centered “LGBTQ” primary care, 29(1):156–160  
 institutions with fewer residencies produce more physicians, 29(3):301–302  
 integrating behavioral health using lean workflow analysis, 29(3):385–393  
 kiosk technology for blood pressure, 29(5):620–629  
 limited NIH funding for family medicine research, 29(5):528–530
- local learning collaboratives for chronic kidney disease, 29(5):543–552
- multiple myeloma and diagnostic delay, 29(6):702–709
- participation in weight loss programs, 29(5):572–580
- patient-centered “LGBTQ” care, 29(1):156–160
- patient portals and blood pressure control, 29(4):452–459
- PCMH recognition and chronic pain management, 29(4):474–481
- people are primary, 29(4):S40–S44
- personal doctoring manifesto, 29(4):S64–S68
- Pisacano scholars' reflections from the Starfield Summit, 29(6):793–804
- prediabetes diagnosis and treatment, 29(2):283–285
- preexposure prophylaxis for HIV prevention, 29(1):143–151
- primary care panel size, 29(4):444–451, 29(4):496–499
- problem drug-related behavior, 29(6):718–726
- providers' perspectives on palliative care, 29(6):748–758
- recharging family medicine, 29(4):S15–S18
- reducing functional decline in frail elderly, 29(2):209–217
- reimagining relationships with patients, 29(4):S1–S11
- relationships in tomorrow's health system, 29(4):S54–S59
- same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- screening for adverse childhood experiences, 29(3):303–307
- seeking relationships in, 29(4):S19–S23
- self-report of alcohol treatment referrals, 29(6):682–687
- social determinants of health and, 29(3):297–300
- social determinants of health in a clinic setting, 29(3):414–418
- social justice and family medicine, 29(4):S69–S71
- strategies for stakeholders to improve EHRs, 29(1):126–134
- teamlets in primary care, 29(1):135–138
- time to remission for depression with CCM, 29(1):10–17
- tribute to G. Gayle Stephens, 29(4):S12–S14
- underuse of primary care in China, 29(2):240–247
- vs PCMHs, learning more about, 29(1):4–7
- Privacy**, technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Private practice**, parents' and pediatricians' views of partnership, 29(5):563–571
- Probability**  
 C-reactive protein level as marker in febrile children, 29(4):460–468
- watchful waiting and diagnostic testing, 29(6):710–717
- Professionalism**  
 complexity of “place” in primary care, 29(4):S24–S27  
 countercultural heritage of family medicine, 29(4):S45–S48  
 family medicine as a disruptive innovation, 29(4):S60–S63  
 Generation III family physicians, 29(4):S49–S53  
 holding on and letting go, 29(4):S32–S39  
 impact of time and timing, 29(4):S28–S31  
 people are primary, 29(4):S40–S44  
 personal doctoring manifesto, 29(4):S64–S68  
 recharging family medicine, 29(4):S15–S18  
 reimagining relationships with patients, 29(4):S1–S11  
 relationships in tomorrow's health system, 29(4):S54–S59  
 seeking relationships in primary care, 29(4):S19–S23  
 social justice and family medicine, 29(4):S69–S71  
 tribute to G. Gayle Stephens, 29(4):S12–S14
- Prognosis**, multiple myeloma and diagnostic delay, 29(6):702–709
- Proportional hazards model**, patient portals and blood pressure control, 29(4):452–459
- Prospective studies**, C-reactive protein level as marker in febrile children, 29(4):460–468
- Prunus dulcis**, almond “appetizer” effect on GTT results, 29(6):759–766
- Public health**  
 access to care associated with lower obesity rates, 29(2):182–190  
 taking the community pulse while caring for patients, 29(3):419–422
- PubMed**  
 avoiding myalgia with statins in exercising adults, 29(6):727–740  
 nonbacterial causes of lymphangitis, 29(6):808–812
- Qualitative research**  
 academic detailing intervention for cancer screening, 29(5):533–542  
 care coordination in PCMHs, 29(1):90–101  
 choosing words wisely in the PCMH, 29(2):248–253  
 medical home implementation in small practices, 29(6):767–774  
 providers' perspectives on palliative care, 29(6):748–758  
 strategies for stakeholders to improve EHRs, 29(1):126–134
- Quality improvement**  
 academic detailing intervention for cancer screening, 29(5):533–542  
 advisory councils and patient-centered care, 29(6):775–784



- integrating behavioral health using lean workflow analysis, 29(3):385–393
- medical home implementation in small practices, 29(6):767–774
- Quality of health care**
- case mix severity impact on quality improvement in a PCMH, 29(1):116–125
- family medicine: bridge to life, 29(1):161–164
- Quality of life**, avoiding myalgia with statins in exercising adults, 29(6):727–740
- Radiology**, insufficiency fractures and bisphosphonate therapy, 29(3):404–407
- Randomized controlled trials as topic**, patient portals and blood pressure control, 29(4):452–459
- Referral and consultation**
- C-reactive protein level as marker in febrile children, 29(4):460–468
- triage amalgamated dermoscopic algorithm, 29(6):694–701
- Registries**
- patient portals and blood pressure control, 29(4):452–459
- problem drug-related behavior, 29(6):718–726
- technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Regression analysis**, primary care panel size, 29(4):444–451
- Reminder systems**, academic detailing intervention for cancer screening, 29(5):533–542
- Renal insufficiency**, local learning collaboratives for chronic kidney disease, 29(5):543–552
- Replacement arthroplasty**, antibiotic prophylaxis and total joint replacement, 29(4):500–507
- Research**
- evaluating Choosing Wisely™ using SORT, 29(4):512–515
- federal research funding for family medicine, 29(5):531–532
- limited NIH funding for family medicine research, 29(5):528–530
- primary care panel size, 29(4):496–499
- researchers' experience with data sharing, 29(6):805–807
- Research design**, PCMH impact on chronically ill patients' utilization, 29(4):482–495
- Research personnel**
- experience with data sharing, 29(6):805–807
- Pisacano scholars' reflections from the Starfield Summit, 29(6):793–804
- Research support as topic**, bridging communities, funders, and policymakers for PBRNs, 29(5):630–635
- Residence characteristics**
- providers' perspectives on palliative care, 29(6):748–758
- taking the community pulse while caring for patients, 29(3):419–422
- Residencies**
- institutions with fewer residencies produce more physicians, 29(3):301–302
- PCMH recognition and chronic pain management, 29(4):474–481
- Respiratory failure**, perioperative care of obstructive sleep apnea, 29(2):263–275
- Respiratory tract diseases**, perioperative care of obstructive sleep apnea, 29(2):263–275
- Retrospective studies**
- electronic messaging and number of incoming telephone calls, 29(5):613–619
- electronic portal use among vulnerable patients, 29(5):592–603
- FIT for colon cancer screening, 29(6):672–681
- multiple myeloma and diagnostic delay, 29(6):702–709
- primary care panel size, 29(4):444–451
- Review, systematic**, diagnosis and management of breast milk oversupply, 29(1):139–142
- Risk**
- hospital readmission, admission data predict, 29(1):50–59
- watchful waiting and diagnostic testing, 29(6):710–717
- Risk assessment**
- clinical decision support for chronic kidney disease, 29(5):604–612
- technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Routine care**, watchful waiting and diagnostic testing, 29(6):710–717
- Rural communities**, improving patient-centered “LGBTQ” primary care, 29(1):156–160
- Safety-net providers**
- patients with diabetes served by safety-net centers, 29(3):356–370
- problem drug-related behavior, 29(6):718–726
- telehealth as useful tool, 29(4):430–431
- use of telehealth in primary care, 29(4):432–433
- School health services**, “expanded medical home,” 29(3):339–347
- Screening**
- academic detailing intervention for cancer screening, 29(5):533–542
- for adverse childhood experiences, 29(3):303–307
- clinical risk tools for predicting osteoporosis, 29(2):233–239
- lung cancer screening guidelines, 29(1):152–155
- perioperative care of obstructive sleep apnea, 29(2):263–275
- Seasons**, FIT for colon cancer screening, 29(6):672–681
- Self-care**
- almond “appetizer” effect on GTT results, 29(6):759–766
- using patient navigators to improve diabetes outcome, 29(1):78–89
- Self-report**
- alcohol treatment referrals, 29(6):682–687
- participation in weight loss programs, 29(5):572–580
- physician use of low-value clinical services, 29(6):785–792
- same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Sensitivity and specificity**, triage amalgamated dermoscopic algorithm, 29(6):694–701
- Sexual partners**, same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Sinusitis**, physician use of low-value clinical services, 29(6):785–792
- Skin**, nonbacterial causes of lymphangitis, 29(6):808–812
- Skin cancer**
- basal cell blancher, 29(3):408–410
- triage amalgamated dermoscopic algorithm, 29(6):694–701
- Sleep disorders**, perioperative care of obstructive sleep apnea, 29(2):263–275
- Smoking**
- patient portals and blood pressure control, 29(4):452–459
- physician advice for e-cigarette use, 29(6):741–747
- same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Smoking cessation**, physician advice for e-cigarette use, 29(6):741–747
- Snoring**, perioperative care of obstructive sleep apnea, 29(2):263–275
- Social class**, patient portals and blood pressure control, 29(4):452–459
- Social determinants of health**
- in a clinic setting, 29(3):414–418
- community engagement in impoverished communities, 29(3):325–338
- doula support during pregnancy, 29(3):308–317
- patients with diabetes served by safety-net centers, 29(3):356–370
- primary care continuity improves diabetic health, 29(3):318–324
- reducing health inequities using sociodemographic data, 29(3):348–355
- school-based health centers, 29(3):339–347
- screening for adverse childhood experiences, 29(3):303–307
- social determinants of health and primary care, 29(3):297–300
- taking the community pulse while caring for patients, 29(3):419–422
- Social justice**
- complexity of “place” in primary care, 29(4):S24–S27

- countercultural heritage of family medicine, 29(4):S45–S48
- family medicine as a disruptive innovation, 29(4):S60–S63
- Generation III family physicians, 29(4):S49–S53
- holding on and letting go, 29(4):S32–S39
- impact of time and timing, 29(4):S28–S31
- people are primary, 29(4):S40–S44
- personal doctoring manifesto, 29(4):S64–S68
- recharging family medicine, 29(4):S15–S18
- reimagining relationships with patients, 29(4):S1–S11
- relationships in tomorrow's health system, 29(4):S54–S59
- seeking relationships in primary care, 29(4):S19–S23
- social justice and family medicine, 29(4):S69–S71
- tribute to G. Gayle Stephens, 29(4):S12–S14
- Social networks**, IUD use based on who initiated discussion of placement, 29(1):24–28
- Social stigma**, same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Special communications**
- “A Paradox Persists When the Paradigm Is Wrong”: Pisacano Scholars' Reflections from the Inaugural Starfield Summit, 29(6):793–804
- A Primary Care Panel Size of 2500 Is neither Accurate nor Reasonable, 29(4):496–499
- Researchers' Experience with Clinical Data Sharing, 29(6):805–807
- Strategies for Primary Care Stakeholders to Improve Electronic Health Records (EHRs), 29(1):126–134
- Teamlets in Primary Care: Enhancing Patient and Clinician Experience, 29(1):135–138
- Spider bites**, nonbacterial causes of lymphangitis, 29(6):808–812
- Statins**, avoiding myalgia with, in exercising adults, 29(6):727–740
- Substance abuse detection**, problem drug-related behavior, 29(6):718–726
- Substance-related disorders**
- PCMH recognition and chronic pain management, 29(4):474–481
- problem drug-related behavior, 29(6):718–726
- same- vs. opposite-sex-partnered patients' medical diagnoses, 29(6):688–693
- Subtrochanteric fractures**, bisphosphonate therapy and, 29(3):404–407
- Suicide**, family medicine: bridge to life, 29(1):161–164
- Surgery**, perioperative care of obstructive sleep apnea, 29(2):263–275
- Surveys and questionnaires**
- academic detailing intervention for cancer screening, 29(5):533–542
- kiosk technology for blood pressure, 29(5):620–629
- parents' and pediatricians' views of partnership, 29(5):563–571
- physician advice for e-cigarette use, 29(6):741–747
- physician attitudes toward diabetes prevention, 29(6):663–671
- physician use of low-value clinical services, 29(6):785–792
- technology-based patient engagement strategies within PBRNs, 29(5):581–591
- use of telehealth in primary care, 29(4):432–433
- Telemedicine**
- use in primary care, 29(4):432–433
- as useful tool, 29(4):430–431
- Telephone**
- communicating delays in diagnostic evaluation, 29(4):469–473
- electronic messaging and number of incoming telephone calls, 29(5):613–619
- providers' perspectives on palliative care, 29(6):748–758
- self-report of alcohol treatment referrals, 29(6):682–687
- Temperature**, FIT for colon cancer screening, 29(6):672–681
- Tobacco products**, physician advice for e-cigarette use, 29(6):741–747
- Tobacco use disorder**, physician advice for e-cigarette use, 29(6):741–747
- Total joint replacement**, patients with, antibiotic prophylaxis for, 29(4):500–507
- Transgender persons**, improving patient-centered “LGBTQ” primary care, 29(1):156–160
- Translational medical research**, decision-to-implement worksheet, 29(5):553–562
- Transportation**, FIT for colon cancer screening, 29(6):672–681
- Triage**, triage amalgamated dermatoscopic algorithm, 29(6):694–701
- Underserved populations**
- community engagement in impoverished communities, 29(3):325–338
- doula support during pregnancy, 29(3):308–317
- primary care continuity improves diabetic health, 29(3):318–324
- United States**
- childhood obesity intervention, 29(4):434–443
- evaluating Choosing Wisely™ using SORT, 29(4):512–515
- Urinary tract infections**, levofloxacin in males with, 29(6):654–662
- U.S. Agency for Healthcare Research and Quality**, technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Vitamin D**, deficiency by body weight and race/ethnicity, 29(2):226–232
- Vulnerable populations**, electronic portal use among, 29(5):592–603
- Watchful waiting**, diagnostic testing and, 29(6):710–717
- Weight loss**
- multiple myeloma and diagnostic delay, 29(6):702–709
- programs, participation in, 29(5):572–580
- Women's health**
- clinical risk tools for predicting osteoporosis, 29(2):233–239
- diagnosis and management of breast milk oversupply, 29(1):139–142
- doula support during pregnancy, 29(3):308–317
- fertility apps marketed to avoid pregnancy, 29(4):508–511
- IUD use based on who initiated discussion of placement, 29(1):24–28
- treatment of acne in pregnancy, 29(2):254–262
- Work**, use of telehealth in primary care, 29(4):432–433
- Workflow**
- academic detailing intervention for cancer screening, 29(5):533–542
- clinical decision support for chronic kidney disease, 29(5):604–612
- kiosk technology for blood pressure, 29(5):620–629
- Workforce**
- family medicine residents and educational debt, 29(2):180–181
- impact of debt on young family physicians, 29(2):177–179
- Workload**
- electronic messaging and number of incoming telephone calls, 29(5):613–619
- primary care panel size, 29(4):496–499
- technology-based patient engagement strategies within PBRNs, 29(5):581–591
- Writing**, Pisacano scholars' reflections from the Starfield Summit, 29(6):793–804