

We try to publish authors' responses in the same edition with readers' comments. Time constraints might prevent this in some cases. The problem is compounded in a bimonthly journal where continuity of comment and redress are difficult to achieve. When the redress appears 2 months after the comment, 4 months will have passed since the article was published. Therefore, we would suggest to our readers that their correspondence about published papers be submitted as soon as possible after the article appears.

Narcotic Theft

To the Editor: We report a case of narcotic theft by a man who disguised himself as a visiting home nurse at a patient's home.

Our patient was an 89-year-old woman who had sustained a right femoral neck fracture 20 months previously. Severe coronary artery disease found by angiography precluded surgery. The patient was cared for at home and was receiving one tablet of hydrocodone bitartrate 5 mg with acetaminophen 500 mg (Vicodin) each night for the last 17 months. She was living with her daughter, who was in regular communication with our clinic. The patient also was receiving cardiac medications for heart failure, warfarin for life-long anticoagulation, and thyroxine for hypothyroidism. Patient had regular follow-up with home nurse visits, home physician visits, and periodic visits at our clinic and an orthopedist's office. The patient had an excellent compliance record for all her medications, and she never requested tablets of hydrocodone in addition to those regularly prescribed to her.

One day a man appeared at the patient's home identifying himself as a visiting home nurse. The patient wondered about the visiting nurse coming to her home on that particular day because no such visit was previously scheduled. She nonetheless admitted the impostor into her home. The intruder told the patient that he had come to check on her medications. He then examined all the medication bottles and told her that because she was on so many costly medications, he could get the hydrocodone for her free. The intruder also assured the patient that although he would take the current amount of hydrocodone in the bottle with him, he would refill the prescription and return with additional tablets of hydrocodone at no charge to the patient. The intruder did not take any other medication bottles with him. Obviously the intruder never returned. When the patient's daughter arrived home after work and learned about the incident, she immediately surmised that the man was a thief and called our clinic and the police to report the theft.

The reason the thief targeted our patient's home is unknown. He might have become aware that our patient was taking narcotic pain medications, or he might just have been targeting homes with a wheelchair access

to find patients who might be taking narcotic medications. Sadly the patient died from a sudden cardiac arrest at home 2 weeks after the incident.

Our case illustrates yet another shrewd method of stealing narcotics. To our knowledge, this method has not previously been reported. Our case should serve as a reminder that patients should be advised not to let unscheduled visiting nurses into their home. When a visiting nurse does arrive at the house, the patient should ask for the name to establish that the visiting nurse has been previously scheduled before letting that person into the house. This recommendation should prevent or minimize narcotic theft and possibility of personal danger to the fragile and ill patients living at home.

Ram Kakaiya, MD

Jennifer Wenzel-Wamhoff, PharmD
University of Illinois Primary Care Clinic
Rockton, Ill

Hormonal Therapy and Management of Premenstrual Syndrome

To the Editor: I was pleased to see an article by a family physician on this very important topic.¹ Premenstrual syndrome (PMS) is more common than is thought, and there are no good studies that have tried to elucidate its prevalence, age at onset, and natural history. Very little is known about the neurohormonal mediators as well. Even so, it remains a common condition that affects young reproductive-age women.

This article discusses the use of hormonal therapy in the management of PMS. Schmidt et al² suggest that PMS might represent an abnormal response to normal hormonal changes. FitzGerald et al³ conclude that premenstrual dysphoric disorder appears to be associated with serotonergic deficiency. High concentrations of luteal phase estradiol and luteinizing hormone are related to the severity of premenstrual symptoms.

Halbreich⁴ discusses the possibility of internationally recognized interdisciplinary diagnostic criteria for PMS. PMS can be preceded by a prodromal aura. The fourth edition of the *Diagnostic and Statistical Manual of Mental Disorders* defines PMS as different from a premenstrual dysphoria disorder or luteal phase dysphoric disorder and late luteal phase dysphoria disorder. It might be possible to categorize the variety of symptoms as (1) PMS with primary mood disorder, (2) PMS with fluid retention, (3) PMS with musculoskeletal symptoms, eg, arthralgia, myalgia, and (4) PMS with generalized symptoms eg, fatigue, headaches, and breast tenderness.

Therapy for this disease could be targeted to the actual symptoms and started at the first indication of aura. General measures are to encourage a healthy lifestyle, exercise, and a balanced diet. The commonest medications used are pain relievers.⁵ One way to prevent PMS might be to use oral contraceptive pills to eliminate for-