

We try to publish authors' responses in the same edition with readers' comments. Time constraints might prevent this in some cases. The problem is compounded in a bimonthly journal where continuity of comment and redress are difficult to achieve. When the redress appears 2 months after the comment, 4 months will have passed since the original article was published. Therefore, we would suggest to our readers that their correspondence about published papers be submitted as soon as possible after the article appears.

Treatment of Otitis Externa

To the Editor: I would like to expand on two points made in Halpern et al's recent article about otitis externa treatment patterns.¹ Physician behaviors seem to vary in actual practice from what is accepted practice.² I am certain that many readers reviewed these results with a thoughtful reevaluation of their own practice habits.

In describing a high rate of systemic antibiotic usage in treating otitis externa, the author did mention a comorbid condition that would justify aggressive use of antipseudomonal antimicrobials. Malignant external otitis is an important concern in elderly diabetic patients with external otitis.³ This uncommon but serious condition can be life-threatening, causing erosion and osteomyelitis of the skull base, which can lead to meningitis, brain abscess, and even death. Progressive cranial nerve palsies have also been reported. Family physicians must be aware of this association and begin treatment urgently with appropriate consultation for careful surgical debridement and management.

The second point of interest relates to the lack of "meticulous cleaning of the external auditory meatus" performed by the study group of physicians; of physicians studied, only 7 percent of adults and 2 percent of children were treated with this procedure. The authors mention the lack of documented efficacy of external auditory canal irrigation in the literature as a theoretical justification for not performing removal of discharge and debris. Additional morbidity has been associated with external auditory canal irrigation, however, resulting in reported malignant external otitis in an elderly diabetic patient.⁴

Further investigation of the efficacy of external auditory canal irrigation, comparing outcomes with those of a control group, appears to be indicated if this procedure is to be recommended in future clinical guidelines.

Paul Evans, DO
College of Osteopathic Medicine
Oklahoma State University, Tulsa

References

1. Halpern MT, Palmer CS, Seidlin M. Treatment patterns for otitis externa. *J Am Board Fam Pract* 1999;12:17.

2. Evans P. Otitis externa. In: Taylor RB editor. *Manual of family practice*. Boston: Lippincott-Raven, 1996.
3. Evans P, Hofmann LA. Malignant external otitis - a case report and review. *Am Fam Physician* 1994;49:427-31.
4. Ford GR, Courteney-Harris RG. Another hazard of ear syringing: malignant external otitis. *J Laryngol Otol* 1990;104:70910.

The above letter was referred to the authors of the article question, who offer the following reply.

To the Editor: We would like to thank Dr. Evans for his thoughtful comments on our recent manuscript. As Dr. Evans indicates, malignant otitis externa is a potentially life-threatening condition. As the main risk group for this condition is elderly patients with diabetes, we agree that aggressive treatment of otitis externa in this population with oral or intravenous antibiotics is very appropriate. As Dr. Evans further points out, external auditory canal irrigation should also be used cautiously in this population.

Malignant external otitis media, however, is a very rare condition. The overwhelming majority of patients who have with external otitis media do not fall into the specified high-risk group for the malignant condition. Thus, the high level of treatment of otitis externa with oral as well as topical medications observed in our study (at more than 40 percent of the physician visits) is not only unnecessary but also increases costs and the likelihood of side effects.

Michael T. Halpern, MD, PhD
Cynthia S. Palmer, MS
MEDTAP International
Mindell Seidlin, MD
Daiichi Pharmaceutical Corporation

Reimbursement for Flexible Sigmoidoscopy

To the Editor: Thanks to Dr. Mulvey¹ for the opportunity to provide additional information and comments regarding the health care system, which discriminates against family physicians who wish to provide diagnostic and therapeutic services such as flexible sigmoidoscopy, endoscopic biopsy, and colonoscopy. His letter accurately reflects yet another dimension of this problem. I have further referenced the consequences of regulatory unfairness in the area of reimbursement for flexible sigmoidoscopy.²

The question remains regarding the political will of physicians within the medical specialty of family practice. Certainly the American Academy of Family Physicians has been a source of support for family physicians who wish to maintain the right to incorporate emerging diagnostic and therapeutic skills into their practice. The Residency Review Committee through their accreditation process attempts to maintain the breadth of the specialty despite a variety of

single-interest groups who wish to eliminate the presence of family physicians in the hospital.³ Maternity care services are also threatened when unfair credentialing or a lack of consistent training makes the delivery of babies difficult for family physicians.⁴⁻⁶ The American Medical Association has been an advocate protector of specialty-neutral credentialing.

Despite heroic efforts the medical profession is at risk for being painted into a corner by academic health centers, managed care organizations, and government regulations. Even so, the tide can be turned if physicians speak up and act at the local level. Oppose discriminatory practices that place artificial barriers in front of cancer prevention services. Become involved at the grass roots level and encourage others to do the same.

Wm. MacMillan Rodney, MD
Department of Family Medicine
University of Tennessee, Memphis

References

1. Mulvey FR. Reimbursement for flexible sigmoidoscopy. *J Am Board Fam Pract* 1999;12:182.
2. Rodney WM. Flexible sigmoidoscopy: the unkept promise of cancer prevention. *Am Fam Physician* 1999;59:270, 273.
3. Rodney WM. Should any hospital-based training for family physicians persist? *Fam Med* 1998;30:398-9.
4. Rodney WM. Obstetrics-enhanced family practice: an endangered species worth saving! *Fla Fam Physician* 1993;43(1):8-9.
5. Harper MB, Mayeaux EJ Jr, Pope JB, Goel R. Procedural training in family practice residencies: current status and impact on resident recruitment. *J Am Board Fam Pract* 1995;8:189-94.
6. Sierpina VS, Volk RJ. Teaching procedures: most common settings, evaluation methods, and training barriers in family practice residencies. *Fam Med* 1998;30:421-3.
7. Weissman AN, Dawson JD, Fox DO. Variability in the learning experiences of family practice residents during an obstetric rotation. *Fam Med* 1999;31:28-33.