

I would like to suggest that this proposed strategy is inefficient, costly, and unlikely to be successful. An alternative strategy to offer preventive services during acute care visits will efficiently increase delivery to a greater proportion of patients at a lower cost. This latter strategy has proved to be effective in a single primary care practice^{3,4} and is currently being tested in a large community-based multisite, multispecialty group practice.² A randomized trial of the competing strategies would also appear to be feasible.

I also agree with Dr. Paul Frame, who stated in an accompanying editorial that "a system for delivering preventive services should be a requirement for accreditation of family practice residency programs."⁵

David L. Hahn, MD
Arcand Park Clinic
Madison, Wisc

References

1. Ruffin MT 4th, Gorenflo DW, Woodman B. Predictors of screening for breast, cervical, colorectal, and prostatic cancer among community-based primary care practices. *J Am Board Fam Pract* 2000;13:1-10.
2. Hahn DL, Olson N. The delivery of clinical preventive services: acute care intervention. *J Fam Pract* 1999;48:785-9.
3. Hahn DL, Berger MG. Implementation of a systematic health maintenance protocol in a private practice. *J Fam Pract* 1990;31:492-504.
4. Hahn DL. Systematic cholesterol screening during acute care visits. *J Am Board Fam Pract* 1993;6:529-36.
5. Frame PG. Implementing clinical preventive medicine: time to fish or cut bait. *J Am Board Fam Pract* 2000;13:84-5.

The above letter was referred to the first author of the article in question, who offers the following reply.

To the Editor: Dr. Hahn raises a valid issue. Is it feasible to promote visits to primary care offices just for preventive issues? Dr. Hahn has written that it is not feasible and still have time to care for sick patients.¹ This conclusion was arrived at by assuming that each primary care physician would spend 30 additional minutes each year performing a complete physical examination for approximately 54,000 patients currently seen for acute care. This conclusion, however, has several assumptions. The first is that the acute care visits would remain the same. There is no evidence to support this assumption, and many have hypothesized that acute care visits would decrease. The second is that a complete physical examination is preventive service. On the contrary, preventive services are not complete physical examinations; a complete physical examination has not proved to be an effective preventive service. Third, one appointment for preventive services might be enough to facilitate the delivery of preventive services in future acute care visits. The literature has reported that ever having been seen for a health maintenance examination is predictive of getting preventive services and being current. So, it might take

only one such visit to implement a system that can address preventive services at other contacts.

All of us who struggle in the field of increasing the delivery of preventive services must be cautious with our interpretation of the published data. The literature has many examples of interventions that made significant changes—in one office,²⁻⁵ in academic settings,^{6,7} when focused on one specific preventive service,⁸⁻¹⁰ or within a short period of time. In contrast, the large randomized controlled trials of theoretically sound interventions have shown no effects to minimal changes in the delivery of preventive services in community-based primary care offices across several years.¹¹⁻¹³ This was recently reconfirmed at the annual meeting of the North American Primary Care Research Group. Four presentations focused on randomized clinical trials of different interventions to increase the delivery of preventive services. All reported no effect to minimal increases. All the published and presented studies have not been trying to increase office visits solely for preventive services. All have taken the approach of increasing the delivery of preventive services at all encounters.

With these failures, I conclude it is time for some radical reexamination of preventive services and changing primary care practices. From this perspective, one questions Dr. Hahn's conclusions that it is not feasible to promote encounters only for preventive services. It might actually decrease acute care visits and increase preventive services. Dr. Hahn's limited trial warrants replication in larger settings for a longer period. In addition, our understanding of the black box of practice behavior and changing practice behaviors is in the infancy stage. There is a need for more basic research into the variables that contribute to the behaviors of a community-based primary care practice. This information will guide the next generation of interventions. I agree with Dr. Frame, it is time for residency accreditation agencies and practicing physician certification groups to focus on measures of health status among the populations served by family physicians of which preventive services delivered is critical.

Mack T. Ruffin IV, MD, MPH
University of Michigan
Ann Arbor

References

1. Hahn D, Olson N. The delivery of clinical preventive services: acute care intervention. *J Fam Pract* 1999;48:785-9.
2. Hahn DL, Berger MG. Implementation of a systematic health maintenance protocol in a private practice. *J Fam Pract* 1990;31:492-504.
3. Frame PS, Zimmer JG, Werth PL, Martens WB. Description of a computerized health maintenance tracking system for primary care practice. *Am J Prev Med* 1991;7:311-8.
4. Frame PS. Health maintenance in clinical practice: strategies and barriers. *Am Fam Physician* 1992;45:1192-200.
5. Frame PS, Zimmer JG, Werth PL, Hall WJ, Eberly SW. Computer-based vs manual health maintenance tracking. A controlled trial. *Arch Fam Med* 1994;3:581-8.

6. McPhee SJ, Richard RJ, Solkowitz SN. Performance of cancer screening in a university general internal medicine practice: comparison with the 1980 American Cancer Society Guidelines. *J Gen Intern Med* 1986;1:275-81.
7. McPhee SJ, Bird JA, Jenkins CN, Fordham D. Promoting cancer screening. A randomized, controlled trial of three interventions. *Arch Intern Med* 1989;149:1866-72.
8. Schapira DV, Kumar NB, Clark RA, Yag C. Mammography screening credit card and compliance. *Cancer* 1992;70:509-12.
9. Yarnall KS, Michener JL, Broadhead WE, Tse CK. Increasing compliance with mammography recommendations: health assessment forms. *J Fam Pract* 1993;36:59-64.
10. Eilers GM, Swanson TK. Using reminder systems to improve Papanicolaou test follow-up. An example of continuous quality improvement. *Arch Fam Med* 1993;2:1136-40.
11. Dietrich AJ, Tobin JN, Sox CH, et al. Cancer early-detection services in community health centers for the underserved. A randomized controlled trial. *Arch Fam Med* 1998;7:320-7.
12. Manfredi C, Czaja R, Freels S, Trubitt M, Warnecke R, Lacey L. Prescribe for health. Improving cancer screening in physician practices serving low-income and minority populations. *Arch Fam Med* 1998;7:329-37.
13. Williams RB, Boles M, Johnson RD. A patient-initiated system for preventive health care. A randomized trial in community-based primary care practices. *Arch Fam Med* 1998;7:338-45.