

and consultations provided by them were not included in the report. This study included 3 family physicians' data for 12 months and had a higher rate of consultation (3.9 percent) despite the exclusions listed above. The work of Pagel,⁶ a former fellow Huntsvillian, was of course well known to us. The topic by Pagel and Wood was the heroic effort in remote Alaska where there was no system of roads and focused on the issue of air transport of sick patients. Dr. Lawler's own work,⁷ which was published after ours was written, again reports the experience from the Fulton, Missouri, training site. Consultations from faculty and a nurse practitioner were excluded, and 3 years of data, including 25,000 patient visits, were included. His finding of a referral rate of 1.31 percent is again very similar to ours.

It is truly heartening to see one's published work carefully scrutinized, as evidenced by Dr. Lawler's effort to suggest a more complete literature survey. It is the mark of a maturing discipline that active scientific debate occurs in the pages of its best journals. We hope this review has added to the reader's understanding of this important topic, and we are pleased to agree with Dr. Young that our report, with 9 years of data and almost 178,000 patient visits, "represents one of the largest reported series of observations regarding outpatient consultations emanating from a family practice teaching program."

William J. Crump, M.D.

Patricia Massengill, B.S.

University of Alabama in Huntsville
Huntsville, AL

References

1. Glenn JK, Hofmeister RW, Neikirk H, Wright H. Continuity of care in the referral process: an analysis of family physicians' expectations of consultants. *J Fam Pract* 1983; 16:329-34.
2. Dolezel JM, Amundson LH, Sinning NJ, Hoody HJ. PriCare and ambulatory referrals. *Cont Educ* 1980; 12:84-94.
3. Hines RM, Curry DJ. The consultation process and physician satisfaction: review of referral patterns in three urban family practice units. *Can Med Assoc J* 1978; 118:1065-73.
4. Moscovice I, Schwartz CW, Shortell SM. Referral patterns of family physicians in an underserved area. *J Fam Pract* 1979; 9:677-82.
5. Mayer TR. Family practice referral patterns in a health maintenance organization. *J Fam Pract* 1982; 14:315-9.
6. Pagel J, Wood T. Transport patterns and complications in an isolated Alaska practice. *J Fam Pract* 1983; 16:957-62.
7. Lawler FH. Referral rates of senior family practice residents in an ambulatory care clinic. *J Med Educ* 1987; 62:177-82.
8. Young PR. Editorial comment. *J Am Bd Fam Pract* 1988; 1:166.

Flexible Sigmoidoscopy

To the Editor: In their article on "Flexible Sigmoidoscopy" in the July-September 1988 issue, Dr. John E. Hocutt, Jr., et al. point out the many advantages to the family physician for performing flexible sigmoidoscopy on his or her patients.¹ I was alarmed, however, that they seem to imply that attending one of the numerous 1-day seminars in flexible sigmoidoscopy might qualify one to begin performing the procedure on patients. Many authors have shown that the procedure requires a number of supervised examinations before the examiner exhibits competence. In fact, the argument has revolved around just how many supervised procedures are necessary before performing the examination alone. Merely performing an examination does not necessarily mean that it was done properly. And with greater charges for flexible sigmoidoscopy versus rigid, how does one justify a limited or incomplete examination done while "self-training?"

With the ever increasing pressure about documentation for privileges, quality of medical care issues, and the competition among specialties, we as family physicians do not want to encourage our members to perform procedures without adequate training. Certainly, the Academy recognizes the need to promote hands-on training for flexible sigmoidoscopy because it expended a great deal of effort in setting up an extensive network of preceptors. Therefore, I would urge family physicians who wish to perform flexible sigmoidoscopy in their practice to arrange for hands-on training, if not through the Academy's programs, perhaps with the help of the faculty of a nearby family practice residency program.

Steven J. Glinka, M.D.

Ventura County Medical Center
Ventura, CA

References

1. Hocutt JE Jr, Hainer B, Jackson M. Flexible fiberoptic sigmoidoscopy: its use in family medicine. *J Am Bd Fam Pract* 1988; 1:189-93.

The above letter was referred to the authors of the article in question, who offer the following reply: