symptom attacks.¹⁶ Therefore, it may be that Fava, et al. failed to identify infrequent or limited symptom attacks that began prior to the onset of phobic avoidance. In my study, although there were several patients in whom the onset of panic attacks and phobic avoidance coincided, the remainder of patients had a significant lag time (mean = 3.18 years) between the onset of panic attacks and phobic axis and phobic attacks and phobic attacks.

The ultimate answer to the natural history of panic disorder and agoraphobia will have to wait for longitudinal studies in which patients at risk for the development of panic disorder/agoraphobia are followed over time for the development of each condition.

> David A. Katerndahl, M.D. The University of Texas Health Science Center San Antonio, TX

References

- 1. Klein DF. Anxiety reconceptualized. In: Klein DF, Rabkin JG, eds. Anxiety: new research and changing concepts. New York: Raven Press, 1981:235-65.
- Garvey MJ, Tuason VB. The relationship of panic disorder to agoraphobia. Compr Psychiatry 1984; 25: 529-31.
- 3. Torgersen S. Genetic factors in anxiety disorders. Arch Gen Psychiatry 1983; 40:1085-9.
- Noyes R Jr, Crowe RR, Harris EL, Hamra BJ, McChesney CM, Chaudhry DR. Relationship between panic disorder and agoraphobia. A family study. Arch Gen Psychiatry 1986; 43;227-32.
- Thyer BA, Himle J, Curtis GC, Cameron OG, Nesse RM. A comparison of panic disorder and agoraphobia with panic attacks. Compr Psychiatry 1985; 26:208-14.
- 6. Klein DF, Ross DC, Cohen P. Panic and avoidance in agoraphobia. Application of path analysis to treatment studies. Arch Gen Psychiatry 1987; 44:377-85.
- Turner SM, Williams SL, Beidel DC, Mezzich JE. Panic disorder and agoraphobia with panic attacks: covariation along the dimensions of panic agoraphobic fear. J Abnorm Psychol 1986; 95:384-8.
- 8. Gelder MG, Marks IM. Severe agoraphobia: a controlled prospective trial of behaviour therapy. Br J Psychiatry 1966; 112:309-19.
- 9. Klein DF. Panic attacks in phobia treatment studies [letter]. Arch Gen Psychiatry 1983; 40:1151-2.
- Marks I. Panic attacks in phobic treatment studies [letter]. Arch Gen Psychiatry 1983; 40:1151.
- Fava GA, Grandi S, Canestrari R. Prodromal symptoms in panic disorder with agoraphobia. Am J Psychiatry 1988; 145:1564-7.
- 12. Sheehan DV, Sheehan KH. Classification of anxiety and hysterical states. Part II. Toward a more heuristic classification. J Clin Psychopharmacol 1982; 2:386-93.
- 13. Katerndahl DA. Significance of infrequent and limited symptom panic attacks. J Nerv Ment Dis (in press).
- 14. Norton GR, Harrison B, Hauch J, Rhodes L. Characteristics of people with infrequent panic attacks. J Abnorm Psychol 1985; 94:216-21.
- Sheehan DV, Coleman JH, Greenblatt DJ, et al. Some biochemical correlates of panic attacks with agoraphobia and their response to a new treatment. J Clin Psychopharmacol 1984; 4:66-75.
- Weissman MM, Leaf PJ, Holzer CE 3d, Merikangas KR. The epidemiology of anxiety disorders: a highlight of recent evidence. Psychopharmacol Bull 1985; 21:538-41.

Postpartum Pap Smear

To the Editor: In the article by Weiss, et al., "The Postpartum Papanicolaou Smear," in the January-March 1989 issue, cases of abnormal cervical cytology on the postpartum examination that were not predicted on the prenatal Pap results were presented. While I support the conclusion of the study, to perform Pap smears during prenatal care and again at the postpartum examination, even when the prenatal Pap is normal, I question the 4.9 percent "conversion" rate to abnormal cytology. When defining specimen collection techniques and interpretation of results, no mention was made of assessing adequacy of smears by the presence or absence of endocervical cells. The presence of endocervical cells is an important indicator of the adequacy of a Pap smear.¹ One can only be sure that the entire transformation zone has been accurately sampled if endocervical cells are present on the smear.² Furthermore, the rate of epithelial abnormalities has been reported to be lower in smears that contain no endocervical cells.³ With pregnancy, changes occur in the anatomy of the cervix, and the endocervical canal is filled with a tenacious mucus that can block access to the columnar cells underneath, causing a lower yield of endocervical cells in pregnant women.4

Because the progression from normal endocervical cells to dysplasia to carcinoma-in-situ is more rapid than formerly thought, Dr. Weiss's study could hold important clinical implications. However, without the adequacy of sampling technique assessed, one would wonder about the degree of false-negative readings on the prenatal examination in the study group. This would potentially lessen the impact of the conclusion drawn in this paper.

> Elizabeth G. Baxley, M.D. Anderson, SC

References

- 1. American College of Obstetricians and Gynecologists. Cervical cytology. Technical Bulletin 81. Chicago: American College of Obstetricians and Gynecologists, October 1984.
- Gondos B, Marshall D, Ostergard DR. Endocervical cells in cervical smears. Am J Obstet Gynecol 1972; 114: 833-4.
- 3. Elias A, Linthorst G, Bekker B, Vooijs PG. The significance of endocervical cells in the diagnosis of cervical epithelial changes. Acta Cytol 1983; 27:225.
- Hamblin JE, Brock CD, Litchfield L, Dias J. Papanicolaou smear adequacy: effect of different techniques in specific fertility states. J Fam Pract 1985; 20:257-60.

The above letter was referred to the authors of the article in question, who offer the following reply.

To the Editor: Dr. Baxley is correct. The apparent "conversion" of some normal Pap smears into abnormal Pap smears during the course of pregnancy may, in reality,