

CLINICAL REVIEW

The Well-Man Visit: Addressing a Man's Health to Optimize Pregnancy Outcomes

Jonathan N. Warner, MD, and Keith A. Frey, MD

Male health and health care before conception has received little attention by many physicians. Only recently has the importance of the health of men before conception been explored. The benefits of preconception care are important, including ensuring that all pregnancies are planned and wanted, improving the man's genetic and biologic contributions to the pregnancy, improving reproductive health and outcomes for female partners, improving the man's capacity for and understanding of parenthood, and enhancing the man's overall health through access to primary care. (J Am Board Fam Med 2013;26: 196–202.)

Keywords: Men's Health, Preconception Care

Male health and health care before conception has received little attention by many practitioners. It is also rare for a male patient to come to a clinic for preconception information. Only recently has the importance of preconception health for men been explored.^{1,2} The benefits of preconception care are important, including ensuring that all pregnancies are planned and wanted, improving the man's genetic and biologic contributions to the pregnancy, improving reproductive health and outcomes for female partners, improving the male's capacity and understanding of parenthood, and enhancing the man's overall health through access to primary care.² Health care professionals evaluating and advising men regarding their overall reproductive health need a comprehensive framework, which is outlined herein.

Evaluation

Reproduction Life Plan

First, it is important to assess the goals and intentions of the male patient regarding reproduction

goals. The Centers for Disease and Control and Prevention recommends that all men, women, and couples to have a reproductive life plan.³ This is defined as a set of goals about having children or not having children based on personal values and resources and a plan to achieve these goals. Like women,⁴ the male patient should be questioned regarding any previous children, the desire to have any more children, whether there has ever been trouble conceiving, and what time frame the couple has for conception. A recent study noted that <10% of men can recall their primary care physician discussing preconception health care with them.⁵ If the couple is not currently interested in having children, it is important that the male patient be involved with contraception, and a discussion regarding proper methods should be addressed.

Maximizing the Health of the Male Patient

To maximize outcomes, herein we aid in the assessment of the male patient who is considering having a child. If the patient has been trying to conceive without results for some time, a formal fertility evaluation is warranted for both the male patient (by a male infertility specialist) and his partner.

Medical and Surgical History

As with all patient encounters, a detailed medical and surgical history is important for the evaluation of a man's reproductive health. Tables 1 and 2 list

This article was externally peer reviewed.
Submitted 1 June 2012; revised 2 November 2012; accepted 7 November 2012.


From the Mayo Clinic, Thunderbird Primary Care Center, Scottsdale, AZ.

Funding: none.

Conflict of interest: none declared.

Corresponding author: Keith A. Frey, MD, Mayo Clinic, Thunderbird Primary Care Center, 13737 N 92nd St, Scottsdale AZ 85260 (E-mail: Frey.Keith@mayo.edu).

Figure 1. Discussion outline for male reproduction life plan. FSH, follicle stimulating hormone; HCG, human chorionic gonadotropin; TSH, thyroid stimulating hormone.



several medical and surgical conditions that are known to impact fertility. If these conditions exist, it is prudent to perform quantitative and qualitative analyses of semen. If abnormalities are found, referral to an infertility specialist is warranted. It is important to evaluate the patient's mental health as well. Although this may not directly impact the man's ability to father a child, there is

evidence that the child of a father with depression may have behavioral and emotional developmental problems.²¹

Medications

The patient's medication history should be evaluated, including past and current prescription medications, nonprescription medications, herbal remedies, and

Table 1. Medical Conditions Associated With Infertility

Cryptorchidism, testicular torsion, and trauma ⁶
Postpubertal mumps orchitis ⁷
Congenital adrenal hyperplasia
Obesity ⁸
Diabetes and poor glycemic control ⁹
Erectile dysfunction ^{10,11}
Multiple sclerosis and spinal cord injury ¹²
Hyper or hypothyroidism ¹³
Varicocele ¹⁴
Prostatitis, epididymitis, sexually transmitted diseases ¹⁵
Malignancies (Hodgkin ¹⁶ and testicular cancer ¹⁷) and any endocrine or metabolic effects of cancer ¹⁸

recreational drugs. If the patient is taking medications that are known to impact fertility (Table 3), an alternative should be sought or the patient should be monitored appropriately. Medications may alter the hypothalamic-pituitary-gonadal axis, impair libido, affect erectile function, or have a direct toxic effect on the sperm.

Family History and Genetic Risks

As part of the family history, a detailed genetic history should be obtained. Because sex-linked or autosomal disorders may skip generations, a 3-generation history is important. Conditions to screen for include cystic fibrosis, which impairs infertility through a congenital absence of the vas deferens,²⁸ and Klinefelter syndrome, which impairs fertility through sclerosis of the seminiferous tubules, resulting in azoospermia and small testicles.²⁹ Families with intersex disorders should lead to the evaluation of the possibility of androgen receptor abnormalities.

Social History

As noted earlier, the risks of tobacco, alcohol, and illicit drug use can directly affect spermatogenesis and impact erectile function. However, a social history is also important to determine whether the patient is exposed to environmental hazards, including ionizing radiation, radiant heat, pollutants in the diet, lead, mercury, dichlorodiphenyltrichloroethane, carbon disulfide, ethylene glycol ethers, dibromochloropropane, polycyclic aromatic hydrocarbons, polychlorinated biphenyls, dioxins, phthalates, and acrylamide.³⁰

A review of patient's hobbies should also be performed. Woodworking with oils used to finish

furniture, painting, pottery with lead-based glazes, stained glass work, and gun cleaning and shooting may lead to toxic exposures to lead and other heavy metals that are harmful to sperm production.

Physical Examination

A thorough physical examination should be performed as part of the preconception evaluation. This includes screening for hypertension and obesity through body mass index measurements. Overall body habitus should also be evaluated because it can offer clues about androgen abnormalities. Tall stature, gynecomastia, and decreased body hair could suggest Klinefelter syndrome. Hyperpigmentation may be a sign of underlying adrenal insufficiency; whereas abdominal striae, increased abdominal girth, and enlargement of the fat pad along the back of the neck and collar bone could be suggestive of Cushing syndrome. Excess sweating and essential tremor could be indicative of hyperthyroidism.

An examination of visual fields should be performed; hemianopsia could indicate a pituitary adenoma. The thyroid gland should be evaluated because enlargement could be indicative of underlying hyper- or hypothyroidism.

Gynecomastia may be a sign of Klinefelter disease, excess estrogen, or androgen insensitivity, which may be associated with infertility. Underlying respiratory disease may be indicative of ciliary disease such as Kartagener syndrome, cystic fibrosis, or Young syndrome, which may inhibit fertilization through absence or obstruction of the vas deferens. A cardiovascular examination should be performed to identify any underlying arterial compromise, which may contribute to erectile dysfunction.

An abdominal examination should be performed to evaluate for any surgical scars, particularly in the lower abdomen or inguinal region, for any evidence of a previous operation that may have affected the patency of the vasa. Hepatomegaly may indicate un-

Table 2. Surgical Conditions Associated With Infertility

Inguinal hernia repairs ¹⁹
Scrotal or pelvic surgery
Retroperitoneal surgery in the setting of testicular cancer ²⁰
Incision of the bladder neck and transurethral resection of the prostate

Table 3. Medications That Can Affect Fertility

Medication	Effect
Exogenous testosterone	Decreased spermatogenesis
Androgen supplements	Impaired HPG axis
5- α reductase inhibition	ED, impaired sperm production ²²
α -blockers	Retrograde ejaculation
Cimetidine, spirinolactone, and ketoconazole	Direct inhibition of androgen production
Erythromycin, neomycin, tetracycline, nitrofurantoin, and gentamycin	Direct impairment of sperm production
Selective serotonin receptor inhibitors, monoamine oxidase inhibitors, lithium, and phenothiazines	Decreased libido, ED, and alteration of the HPG axis ²³
Chemotherapeutic agents	Direct impairment of spermatogenesis
Marijuana, ²⁴ cocaine, ²⁵ alcohol, ²⁶ and tobacco ²⁷	May impair the HPG axis, spermatogenesis, and erectile function

ED, erectile dysfunction; HPG, hypothalamic-pituitary-gonadal.

derlying hepatic dysfunction, which may impair the metabolism of sex steroids.

A genital examination should be performed (Table 4), beginning with the phallus. Penile lesions could indicate an underlying sexually transmitted disease. Hypospadias, penile curvature, or chordee may impact the delivery of semen during intercourse. The scrotal skin and contents should be evaluated. The presence of scars could indicate previous scrotal surgery, which may impact patency of the vas deferens. Small testicles could indicate testicular failure, either primarily as is seen in Klinefelter syndrome, or secondarily as seen with use of exogenous testosterone. Normal testicular size is 4 \times 3 cm or a 20-mL volume.³¹ Undescended testicles or absent testicles should also be noted. Testicular masses may indicate an underlying neoplasm. Epididymal tenderness may be indicative of untreated epididymitis, which can affect sperm delivery. The presence of the vas deferens should be evaluated to rule out congenital absence of the vas deferens. Vasal nodularity or thickening could indicate a prior infection such as tuberculosis, whereas thickening may be indicative of obstruction. Finally, the patient should be examined in the supine and standing positions to determine the presence of a varicocele. Varicoceles are seen in 15% of normal men and are present in 19% to 41% of men with primary infertility and up to 81% of those with secondary infertility.¹⁴

Laboratory Testing

Basic screening, as recommended by the U.S. Preventive Service Task Force, should be offered to all male patients. This includes screening for lipid disorders beginning at age 35 (and for men ages 20

to 35 if they have diabetes, a family history of coronary artery disease or familial hyperlipidemia, or multiple risk factors for coronary heart disease) and screening for diabetes.³²

Targeted laboratory evaluation can be directed toward specific clinical signs and symptoms. Prolactin levels may be checked if there is any evidence of bitemporal hemianopsia. Thyroid function tests should be ordered if there are signs and symptoms of hyper- or hypothyroidism. Adrenal function tests may be necessary with any signs of Addison or Cushing syndrome. Liver function testing should be completed if there is evidence of hepatomegaly. Testosterone, estrogen, follicle-stimulating hormone, and luteinizing hormone should be evaluated if there are small testicles, gynecomastia, eunuchoid body habitus, or signs of androgen resistance.

Recommendations for Health Promotion

Achieving and maintaining a healthy weight is the starting point for discussion of health promotion. Obesity has a number of detrimental health effects. In fact, every 20 lb overweight can lead to a 10% increase in the odds of infertility.⁸ As part of the preconception evaluation, overweight and obese men should be counseled on the health and fertility benefits of weight loss; they should set goals and be offered the resources to achieve those goals. This may include referral to a nutritionist, a structured weight loss program, and promoting regular exercise.

Stress reduction is also important to emphasize in male patients. Although the mechanisms are not fully understood, there is a clear impact of stress on

Table 4. Physical Examination

Examination	Possible Condition
General	
Tall stature	Klinefelter syndrome
Long arms	Klinefelter syndrome
Absence of body hair	Adrenal disease
Hyperpigmentation	Adrenal disease
Obesity	Impaired testosterone and high estrogen
HEENT	
Bitemporal hemianopsia	Pituitary adenoma
Goiter	Thyroid disease
Anosmia	Kallman syndrome
Chest	
Gynecomastia	Klinefelter syndrome or HPG axis defect
Buffalo hump	Adrenal disease
Respiratory rales and crackles	Cystic fibrosis or celiac disease
Abdomen	
Surgical scars in the pelvis	Possible nerve damage or vasal damage
Inguinal scars	Vasal obstruction
Striae	Adrenal disease
Truncal obesity	Adrenal disease
Hepatomegaly	Liver disease
Genitals	
Penile lesions	Possible STD, which impacts fertility
Inguinal adenopathy	Possible infection or malignancy
Impalpable vas deferens	Possible CBAVD or cystic fibrosis
Vas deferens nodularity	Possible vasal obstruction
Varicocele	Impairs fertility
Scrotal scars	Possible testicular or vasal injury
Epididymitis, prostatitis	Active infection can impair fertility

CBAVD, congenital absence of the vas deferens; HEENT, head, ears, eyes, nose, and throat; HPG, hypothalamic-pituitary-gonadal; STD, sexually transmitted disease.

sperm morphology³³ and concentration³⁴ and decreasing testosterone.³⁵ The reason for these changes is believed to be mediated by impacts on the hypothalamic-pituitary-gonadal axis secondary to increased release of epinephrine and corticotrophin-releasing hormone, which impacts the natural pulsatility of luteinizing hormone.^{36,37} Approaches to stress reduction include regular exercise, proper sleep habits, and proper nutrition.

The benefits of avoidance of tobacco, alcohol, and recreational drugs have been noted. Providers should

also encourage those patients with a profession with hazardous environmental exposures to take appropriate precautions to minimize risk. This includes the use of leaded aprons for those with regular exposure to x-rays and proper use of gloves and masks when exposed to harmful toxins. Nutritional recommendations are not only an important part of helping a patient maintain a healthy weight and reducing stress, but certain nutrients actually have been shown to affect the viability of sperm; zinc sulfate and folic acid have shown a positive effect on sperm production.³⁸

Additional Considerations

Men with findings during a physical examination that suggest hormonal or testicular abnormalities should be referred to an infertility expert. In addition, the couple who has been attempting unsuccessfully to conceive should undergo an infertility evaluation.³⁹ It is vital that male patients be included in such an infertility evaluation.^{39,40}

Male patients should also be counseled regarding the growing evidence of increased risks associated with advanced paternal age. The risk of a child with schizophrenia is twice as likely if the father is 45 or older compared with those in their 20s and 3 times more likely after the paternal age of 50.⁴¹ Autism is 5.75 times more likely if the father is aged 40 or older compared with those men who father a child when they are younger than 40.^{41,42} Autosomal dominant disorders also have been associated with advanced paternal age, including achondroplasia, Marfan syndrome, and osteogenesis imperfecta.⁴³

Conclusions

Preconception care and planning is important for improving pregnancy outcomes, ensuring all pregnancies are planned and wanted, and enhancing the health of female partners and, particular to this review, of the male patient. The preconception examination is an excellent window to promote not only reproductive health but also to discuss overall health issues with the male patient. This includes promotion of proper nutrition and regular exercise, treatment and prevention of diseases, stress reduction, and avoidance of dangerous behaviors.

References

1. Frey KA. Male reproductive health and infertility. *Prim Care* 2010;37:643–52, x.
2. Frey KA, Navarro SM, Kotelchuck M, Lu MC. The clinical content of preconception care: preconcep-

- tion care for men. *Am J Obstet Gynecol* 2008;199: S389–95.
3. Johnson K, Posner SF, Biermann J, et al; CDC/ATSDR Preconception Care Work Group; Select Panel on Preconception Care. Recommendations to improve preconception health and health care—United States. A report of the CDC/ATSDR Preconception Care Work Group and the Select Panel on Preconception Care. *MMWR Recomm Rep* 2006;55(RR-6):1–23.
4. Files JA, Frey KA, David PS, Hunt KS, Noble BN, Mayer AP. Developing a reproductive life plan. *J Midwifery Womens Health* 2011;56:468–74.
5. Frey KA, Engle R, Noble BB. Preconception health-care: what do men know and believe? *J Mens Health* 2012;9:25–35.
6. Puri P, Barton D, O'Donnell B. Prepubertal testicular torsion: subsequent fertility. *J Pediatr Surg* 1985;20:598–601.
7. Werner CA. Mumps orchitis and testicular atrophy; a factor in male sterility. *Ann Intern Med* 1950;32: 1075–86.
8. Sallmen M, Sandler DP, Hoppin JA, Blair A, Baird DD. Reduced fertility among overweight and obese men. *Epidemiology* 2006;17:520–3.
9. Padron RS, Dambay A, Suarez R, Mas J. Semen analyses in adolescent diabetic patients. *Acta Diabetol Lat* 1984;21:115–21.
10. McCulloch DK, Campbell IW, Wu FC, Prescott RJ, Clarke BF. The prevalence of diabetic impotence. *Diabetologia* 1980;18:279–83.
11. Sexton WJ, Jarow JP. Effect of diabetes mellitus upon male reproductive function. *Urology* 1997;49: 508–13.
12. Sonksen J, Biering-Sorensen F. Fertility in men with spinal cord or cauda equina lesions. *Semin Neurol* 1992;12:106–14.
13. Velazquez EM, Bellabarba Arata G. Effects of thyroid status on pituitary gonadotropin and testicular reserve in men. *Arch Androl*. 1997;38:85–92.
14. Agarwal A, Deepinder F, Cocuzza M, et al. Efficacy of varicocelectomy in improving semen parameters: new meta-analytical approach. *Urology* 2007;70:532–8.
15. Aitken RJ, De Iuliis GN. Value of DNA integrity assays for fertility evaluation. *Soc Reprod Fertil Suppl* 2007;65:81–92.
16. Rueffer U, Breuer K, Josting A, et al. Male gonadal dysfunction in patients with Hodgkin's disease prior to treatment. *Ann Oncol* 2001;12:1307–11.
17. Petersen PM, Skakkebaek NE, Vistisen K, Rorth M, Giwercman A. Semen quality and reproductive hormones before orchiectomy in men with testicular cancer. *J Clin Oncol* 1999;17:941–7.
18. Costabile RA, Spevak M. Cancer and male factor infertility. *Oncology (Williston Park)* 1998;12:557–62, 565; discussion 566–8, 570.
19. Shin D, Lipshultz LI, Goldstein M, et al. Herniorrhaphy with polypropylene mesh causing inguinal vasal obstruction: a preventable cause of obstructive azoospermia. *Ann Surg* 2005;241:553–8.
20. Kedia KR, Markland C, Fraley EE. Sexual function after high retroperitoneal lymphadenectomy. *Urol Clin North Am* 1977;4:523–8.
21. Ramchandani P, Stein A, Evans J, O'Connor TG. Paternal depression in the postnatal period and child development: a prospective population study. *Lancet* 2005;365:2201–5.
22. Giuliano F. Impact of medical treatments for benign prostatic hyperplasia on sexual function. *BJU Int* 2006;97(Suppl 2):34–8; discussion 44–5.
23. Nudell DM, Monoski MM, Lipshultz LI. Common medications and drugs: how they affect male fertility. *Urol Clin North Am* 2002;29:965–73.
24. Close CE, Roberts PL, Berger RE. Cigarettes, alcohol and marijuana are related to pyospermia in infertile men. *J Urol* 1990;144:900–3.
25. Hurd WW, Kelly MS, Ohl DA, Gauvin JM, Smith AJ, Cummins CA. The effect of cocaine on sperm motility characteristics and bovine cervical mucus penetration. *Fertil Steril* 1992;57:178–82.
26. Muthusami KR, Chinnaswamy P. Effect of chronic alcoholism on male fertility hormones and semen quality. *Fertil Steril* 2005;84:919–24.
27. Vine MF, Tse CK, Hu P, Truong KY. Cigarette smoking and semen quality. *Fertil Steril* 1996;65: 835–42.
28. Anguiano A, Oates RD, Amos JA, et al. Congenital bilateral absence of the vas deferens. A primarily genital form of cystic fibrosis. *JAMA* 1992;267: 1794–7.
29. Lanfranco F, Kamischke A, Zitzmann M, Nieschlag E. Klinefelter's syndrome. *Lancet* 2004;364:273–83.
30. Bonde JP. Male reproductive organs are at risk from environmental hazards. *Asian J Androl* 2010;12: 152–6.
31. Charny CW. The spermatogenic potential of the undescended testis before and after treatment. *J Urol* 1960;83:697–705.
32. Guide to Clinical Preventive Services, 2nd edition. Rockville, MD: Agency for Healthcare Research and Quality; 2010. Available from: <http://odphp.osophs.dhhs.gov/pubs/guidecps/>. Accessed January 3, 2013.
33. Giblin PT, Poland ML, Moghissi KS, Ager JW, Olson JM. Effects of stress and characteristic adaptability on semen quality in healthy men. *Fertil Steril* 1988;49:127–32.
34. Zorn B, Auger J, Velikonja V, Kolbezen M, Meden-Vrtovec H. Psychological factors in male partners of infertile couples: relationship with semen quality and early miscarriage. *Int J Androl* 2008;31:557–64.
35. Nilsson PM, Moller L, Solstad K. Adverse effects of psychosocial stress on gonadal function and insulin

- levels in middle-aged males. *J Intern Med* 1995;237: 479–86.
36. McGrady AV. Effects of psychological stress on male reproduction: a review. *Arch Androl* 1984;13:1–7.
37. Schweiger U, Deuschle M, Weber B, et al. Testosterone, gonadotropin, and cortisol secretion in male patients with major depression. *Psychosom Med* 1999;61:292–6.
38. Wong WY, Merkus HM, Thomas CM, Menkveld R, Zielhuis GA, Steegers-Theunissen RP. Effects of folic acid and zinc sulfate on male factor subfertility: a double-blind, randomized, placebo-controlled trial. *Fertil Steril* 2002;77:491–8.
39. Jose-Miller AB, Boyden JW, Frey KA. Infertility. *Am Fam Physician* 2007;75:849–56.
40. Kolettis PN. Evaluation of the subfertile man. *Am Fam Physician* 2003;67:2165–72.
41. Malaspina D, Harlap S, Fennig S, et al. Advancing paternal age and the risk of schizophrenia. *Arch Gen Psychiatry* 2001;58:361–7.
42. Reichenberg A, Gross R, Weiser M, et al. Advancing paternal age and autism. *Arch Gen Psychiatry* 2006; 63:1026–32.
43. Stewart AF, Kim ED. Fertility concerns for the aging male. *Urology* 2011;78:496–9.