

ORIGINAL RESEARCH

Effects of Ginger for Nausea and Vomiting in Early Pregnancy: A Meta-Analysis

Maggie Thomson, MD, Renee Corbin, MSc, and Lawrence Leung, MBBChir, MFMClin

Background: Nausea and vomiting in early pregnancy (NVEP) is commonly encountered in family medicine. Ginger (*Zingiber officinale*) is a popular nonpharmacological treatment but consensus of its use is lacking.

Methods: We conducted a meta-analysis of clinical trials using ginger for NVEP as published in PubMed and EMBASE, CINAHL, Cochrane Library, and all EBM reviews. Studies satisfying 3 criteria were selected: (1) randomized placebo-controlled design; (2) use of ginger or *Z. officinale*; and (3) extractable data on improvement in NVEP. Data were synthesized into pooled odd ratios based on the random effects model, and results were tabulated with the aid of Forest plots.

Results: We identified 135 potentially relevant records; only 6 studies met the final criteria. Of the total 508 subjects, 256 and 252 subjects were randomly assigned to receive ginger and placebo, respectively. The use of ginger (~1 g daily) for at least 4 days is associated with a 5-fold likelihood of improvement in NVEP. Heterogeneity among the clinical studies were acknowledged in the final interpretation of results.

Conclusions: Despite the widespread use of ginger in the diet, its clinic value and safety profile in treating NVEP is still unknown. Our meta-analysis suggests that ginger is an effective nonpharmacological treatment for NVEP. (J Am Board Fam Med 2014;27:115–122.)

Keywords: Alternative Medicine, Complementary Medicine, Pregnancy, Prenatal Care, Primary Health Care

Pregnancy-induced nausea and vomiting is commonly seen in family medicine, and 50% to 90% of pregnancies are affected by nausea with or without vomiting. According to a recent study, up to 63% of women experience nausea and vomiting up to 24 weeks' gestation.¹ While only 0.3% to 2% of these cases are considered severe (called hyperemesis gravidarum, leading to a loss of >5% of prepregnancy body weight), all forms of pregnancy-related nausea can affect quality of life.² Compared with the volume of literature regarding the pathogenesis and treatment of pregnancy-induced nausea and

vomiting in general, few studies look at nausea and vomiting in early pregnancy (NVEP).

Epidemiology, Risk Factors, and Pathogenesis of NVEP

Nausea and vomiting during pregnancy is known to be more common in younger primigravid women, as well as in Western countries and urban areas.² Risk factors for its development include history of estrogen-based medication causing nausea and motion or migraine causing nausea. Multiple gestation pregnancies,³ women who did not take multivitamins before conception,⁴ those with acid reflux,⁵ and those with a hydatidiform mole⁶ also are known to be at increased risk. There seems to be a genetic component to developing hyperemesis gravidarum; several studies demonstrate that daughters of women who experienced the condition are at increased risks.^{7,8} It is interesting that alcohol consumption and cigarette smoking have both been demonstrated to be protective.⁹ The exact pathogenesis of NVEP remains unclear; however, hormonal changes including elevated serum human chorionic gonado-

This article was externally peer reviewed.

Submitted 29 May 2013; revised 17 July 2013; accepted 22 July 2013.

From the Department of Family Medicine (MT, LL) and the Centre of Studies in Primary Care (RC, LL), Queen's University, Kingston, Ontario, Canada.

Funding: none.

Conflict of interest: none declared.

Corresponding author: Lawrence Leung, MBBChir, MFMClin, Department of Family Medicine, Queen's University, 220 Bagot St, PO Bag 8888, Kingston ON K7L 5E9, Canada (E-mail: leungl@queensu.ca).

tropin have been implicated,² as have psychological factors and stress response.¹⁰ Delayed or dysrhythmic gastric motility also has been postulated as a likely cause of NVEP.²

Diagnosis and Treatment of NVEP

Diagnosis of NVEP is purely clinical. There exists no defined diagnostic criteria for the condition. Mean onset is known to be between 5 to 6 weeks of gestation, with the majority of symptoms resolving by 20 weeks.² Approximately 20% of women continue to have symptoms into their second and third trimesters.² Contrary to popular belief, symptoms occur throughout the day and are often not limited to the morning as the term *morning sickness* would imply.²

Goals of treatment include reducing maternal symptoms and complications and mitigating any effects on the fetus. Antiemetics for NVEP that have been studied include Dicletin (Duchesnay Inc., Blainville, QC, Canada), ondansetron, ginger, various antihistamines, vitamin B₆, metoclopramide, and phenothiazines.² A recent randomized, double-blind, placebo-controlled trial¹¹ demonstrated that Dicletin was effective for nausea and vomiting during pregnancy, and this drug is currently recommended as first-line pharmacological treatment for NVEP in Canada. Ondansetron is commonly used as an antiemetic during chemotherapy, with good evidence.¹² However, it has been used for some time in the United States as a first-line (albeit off-label) treatment for pregnancy-related nausea and vomiting.¹³ While evidence exists for other pharmacological and nonpharmacological treatments (eg, dietary adjustments, acupressure, and acupuncture), they are beyond the scope of this article. Our study aimed to critically examine the efficacy of ginger in the treatment of NVEP through meta-analysis of published clinical trials.

Ginger and Its Medicinal Effects

The antinausea effect of ginger was first described in one of the canons of traditional Chinese medicine—the Synopsis of the Prescriptions of Golden Chamber¹⁴—in 200 AD. Ginger is the underground stem (or rhizome) of the perennial plant *Zingiber officinale*, which is indigenous to China and India but is cultivated all over the world. From the body of ginger sprouts the pseudostems, which

branch off to leaves that can reach 2 feet in height.¹⁵ Analysis of ginger reveals 2 major classes of phytochemicals: volatile oils, which give ginger its pleasant smell,¹⁶ and the nonvolatile compounds (eg, gingerols and zingerones), which account for its piquant taste and its pharmacological effects.¹⁷ Many studies of the antiemetic nature of ginger for various conditions have been published but with mixed results. While evidence supports benefits of ginger for seasickness,¹⁸ motion sickness,¹⁹ and postoperative nausea and vomiting,²⁰ its use in preventing chemotherapy-induced nausea and vomiting is still conflicting.^{21,22} Ginger also has been demonstrated to be effective in treating nausea and vomiting in pregnancy to the extent that it is as effective as vitamin B₆ alone.^{23–27} The exact antiemetic mechanisms of ginger are still unknown, but in vitro studies revealed antagonistic effects of gingerols on serotonergic 5-HT₃^{28,29} and cholinergic M receptors.²⁹ So far, no direct adverse effects on human fetuses or the course of pregnancy have been demonstrated.^{24–27} However, there had been concerns about interfering with fetal development, which led to the issuance within Finland and Denmark of warning labels for all supplements containing ginger.³⁰ It is also recognized that ginger has potent anticoagulant effects, which may enhance bleeding and miscarriages and interact with other medications.³⁰ At present, there are no large-scale studies ascertaining the safety of ginger. That said, in Europe and North America the current consensus for the maximum safe dose of ginger is 2 g/day in divided doses of 250 mg, even during pregnancy.³⁰

Methods

Study Aim

NVEP is a commonly encountered condition in family medicine, and ginger has been used as a nonpharmacological remedy. Our meta-analysis aimed to critically examine and synthesize available data from good-quality randomized clinical trials to evaluate the efficacy of ginger in treating NVEP.

Eligibility Criteria

Our primary interest was the treatment of NVEP using ginger as the therapeutic intervention. To minimize heterogeneity, we limited our scope to randomized, placebo-controlled trials with a satis-

factory score on the Cochrane Risk of Bias assessment tool.

Search Strategy

A literature search of published medical reports was performed in all languages using PubMed, EMBASE, CINAHL, the Cochrane Library, and all evidence-based medicine reviews using the OVID Portal of Queen's University, Kingston, Ontario. Abstracts were initially obtained using MeSH keywords of *early pregnancy*, *nausea*, *vomiting*, and *ginger*. Manual searches of references and review articles supplemented the computerized search.

Study Selection, Data Extraction, and Quality Assessment

Two reviewers (MT and RC) went through the initial abstracts. A simple form was adopted to select trials that satisfied the eligibility criteria stated above. Selected studies were evaluated according to the Cochrane Risk of Bias tool with regard to quality of the study, randomization protocol, adequacy of concealment and blinding, and rigor of follow-up for drop-outs. Information regarding the demography of the study population, duration of the study, the number of affected subjects who improved with treatment and placebo, and finally a numeric score for the Cochrane risk of bias was extracted and tabulated in spreadsheets. Of note, we found only 6 randomized, placebo-controlled trials of ginger that displayed data required for our meta-analysis (see Table 1). The primary outcome was improvement of pregnancy-related nausea and vomiting.

Statistical Analysis

All data from the 6 appropriate studies of ginger were synthesized in a meta-analysis, and odds ratios (ORs) were calculated with appropriate confidence intervals (CIs) based on the number of subjects reporting improvement in both the intervention and control groups. Where necessary, the value of 1 was added to any arm with zero outcome events according to the Sheele + 1 rule. To assess heterogeneity, we used the Cochrane Q-statistic with 95% confidence CIs. We assumed a *P* value of <.05 for the Cochran Q-statistic. Forest plots were generated with ORs for each ginger study. Statistical advice was provided by our data analyst at our Centre of Studies in Primary Care.

Results

We initially identified 135 records using the MeSH keywords of *early pregnancy*, *ginger*, *nausea*, and *vomiting*. After removing duplicates and further screening, 8 full articles were retrieved. Two of them were excluded because of irrelevance. Consensus was reached among the reviewers to include the final 6 studies. The 2009 PRISMA checklist was used; the PRISMA flow diagram is given in the Appendix.

In our meta-analysis, a total of 256 patients were randomly assigned to receive ginger, and 252 patients were randomly assigned to receive placebo. Total sample size per study ranged from 23 to 235. The period of intervention lasted from 4 days to 3 weeks. Dose and method of administration of ginger also varied among studies: Basirat et al³¹ used 5 biscuits per day, each containing 500 mg of ginger,

Table 1. Brief Description of the 6 Studies Included in the Meta-Analysis

Study	Intervention	Placebo	Duration of Study	Gestation Stage	Outcome Measure (Nausea and Vomiting)
Basirat et al. ³¹	500-mg ginger powder in biscuit, 5 biscuits/day	Placebo biscuit	4 days	<17 weeks	Nausea scale (VAS) + frequency of vomiting
Ozgoli et al. ³⁵	250-mg ginger capsules QID	Lactose capsules	4 days	<20 weeks	Nausea scale (VAS) + frequency of vomiting
Smith et al. ²⁶	350-mg ginger capsules TID	Vitamin B ₆ (25-mg capsules)	3 weeks	<16 weeks	Nausea and vomiting scales (Rhodes Index)
Keating et al. ³²	250-mg ginger syrup QID	Plain syrup	2 weeks	<12 weeks	4-point nausea scale + frequency of vomiting
Vutyavanich et al. ³⁴	250-mg ginger capsules QID	Placebo capsules	4 days	<17 weeks	VAS + Likert scale
Fischer-Rasmussen et al. ³³	250-mg ginger capsules QID	Lactose capsules	4 days	<20 weeks	Unique scoring system

QID, four times per day; TID, three times per day; VAS, visual analogue scale.

Table 2. Results from Meta-Analysis Comparing Ginger to Placebo in Improvement of Nausea and Vomiting in Early Pregnancy

Study	Intervention Group	Control Group	Odds Ratio	95% CI
Basirat et al. ³¹	28/32	21/30	3.00	0.81–11.1
Fischer-Rasmussen et al. ³³	19/27	4/27	13.66	3.56–52.43
Keating et al. ³²	10/13	2/10	13.33	1.78–100.14
Vutyavanich et al. ³⁴	28/32	10/35	17.50	4.87–62.87
Ozgili et al. ³⁵	27/32	20/35	4.05	1.26–12.99
Smith et al. ²⁶	68/120	69/115	0.87	0.52–1.47
Total (random effects)	180/256	126/252	4.89	1.88–12.73

CI, confidence interval.

whereas the others used either capsules or syrup containing approximately 1 g of ginger daily^{26,32–35} (Table 1).

We applied the Cochrane Risk of Bias assessment to the 6 studies; they achieved a score of at least 3 of 6 and were deemed to be of satisfactory quality.

Our primary outcome was the improvement of pregnancy-related nausea and vomiting, which was reported in all 6 studies: 180 of the 256 subjects in the ginger group and 126 of the 252 subjects in the placebo group reported improvement in symptoms of nausea and vomiting. In view of the interstudy variation in the duration and form of intervention, a random effects model was adopted. The pooled OR was 4.89, with a 95% CI of 1.88 to 12.73 (see Table 2). The Cochrane Q-statistic was significant at 33.72, with a degree of freedom of 5 ($P < .0001$) (Table 3). A synthesis Forest plot was generated (Figure 1), and the corresponding funnel plot is included for reference (Figure 2). Relative risk was calculated at 1.76 (95% CI, 1.18–2.65), and the number needed to treat for a positive effect was calculated to be 5.

Discussion

Our meta-analysis demonstrated that ginger (*Z. officinale*) is better than placebo in improving NVEP when administered at a dosage of approximately 1 g/day for a duration of at least 4 days.

Despite the common intake of ginger in our diet, no large-scale studies have directly assessed the safety profile of ginger during human preg-

nancy. In the early 1990s, Backon³⁶ commented on a theoretical possibility of ginger affecting testosterone receptor binding and sex steroid differentiation in the fetal brain. An antiplatelet effect of ginger through inhibition of thromboxane synthetase has been reported in vitro.³⁷ However, there have been no reports of increased risk of antenatal or postpartum hemorrhage with ginger use during pregnancy. Increased levels of early embryonic loss due to ginger intake³⁸ were reported in rats; however, the dosage was far greater than those routinely used in humans. Subsequent study of rats using human-compatible doses of ginger showed no similar adverse effects.³⁹


A commonly observed side effect of ginger is reflux. In a study by Willets et al,²⁴ 4 of the 120 subjects withdrew because of symptoms consistent with reflux. Reflux has been commonly reported with ginger use outside pregnancy. In a study of the prevention of postoperative nausea and vomiting, 8% of subjects had heartburn after taking 1 g ginger.⁴⁰ Another study of photopheresis-induced nausea demonstrated that 27% of subjects has reflux as a result of ginger use.⁴¹ Apart from gastric discomfort, the side effect of reflux poses no long-term harm or damage, and patients should be warned of this side effect before beginning ginger therapy.

There are a few limitations to our meta-analysis. First, there is variability in the dosage and formulation of ginger. While 4 studies administered ginger in capsules,^{26,33–35} one study administered ginger in syrup³² and the other in the form of biscuits.³¹ Moreover, the durations of intervention also differed. Four studies administered intervention for 4 days,^{31,33–35} while Keating and Chez³² gave ginger for 2 weeks and Smith et al²⁶ for 3

Table 3. Test for Heterogeneity of Studies

Q-statistic	33.72
Degree of freedom	5
Significance level	$P < .0001$


Figure 1. Forest plot for 6 studies using a random effects model. CI, confidence interval.



weeks. There was also variation in the combined sample size in the studies, which ranged from 23 to 235 (see Table 2). All these translated to a level of

heterogeneity that was reflected in the Cochran Q-statistic, with a significance of $P < .0001$. A final limitation of our study is the variability of scores

Figure 2. Funnel plot displaying the heterogeneity of studies. OR, odds ratio.



used to qualify and quantify outcome measures in NVEP. Vutyavanich et al,³⁴ Ozgoli et al,³⁵ and Basirat et al³¹ used a visual analog scale, whereas Fischer-Rasmussen et al³³ developed a unique scoring system. Keating and Chez used a symptoms diary and a 4-point scale and Smith et al used the Rhode's index. While this does add heterogeneity to our meta-analysis, the primary endpoint chosen remained the same (ie, improvement of NVEP).

Given its limitations, our meta-analysis demonstrated that ginger is better than placebo in improving NVEP, with an approximate number needed to treat of 5. Further large-scale, multicenter trials should be undertaken to further examine the efficacies of ginger and detail its safety profiles when treating NVEP.

Conclusions

Ginger is commonly consumed in our diet as an additive in cooking. In traditional Chinese medicine, ginger is indicated specifically as a remedy for NVEP. Based on our meta-analysis, we conclude that ginger is an effective nonpharmacological option for treating NVEP with respect to the inherent heterogeneity of the available studies. Family physicians and other medical professionals should be cognizant of the value of ginger as they contemplate pharmacological options for suitable patients with NVEP.

References

1. Kramer J, Bowen A, Stewart N, Muhajarine N. Nausea and vomiting of pregnancy: prevalence, severity and relation to psychosocial health. *MCN Am J Matern Child Nurs* 2013;38:21–7.
2. Lee NM, Saha S. Nausea and vomiting of pregnancy. *Gastroenterol Clin North Am* 2011;40:309–34, vii.
3. Basso O, Olsen J. Sex ratio and twinning in women with hyperemesis or pre-eclampsia. *Epidemiology* 2001;12:747–9.
4. Emelianova S, Mazzotta P, Einarson A, Koren G. Prevalence and severity of nausea and vomiting of pregnancy and effect of vitamin supplementation. *Clin Invest Med* 1999;22:106–10.
5. Gill SK, Maltepe C, Koren G. The effect of heartburn and acid reflux on the severity of nausea and vomiting of pregnancy. *Can J Gastroenterol* 2009; 23:270–2.
6. Hou JL, Wan XR, Xiang Y, Qi QW, Yang XY. Changes of clinical features in hydatidiform mole: analysis of 113 cases. *J Reprod Med* 2008;53: 629–33.

7. Zhang Y, Cantor RM, MacGibbon K, et al. Familial aggregation of hyperemesis gravidarum. *Am J Obstet Gynecol* 2011;204:230.e1–7.
8. Fejzo MS, Ingles SA, Wilson M, et al. High prevalence of severe nausea and vomiting of pregnancy and hyperemesis gravidarum among relatives of affected individuals. *Eur J Obstet Gynecol Reprod Biol* 2008;141:13–7.
9. Weigel MM, Weigel RM. The association of reproductive history, demographic factors, and alcohol and tobacco consumption with the risk of developing nausea and vomiting in early pregnancy. *Am J Epidemiol* 1988 ;127:562–70.
10. Buckwalter JG, Simpson SW. Psychological factors in the etiology and treatment of severe nausea and vomiting in pregnancy. *Am J Obstet Gynecol* 2002; 186(5 Suppl Understanding):S210–4.
11. Koren G, Clark S, Hankins GD, et al. Effectiveness of delayed-release doxylamine and pyridoxine for nausea and vomiting of pregnancy: a randomized placebo controlled trial. *Am J Obstet Gynecol* 2010; 203:571.e1–7.
12. Tucker ML, Jackson MR, Scales MD, Spurling NW, Tweats DJ, Capel-Edwards K. Ondansetron: pre-clinical safety evaluation. *Eur J Cancer Clin Oncol* 1989;25(Suppl 1):S79–93.
13. Einarson A, Maltepe C, Navioz Y, Kennedy D, Tan MP, Koren G. The safety of ondansetron for nausea and vomiting of pregnancy: a prospective comparative study. *BJOG* 2004;111:940–3.
14. Zhongjing Z. Synopsis of prescriptions of the golden chamber (Jingui Yaolue). Library of Chinese Classics. Middle Island, NY: New World Press; 2008.
15. National Tropical Botanical Garden. Zingiber officinale. Available from: http://ntbg.org/plants/plant_details.php?plantid=11651. Accessed July 1, 2013.
16. Govindarajan VS. Ginger–chemistry, technology, and quality evaluation: part 1. *Crit Rev Food Sci Nutr* 1982;17:1–96.
17. Govindarajan VS. Ginger–chemistry, technology, and quality evaluation: part 2. *Crit Rev Food Sci Nutr* 1982;17:189–258.
18. Grontved A, Brask T, Kambskard J, Hentzer E. Ginger root against seasickness. A controlled trial on the open sea. *Acta Otolaryngol* 1988;105:45–9.
19. Mowrey DB, Clayton DE. Motion sickness, ginger, and psychophysics. *Lancet* 1982;1:655–7.
20. Chaiyakunapruk N, Kitikannakorn N, Nathisuwan S, Leepakoboon K, Leelasettagool C. The efficacy of ginger for the prevention of postoperative nausea and vomiting: a meta-analysis. *Am J Obstet Gynecol* 2006;194:95–9.
21. Haniadka R, Rajeev AG, Palatty PL, Arora R, Baliga MS. Zingiber officinale (ginger) as an anti-emetic in cancer chemotherapy: a review. *J Altern Complement Med* 2012;18:440–4.
22. Marx WM, Teleni L, McCarthy AL, et al. Ginger (Zingiber officinale) and chemotherapy-induced

- nausea and vomiting: a systematic literature review. *Nutr Rev* 2013;71:245–54.
23. Niebyl JR, Goodwin TM. Overview of nausea and vomiting of pregnancy with an emphasis on vitamins and ginger. *Am J Obstet Gynecol* 2002;186(5 Suppl Understanding):S253–5.
 24. Willetts KE, Ekangaki A, Eden JA. Effect of a ginger extract on pregnancy-induced nausea: a randomised controlled trial. *Aust N Z J Obstet Gynaecol* 2003;43:139–44.
 25. Portnoi G, Chng LA, Karimi-Tabesh L, Koren G, Tan MP, Einarson A. Prospective comparative study of the safety and effectiveness of ginger for the treatment of nausea and vomiting in pregnancy. *Am J Obstet Gynecol* 2003;189:1374–7.
 26. Smith C, Crowther C, Willson K, Hotham N, McMillian V. A randomized controlled trial of ginger to treat nausea and vomiting in pregnancy. *Obstet Gynecol* 2004;103:639–45.
 27. Borrelli F, Capasso R, Aviello G, Pittler MH, Izzo AA. Effectiveness and safety of ginger in the treatment of pregnancy-induced nausea and vomiting. *Obstet Gynecol* 2005;105:849–56.
 28. Abdel-Aziz H, Windeck T, Ploch M, Verspohl EJ. Mode of action of gingerols and shogaols on 5-HT₃ receptors: binding studies, cation uptake by the receptor channel and contraction of isolated guinea-pig ileum. *Eur J Pharmacol* 2006;530:136–43.
 29. Pertz HH, Lehmann J, Roth-Ehrang R, Elz S. Effects of ginger constituents on the gastrointestinal tract: role of cholinergic M₃ and serotonergic 5-HT₃ and 5-HT₄ receptors. *Planta Med* 2011;77:973–8.
 30. Tiran D. Ginger to reduce nausea and vomiting during pregnancy: evidence of effectiveness is not the same as proof of safety. *Complement Ther Clin Pract* 2012;18:22–5.
 31. Basirat Z, Moghadamnia A, Kashifard M, Sarifirazavi A. The effect of ginger biscuit on nausea and vomiting in early pregnancy. *Acta Medica Iranica* 2009;47:51–6.
 32. Keating A, Chez RA. Ginger syrup as an antiemetic in early pregnancy. *Altern Ther Health Med* 2002;8:89–91.
 33. Fischer-Rasmussen W, Kjaer SK, Dahl C, Asping U. Ginger treatment of hyperemesis gravidarum. *Eur J Obstet Gynecol Reprod Biol* 1991;38:19–24.
 34. Vutyavanich T, Kraissarin T, Ruangsri R. Ginger for nausea and vomiting in pregnancy: randomized, double-masked, placebo-controlled trial. *Obstet Gynecol* 2001;97:577–82.
 35. Ozgoli G, Goli M, Simbar M. Effects of ginger capsules on pregnancy, nausea, and vomiting. *J Altern Complement Med* 2009;15:243–6.
 36. Backon J. Ginger in preventing nausea and vomiting of pregnancy; a caveat due to its thromboxane synthetase activity and effect on testosterone binding. *Eur J Obstet Gynecol Reprod Biol* 1991;42:163–4.
 37. Srivastava KC. Effect of onion and ginger consumption on platelet thromboxane production in humans. *Prostaglandins Leukot Essent Fatty Acids* 1989;35:183–5.
 38. Wilkinson JM. Effect of ginger tea on the fetal development of Sprague-Dawley rats. *Reprod Toxicol* 2000;14:507–12.
 39. Weidner MS, Sigwart K. Investigation of the teratogenic potential of a zingiber officinale extract in the rat. *Reprod Toxicol* 2001;15:75–80.
 40. Arfeen Z, Owen H, Plummer JL, Ilsley AH, Sorby-Adams RA, Doecke CJ. A double-blind randomized controlled trial of ginger for the prevention of post-operative nausea and vomiting. *Anaesth Intensive Care* 1995;23:449–52.
 41. Meyer K, Schwartz J, Crater D, Keyes B. Zingiber officinale (ginger) used to prevent 8-Mop associated nausea. *Dermatol Nurs* 1995;7:242–4.

Appendix: PRISMA Flow Diagram

