

7. Lumbiganon P, Laopaiboon M, Gülmezoglu AM, et al; World Health Organization Global Survey on Maternal and Perinatal Health Research Group. Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007–08. *Lancet* 2010;375:490–9.

doi: 10.3122/jabfm.2015.01.140269

The above letter was referred to the author of the article in question, who offers the following reply.

Response: Re: Promotion of Family-Centered Birth with Gentle Cesarean Delivery

To the Editor: We thank Dr. Pust for his comments and for reiterating the goal of promoting vaginal birth as the ideal mode of delivery whenever possible.

The context of our gentle cesarean work is a small urban hospital caring for a largely underserved population, including many high-risk obstetric patients. All well babies are placed skin-to-skin after birth regardless of the mode of delivery. Rooming-in is the norm—we have renamed the nursery the “procedure and monitoring room” and use it as such. In 2013, 458 deliveries occurred at our hospital; 29% of these were cesarean births, including 21% primary cesareans. Vaginal birth after cesarean was successfully achieved in 21 of 24 patients undergoing trial of labor after cesarean. Despite our largely underserved, often high-risk patient population, we are pleased that our primary cesarean rates remain the same or lower than national averages.¹

We note, however, that these statistics are not a perfect assessment of labor and delivery practices at our facility. In recent years a number of women desiring skin-to-skin contact after a planned cesarean birth have elected to transfer care to our facility toward the end of their pregnancy to participate in our gentle cesarean program, so that they may keep their baby with them in the operating room, assuming no emergencies arise. These women are included in the total and primary cesarean rate listed above.

We agree that cesarean delivery is a major surgery and should not be recommended as the mainstay. Yet when operative birth is needed, and mother and baby are healthy, we urge providers and hospital administrators to consider the benefits of the family-centered approach to cesarean delivery that we detailed in our article.² We believe early skin-to-skin contact between mother and child is possible for most cesarean deliveries and is a potentially important step in the bonding process, in achieving a successful early latch, and in helping to promote sustained breastfeeding.

Sincerely,

Susanna Magee, MD, MPH
Cynthia Battle, PhD
John Morton, MD
Melissa Nothnagle, MD, MSc
Brown University
Alpert Medical School
Providence, RI

References

- Osterman MJK, Martin JA. Primary cesarean delivery rates, by state: results from the revised birth certificate, 2006–2012. *Natl Vital Stat Rep* 2014;63:1–11.
- Magee SR, Battle C, Morton J, Nothnagle M. Promotion of family-centered birth with gentle cesarean delivery. *J Am Board Fam Med* 2014;27:690–3.

doi: 10.3122/jabfm.2015.01.140315

Re: Family Physicians' Knowledge of Commonly Overused Treatments and Tests

To the Editor: Maurer et al¹ outlined the results of a simple yet elegant study that shows family physicians' knowledge of commonly overused treatments and tests. At first glance, the article seems to demonstrate important gaps in knowledge; however, a detailed reading of the article gives cause for doubt.

Maurer et al¹ and the *Journal of the American Board of Family Medicine* should be lauded for publishing the actual multiple choice questions used in the study in the appendix of the article. Unfortunately, all the questions leave room for improvement. The second question offers the learner a binary choice: yes or no. Someone with no medical knowledge would have a 50% chance of getting this question correct. The test-wise candidate—who knows that he or she is undergoing an assessment to test his or her ability to “choose wisely”—might also feel prompted to give the answer that the examiner expects, and might thus choose the option with the least number of treatments or tests. The remaining questions ask the candidate to choose the most appropriate scenario for the question. Once again, this is less than optimal as it doesn't simulate clinical reality. Questions should ideally mimic clinical practice—hardly ever in clinical primary care medicine do 4 patients in a row present in relation to a possible Papanicolaou test (as in question 3). The other questions contain similar issues. These problems might seem minor; however, they may create questions that are not as valid and reliable as they should be. The art and science of constructing test questions has advanced over many years, and ignoring the advances that have been made seems a shortcoming.²

The good news is that the core content of the questions is likely to be sound and that judicious editing will make them valid and reliable. The questions could then be used in a variety of contexts outside the framework of the study. The questions, or questions like them, could be used efficiently and at low cost in formative or summative assessments, E-learning resources, or simply in promotional campaigns aiming to encourage wiser choices among physicians.³ Increasingly, the single best answer to such questions might be no investigation, no drug treatment, and no surgery.

Kieran Walsh
BMJ Learning
London, UK
kmwalsh@bmj.com

References

1. Maurer D, Stephens M, Reamy B, Crownover B, Crawford P, Chang T. Family physicians' knowledge of commonly overused treatments and tests. *J Am Board Fam Med* 2014; 27:699–703.
2. Case S, Swanson D. Constructing written test questions for the basic and clinical sciences. 3rd ed. Philadelphia: National Board of Medical Examiners; 2002.
3. Sandars J. Cost-effective e-learning in medical education. In: Walsh K, ed. Cost-effectiveness in medical education. Abingdon, UK: Radcliffe Publishing; 2010.

doi: 10.3122/jabfm.2015.01.140275

The above letter was referred to the author of the article in question, who offers the following reply.

Response: Re: Family Physicians' Knowledge of Commonly Overused Treatments and Tests

To the Editor: We appreciate Dr. Walsh's commentary and strongly agree that it is important to publish the actual questions used when survey-based research is published. While we respect his opinion that the questions could have been written differently, great care was taken

when drafting these questions so that the content and context could be understood by practicing clinicians. The questions were tested among several family physicians and were edited based on feedback gained from this process.

While we also appreciate that Dr. Walsh cites the excellent National Board of Medical Examiners guidance for constructing written test questions, a book we consult often when crafting test questions for our medical students, we offer as a caution the introductory guidance to this book from its authors: ". . .this manual will be useful primarily by faculty who are teaching medical students in basic science and clinical clerkships." This study's questions were crafted for an entirely different audience and we therefore chose to use a different approach.

Respectfully,

Douglas Maurer, DO, MPH
Mark Stephens, MD
Brian Reamy, MD
Brian Crownover, MD
Paul Crawford, MD
Tammy Chang, MD

doi: 10.3122/jabfm.2015.01.140287