

ORIGINAL RESEARCH

Reporting and Using Near-miss Events to Improve Patient Safety in Diverse Primary Care Practices: A Collaborative Approach to Learning from Our Mistakes

Steven Crane, MD, Philip D. Sloane, MD, Nancy Elder, MD, Lauren Cohen, MA, Natascha Laughtenschlaeger, MD, Kathleen Walsh, BA, and Sheryl Zimmerman, PhD

Purpose: Near-miss events represent an opportunity to identify and correct errors that jeopardize patient safety. This study was undertaken to assess the feasibility of a near-miss reporting system in primary care practices and to describe initial reports and practice responses to them.

Methods: We implemented a web-based, anonymous near-miss reporting system into 7 diverse practices, collecting and categorizing all reports. At the end of the study period, we interviewed practice leaders to determine how the near-miss reports were used for quality improvement (QI) in each practice.

Results: All 7 practices successfully implemented the system, reporting 632 near-miss events in 9 months and initiating 32 QI projects based on the reports. The most frequent events reported were breakdowns in office processes (47.3%); of these, filing errors were most common, with 38% of these errors judged by external coders to be high risk for an adverse event. Electronic medical records were the primary or secondary cause of the error in 7.8% and 14.4% of reported cases, respectively. The pattern of near-miss events across these diverse practices was similar.

Conclusions: Anonymous near-miss reporting can be successfully implemented in primary care practices. Near-miss events occur frequently in office practice, primarily involve administrative and communication problems, and can pose a serious threat to patient safety; they can, however, be used by practice leaders to implement QI changes. (J Am Board Fam Med 2015;28:452–460.)

Keywords: Medical Errors, Physician's Practice Patterns, Practice Management, Quality of Health Care

Near-miss events, or errors that are corrected before a patient is harmed, represent an opportunity to identify and correct flaws that jeopardize patient safety. Because more than half of all medical ambulatory visits occur in primary care, improved attention to near-miss events could markedly improve overall patient safety.¹ Others have demonstrated that error- and event-reporting systems can be implemented in primary care; however, these rarely focus on near-

misses or the coordination of near-miss reports with quality improvement (QI).^{2–9}

Barriers to reporting events include the additional workload burden, concern over punitive action, lack of confidence that positive change will result, and psychological barriers to admitting an error.^{10–14} Anonymous reporting systems may increase the number of error reports and reduce concerns about punitive actions but might

This article was externally peer reviewed.

Submitted 2 February 2014; revised 30 March 2015; accepted 14 April 2015.

From the Mountain Area Health Education Center, Asheville, NC (SC, NL, KW); the Cecil G. Sheps Center for Health Services Research (PDS, LC, SZ), and Department of Family Medicine and School of Medicine (PS), and School of Social Work (SZ), University of North Carolina—Chapel Hill, Chapel Hill and the University of Cincinnati, Cincinnati, OH (NE).

Funding: This study was funded by grant no. PS/R21 HS19558-01 from the US Agency for HealthCare Research and Quality (“Ambulatory Near Miss Reporting and Tracking to Improve Patient Safety,” Steven Crane, MD, Principal Investigator, September 2010–2011).

Conflict of interest: none declared.

Corresponding author: Steven D. Crane, Mountain Area Health Education Center, University of North Carolina—Chapel Hill, 121 Henderson Rd, Asheville, NC 28803 (E-mail: steven.crane@msj.org).

reduce the detail of the events.^{15,16} There is value in including all office staff in a reporting system, but this strategy may require frequent reminders to keep reporting volumes from dwindling.¹⁷

While errors occur frequently in primary care, few seem to result in significant harm to patients, consistent with the “near-miss” nature of many of these errors.^{18–22} Nevertheless, given the volume of ambulatory visits, even these relatively infrequent adverse events may be associated with a substantial portion of inpatient admissions and other patient harm.²³ A systematic approach to identify and correct near-miss events in primary care could be an important strategy to improve patient safety.

To demonstrate that such a system can be successfully adopted by a broad range of primary care practices, we designed and implemented an anonymous, practice-wide near-miss reporting and improvement tracking system in 7 diverse primary care medical practices. Our goals were to assess the feasibility of regular reporting, better understand the types of near-miss events that occur in ambulatory practices, and observe how medical practices use near-miss reports to initiate QI changes.

Methods

Participants

We recruited 7 diverse practices in western North Carolina to participate in this 1-year study. Practices included 2 family medicine residency practices, a federally qualified health center, a county-owned health department, and 3 private practices (2 family medicine, 1 pediatrics). Together the study practices employed more than 70 medical providers and 200 clinical support staff, provided >2000 office visits per month, and represented the full scope of primary care services (pediatric, geriatric, adult, and obstetric care) in both rural and urban settings. All but 1 used electronic medical records. Table 1 summarizes descriptive data on these practices.

Near-miss Reporting System

Our operational definition of a near-miss event was “an event/situation in which a negative outcome could have occurred but did not, either by chance or because the problem was identified and corrected before a negative outcome occurred.”²⁴ All staff members were invited to anonymously report near-miss events using an online form that had been adapted from previous studies and field tested,

Table 1. Description of the Study Practices

Type of practice	n (%)
Private	1 (14.3)
Part of a hospital system or other health system	3 (42.9)
Community health center	1 (14.3)
County health department	1 (14.3)
Residency program	1 (14.3)
Primary medical specialty represented	n (%)
Family medicine	6 (85.7)
Pediatrics	1 (14.3)
Number of Providers (full-time equivalents)	Mean (SD); Range
Physicians	3.9 (2.8); 1–8.75
Nurse practitioner or physician assistants	2.8 (2.3); 0.5–5.6
Physicians-in-training (residents)	5.2 (10.2); 0–27
Services provided	n (%)
Pediatric care	7 (100)
Obstetric care	3 (42.9)
Geriatric care	5 (71.4)
Number of Patient Encounters Per Year	Mean (SD); Range
Outpatient medical visits	22,589 (11,258.6); 10,000–36,000
Inpatient visits	3,433 (2,528.6); 0–5,688
Obstetric deliveries	230 (122.4); 0–315
Behavioral health visits	2,584 (1,835.0); 0–4,637
Nursing home visits	1,077 (748.5); 0–2,000
Home visits	41.6 (74.6); 0–185
Predominant medical record system for office visits	n (%)
Paper	1 (14.3)
Electronic	6 (85.7)
Percentage of Annual Patient Visits, by Age Category	Mean (SD); Range
< 18 years	52 (44.0); 10–100
18–64 years	47 (31.2); 0–75
≥ 65 years	16 (17.8); 0–50
Percentage of Annual Patient Visits, by Payer Status	Mean (SD); Range
Private insurance	27 (23.4); 0–70
HMO	3 (8.0); 0–22
Medicare	26 (23.2); 0–60
Medicaid	22 (14.5); 0–45
Self-pay	11 (15.1); 0–45
Charity	11 (23.1); 0–63
Percentage of Annual Visits by Patient Race/Ethnicity	Mean (SD); Range
White	88 (9.1); 75–95
African American	9 (6.2); 4–20
American Indian	0 (0.8); 0–2
Asian	1 (1.8); 0–5
Other	2 (3.3); 0–9
Hispanic or Latino Ethnicity	16 (18.8); 1–40

SD, standard deviation.

with an average completion time of 2 minutes per report (See Appendix).²⁵ The online form did not include any patient identifiers, was available electronically from any Internet-enabled computer, and stored reports on a central computer in an encrypted format. Staff attended a standardized, 1-hour orientation and during the study period received an automated E-mail message every 2 weeks inviting them to report any near-miss event they could recall from the previous 2 weeks. Project participation was phased in over 2 months from September 15 to November 30, 2010, and data collection was terminated at the end of June 2011; thus, the project period last 7 to 9 months, depending on practice site.

Near-miss Event Reports

Before being forwarded to the project's central computer, each near-miss report was reviewed by a designated individual in the practice (usually the medical director), who (1) excluded from the study any events that were adverse events causing patient harm; (2) ensured the absence of patient-identifying data in the responses forwarded for analysis; and (3) reviewed the incident for possible initiation of QI efforts in the practice. There was no attempt to standardize across practices which near-miss reports would be assigned for QI; during the structured interview process after the study, medical directors and practice administrators reported that they concentrated on events that seemed to be likely to recur, would have potential serious consequences if harm reached the patient, and seemed to be in their control to change.

QI from Event Reports

Initiation of QI around near misses was encouraged as part of the project. At the time of enrollment in the study, the 7 practices had significant differences in how they approached performance improvement. Several had robust QI teams in place, whereas others reported no formal performance improvement processes; none had incorporated near-miss reporting into the QI process. As part of study orientation, practice leaders each received a short orientation including a brief overview of how to initiate a Plan Do Study Act (PDSA) cycle and how to use the PDSA tracking software that was included in the near-miss system. After 3 months of successful reporting, each practice was expected to initiate at least 1 improvement process based on the

near-miss reports from the practice. All near-miss reports within each practice were reviewed every 2 months during a QI committee meeting. At the end of the project period, leaders from each practice participated in a structured group interview to gather additional information about how they actually responded to the information contained in the near-miss reports.

Each practice was reimbursed \$5,000 for identifying a core implementation team, participating in planning meetings and the all-staff orientation, and completing the baseline survey information. An additional \$1,500 per month was given to each practice when they reported at least 10 near-miss events and identified at least 1 near-miss event to remediate and track. Staff themselves did not receive any direct monetary inducement to submit reports, but several practices introduced small team-based rewards if the practice overall met the monthly reporting target. During the structured group interviews after the reporting period, there were no reports that staff felt pressured to report.

Data Analysis

After standardized training, the narrative portions of the near-miss error reports were coded by a team of 6 physician coders using a published taxonomy of ambulatory care errors.²⁶ For each report, the primary error was defined as "the breakdown in process, or knowledge/skill deficit that led to the reported problem." In addition, up to 4 associated or "cascade" errors and up to 4 contributing factors and possible preventive measures also were coded using the same taxonomy. The coders also provided their own subjective ratings, on a scale of 0 to 100, of the potential seriousness of the near-miss event, where 0 indicated "not very serious" and 100 indicated "extremely serious." They also rated the likelihood of harm and potential cost to the patient had the error actually occurred, as well as the estimated cost to the practice to remedy the system problem identified in the near-miss report, all on a 3-point scale, where 0 = "none/minimal," 1 = "some," and 2 = "a lot." Before coding, study leadership and coders met to achieve a common understanding about what would classify as "very serious" or "a lot of harm or cost." To ensure reliability of the coding and rating, 10% of the reports were coded independently by a second coder without knowledge of the first coder's results. Coder agreement was 70%

at the finest level of detail (3 levels in the 5-level taxonomy) and 87% at 2 levels of detail.

Quantitative data were analyzed using SAS 9.1 software (SAS, Inc., Cary, NC). Continuous data are reported using means and standard deviations (SDs), whereas categorical data are reported as frequencies and percentages. The study protocol was reviewed by and received institutional review board approval from Margaret R. Pardee Hospital.

Results

A total of 632 near misses were reported by the 7 practices. The most common categories of reported near-miss events, overall and by practice, are summarized in Table 2. The most common types of errors were breakdowns in office processes (47.3%), such as filing (25.3%), chart data entry errors (15.0%), problems with patient flow (2.2%), and problems with appointments and referrals (4.8%). The second most common category of errors was in ordering (6.2%), implementing (7.1%), or reporting the results of (12.2%) investigations, representing 25.5% of all near-miss reports. The pattern of near-miss events was similar across practices. Errors involving clinical knowledge or performance represented a very small percentage of errors (1.9%).

Table 3 reports coder ratings of near-miss severity, likelihood of an adverse event (AE) if the near miss had not been identified, the potential financial costs if the near-miss event had resulted in an AE, and the estimated cost to the practice to remedy the problem. Filing errors, the most common single near miss reported, had a mean severity rating of 51.8 (SD, 30.7), with 23.8% (n = 38) of these errors judged to be at high risk for leading to an AE had the error not been identified. Among all error types, those related to reporting investigations were rated as potentially most serious, with a mean severity score of 72.0 (SD, 28.3). Errors involving ordering medication and treatments (ordering, dispensing, or implementing) represented only 14% of near-miss reports but were rated as the second most severe (mean range, 59.1–63.0; SD range, 29.3–31.0).

Practices reported that 14.4% of the errors were secondarily attributable to the electronic medical record (EMR), including 21.9% of the filing errors, an example of which was an EMR interface that did not deliver the results of an important test to the ordering provider and resulted in a delay in addressing the test results. The EMR also was implicated in 40% of

ordering medication or treatment errors and 4.3% of communication with other health care providers sharing patient care errors. These computer-related errors had the third highest mean severity rating (mean, 59.2; SD, 25.2). The EMR was the perceived primary cause of 49 of these errors (7.8% of the total sample).

By the end of the study period, each of the practices had initiated at least 1 practice improvement process directly tied to the near-miss reports. Table 4 summarizes these practice improvement efforts.

Discussion

This study reports the results of the successful introduction of near-miss reporting in 7 primary care practices, each of which generated a substantial number and broad array of events and initiated performance improvement activities as a result.

The most frequent near-miss events recorded involved relatively mundane office processes such as charting data, filing, and computer operation, which is consistent with previous reports.^{2,6} Somewhat surprisingly, however, our data showed that administrative errors were frequently considered to carry with them the potential to lead to significant patient AEs, which supports our approach of encouraging all office staff to be involved in near-miss reporting. The events judged to be associated with the highest potential cost were those involving dispensing medication or implementing treatment (30% judged to involve “a lot” of potential cost) and handling test results (22% judged as “a lot”).

EMRs were directly linked to 14% of near-miss events, including 40% of the errors related to prescribing. Among the filing, data retrieval, and prescribing errors, 21.9%, 28.4%, and 40% of near-miss events, respectively, were attributed to EMR use. This finding reflects what others have found: While EMRs can reduce errors, they can also cause errors.^{27–29} Additional study of this important finding is needed to redesign EMRs to reduce error rates.

Participating practices seemed to have used the data generated from these near-miss reports to implement meaningful practice changes and improvements. Each initiated at least 1 Continuous Quality Improvement (CQI) project as a result of the study, and each identified at least 1 important safety improvement they made as a result of a near-miss report. Our interview data suggest that practice leaders found that immediate action or rapid PDSA

Table 2. Frequency of Near Miss Events Overall and by Participating Primary Care Practice

Near Miss Event	Number of Near Miss Events (% of Reports, by Site)							
	All Practices (n=632)	Practice A (n=43)	Practice B (n=177)	Practice C (n=80)	Practice D (n=139)	Practice E (n=69)	Practice F (n=43)	Practice G (n=81)
Office Process Problem								
Filing ^a	160 (25.3)	7 (16.3)	72 (40.7)	25 (31.3)	18 (13.0)	7 (10.1)	15 (34.9)	16 (19.8)
Chart data ^a	95 (15.0)	6 (14.0)	20 (11.3)	12 (15.0)	14 (10.1)	10 (14.5)	4 (9.3)	29 (35.8)
Patient flow	14 (2.2)	1 (2.3)	2 (1.1)	3 (3.8)	5 (3.6)	1 (1.5)	1 (2.3)	1 (1.2)
Appointment or referral	30 (4.8)	1 (2.3)	4 (2.3)	3 (3.8)	14	7 (10.1)	1 (2.3)	—
Equipment or Building Problem								
Equipment and physical building/surroundings/practice site	8 (1.3)	—	2 (1.1)	1 (1.3)	3 (2.2)	2 (2.9)	—	—
Other specific problems with computer	8 (1.3)	—	1 (0.6)	—	4 (2.9)	3 (4.4)	—	—
Investigations								
Ordering investigations	39 (6.2)	8 (18.6)	6 (3.4)	5 (6.3)	4 (2.9)	8 (11.6)	5 (11.6)	3 (3.7)
Implementing investigations	45 (7.1)	2 (4.7)	9 (5.1)	14 (17.5)	13 (9.4)	4 (5.8)	—	3 (3.7)
Reporting investigations	77 (12.2)	1 (2.3)	16 (9.0)	4 (5.0)	12 (8.6)	8 (11.6)	12 (27.9)	24 (29.6)
Medications or Other Treatments								
Ordering medications or treatments ^a	55 (8.7)	8 (18.6)	22 (12.4)	—	18 (13.0)	2 (2.9)	3 (7.0)	2 (2.5)
Dispensing medications or implementing treatments	36 (5.7)	4 (9.3)	14 (7.9)	1 (1.3)	11	9 (8.7)	—	—
Communication								
Communication with patients	30 (4.8)	1 (2.3)	6 (3.4)	8 (10.0)	8 (5.8)	5 (7.3)	2 (4.7)	—
Communication with other healthcare providers sharing patient care ^a	23 (3.6)	3 (7.0)	2 (1.1)	2 (2.5)	10 (7.2)	3 (4.4)	—	3 (3.7)
Clinical Knowledge or Performance								
Failure to follow standard or recommended practice	12 (1.9)	1 (2.3)	1 (0.6)	2 (2.5)	5 (3.6)	3 (4.4)	—	—

^a Across all practices, 21.9% (filing), 28.4% (chart data), 40.0% (ordering medications or treatments), and 4.3% (communication with other healthcare providers sharing patient care) of these near misses were secondarily attributable to an EMR-related problem.

Table 3. Perceived Severity and Estimated Cost of Selected Near Miss Events in Seven Primary Care Practices

Code	# of Reports	Event Description	Severity Rating ^a Mean (SD)	Likelihood of Adverse Event if Near Miss not Identified ^b n (%)	Potential Financial Cost of Event to Patient ^b n (%)	Estimated Financial Cost of Event to Practice ^b n (%)
Five Most Common Near Miss Events						
1.1.1	160	Filing problems	51.8 (30.7)	38 (23.8)	12 (7.5)	5 (3.1)
1.1.2	95	Chart data problems	35.4 (29.9)	11 (11.6)	8 (8.4)	4 (4.2)
1.3.2	45	Implementing investigations	52.2 (28.2)	10 (22.2)	4 (8.9)	3 (6.7)
1.3.3	77	Reporting investigations	72.0 (28.3)	38 (49.4)	17 (22.1)	1 (1.3)
1.4.1	55	Ordering medication or treatments	59.1 (29.3)	17 (30.9)	6 (10.9)	5 (9.1)
Five Near Miss Events Rated Most Potentially Severe						
1.2.1	8	Other specific problems with computer	59.2 (25.2)	5 (62.5)	1 (12.5)	0 (0)
1.3.3	77	Reporting investigations	72.0 (28.3)	38 (49.4)	17 (22.1)	1 (1.30)
1.4.1	55	Ordering medications or treatments	59.1 (29.3)	17(30.9)	6 (10.9)	5 (9.1)
1.4.2	36	Dispensing medications or implementing treatments	63.0 (31.0)	16 (44.4)	11 (30.6)	3 (8.3)
2.1	12	Failure to follow standard or recommended practice	56.5 (21.7)	2 (16.7)	3 (25.0)	1 (8.3)
Five ^c Near Miss Events Rated Most Potentially Costly to the Practice						
1.1.4	30	Appointment or referral	46.1 (28.6)	6 (20.0)	4 (13.3)	
1.2	8	Equipment and physical building/surroundings/practice site	49.7 (31.8)	2 (25.0)	1 (12.5)	
1.3.2	45	Implementing investigations	52.2 (28.2)	10 (22.2)	4 (8.9)	
1.4.1	55	Ordering medication or treatments	59.1 (29.3)	17 (30.9)	6 (10.9)	
1.4.2	36	Dispensing medications or implementing treatments	63.0 (31.0)	2 (44.4)	11 (30.6)	
2.1	12	Failure to follow standard or recommended practice	56.5 (21.7)	2 (16.7)	3 (25.0)	

^a On a scale of 0–100, where 0=not very serious and 100=extremely serious.

^b Rated categorically with 0 = none/minimal; 1 = some; 2 = a lot. Percent reflects ‘A lot’ responses within each error type.

^c Two events tied for fifth most costly.

SD, standard deviation.

Table 4. Summary of Practice Improvement Projects Implemented by Participating Primary Care Practices during the Study Period

	Primary Care Practice							
	A	B	C	D	E	F	G	
Length of study period in months	8	9	8	9	9	6	6	
Number of Practice Improvement Projects	Initiated	6	6	2	3	15	1	1
	Completed in study period	0	0	0	0	7	0	1
	Still in process at end of study period	6	0	1	1	4	1	0
	On hold at end of study period	0	1	1	1	0	0	0
	Inactivated during study period	0	5	0	1	4	0	0

cycles were used to avert potentially dangerous situations identified by near-miss reports; many expressed surprise about the type and frequency of near-miss errors that occurred in their practice. Indeed, the relatively large volume of near-miss reports generated by each practice suggests the importance of developing a systematic approach to process

improvement driven not only by potential for harm but also by frequency of occurrence.

Although our project included a cash bonus to practices for their participation, this did not seem to be an important issue for practices to continue near-miss reporting; the per capita reporting rate did not seem to vary depending on whether the

practice offered a reporting incentive. In fact, study practices have continued to log near-miss reports even after the project officially ended and the cash bonuses stopped. Practice leader buy-in and encouragement seems to be a key element of a successful reporting system, as has been demonstrated in hospital settings.³⁰

This study has several important limitations. Although we purposively chose practices to represent a diversity of size, ownership, specialty, and range of clinical services, our sample was small; therefore results cannot be generalized to all US primary care practices. Similarly, the frequency and types of near-miss reports in this sample cannot be used to estimate the frequency of actual near-miss events. Furthermore, even under the conditions of the study, some underreporting likely occurred. In addition, because event reporting was anonymous, we could not be certain that some events were not reported more than once (ie, by different individuals).

Our project involved only near-miss reports. We took great care to exclude AEs (where harm came to the patient) because of concerns of legal liability associated with data sharing. The self-report of likelihood of harm resulting from a near-miss event is, therefore, an estimate. According to leaders in our participating practices, near-miss events that affect patient outcomes are rare; therefore, it is possible either that the subjective estimates were exaggerated or, alternatively, that patients may suffer low-level AEs more often than they report. The reporting system did not invite patients to report errors, as some have suggested.³¹

The reporting phase of our project lasted only 7 to 9 months. The short time frame is insufficient to make broad conclusions about how practice change may result from near-miss reporting or how enduring those changes will prove to be. This important question requires further study over a longer period of time.

Conclusions

We demonstrated that an anonymous near-miss reporting system can be successfully implemented in a diverse group of primary care practices in a region. The reports generated indicate that near-miss events occur frequently in office practice, primarily involve administrative and communication problems, and occasionally pose a significant risk of

patient harm. Practice leaders in our project found these reports helpful and used this information to implement meaningful practice improvement. Further study is needed to determine whether these improvements can be sustained.

The near-miss reporting and tracking system was developed by Scott Pierson of WindSwept Solutions, Austin, Texas.

References

1. National Ambulatory Med Care Survey: 2010 summary tables. Available from: http://www.cdc.gov/nchs/data/ahcd/namcs_summary/2010_namcs_web_tables.pdf. Accessed May 19, 2015.
2. National Learning and Reporting Service, National Patient Safety Agency. Being open: communicating patient safety incidents with patients, their families and carers. Reference number 1097B. November 19, 2009. Available from: <http://www.nrls.npsa.nhs.uk/resources/?entryid45=65077>. Accessed May 19, 2015.
3. Hoffmann B, Beyer M, Rohe J, Gensichen J, Gerlach FM. "Every error counts": a web-based incident reporting and learning system for general practice. *Qual Saf Health Care* 2008;17:307–12.
4. Bhasale A. The wrong diagnosis: identifying causes of potentially adverse events in general practice using incident monitoring. *Fam Pract* 1998;15:308–18.
5. Fernald DH, Pace WD, Harris DM, West DR, Main DS, Westfall JM. Event reporting to a primary care patient safety reporting system: a report from the ASIPS collaborative. *Ann Fam Med* 2004;2:327–32.
6. Near-miss project e-newsletter. Albany: New York Chapter American College of Physicians. Available from: <http://www.nyacp.org/i4a/pages/index.cfm?pageid=3558>. Accessed May 19, 2015.
7. Hickner J, Zafar A, Kuo GM, et al. Field test results of a new ambulatory care Medication Error and Adverse Drug Event Reporting System—MEADERS. *Ann Fam Med* 2010;8:517–25.
8. Kaprielian V, Østbye T, Warburton S, Sangvai D, Michner L. A system to describe and reduce medical errors in primary care. In: Henriksen K, Battles JB, Keyes MA, Grady ML, eds. *Advances in patient safety: new directions and alternative approaches*. Vol. 1: Assessment. AHRQ publication no. 08–0034-1. Rockville, MD: Agency for Healthcare Research and Quality; 2008.
9. Zwart DL, Steerneman AH, van Rensen EL, Kalkman CJ, Verheij TJ. Feasibility of centre-based incident reporting in primary healthcare: the SPIEGEL study. *BMJ Qual Saf* 2011;20:121–7.
10. Elder NC, Graham D, Brandt E, et al. Barriers and motivators for making error reports from family medicine offices: a report from the American Academy of Family Physicians National Research Network (AAFP NRN). *J Am Board Fam Med* 2007;20:115–23.

11. Rowin E, Lucier D, Pauker S, Kumar S, Chen J, Salem D. Does error and adverse event reporting by physicians and nurses differ? *Jt Comm J Qual Patient Saf.* 2008;34:537–45.
12. Fisher MA, Mazor KM, Baril J, et al. Learning from mistakes. Factors that influence how students and residents learn from medical errors. *J Gen Intern Med* 2006;21:419–23.
13. Chan KD, Gallagher TH, Reznick, et al. How surgeons disclose medical errors to patients: a study using standardized patients. *Surgery* 2005;138:851–8.
14. Hingorani M, Wong T, Vafidis G. Patients' and doctors attitudes to amount of information given after unintended injury during treatment: cross sectional, questionnaire survey. *BMJ* 1999;318:640–1.
15. Makeham MA, Stromer S, Kidd MR, Lessons from the TAPS study-reducing the risk of patient harm. *Aust Fam Physician* 2008;37:339–40.
16. Pace WD, Staton EW, Higgins GS, Main DS, West DR, Harris DM; ASIPS Collaborative. Database design to ensure anonymous study of medical errors: a report from the ASIPS Collaborative. *J Am Med Inform Assoc* 2003;10:531–40.
17. Kennedy AG, Littenberg B, Senders JW. Using nurses and office staff to report prescribing errors in primary care. *Int J Qual Health Care* 2008;20:238–45.
18. Sandars J, Esmail A. The frequency and nature of medical error in primary care: understanding the diversity across studies. *Fam Pract* 2003;20:231–6.
19. Rubin G, George A, Chinn DJ, Richardson C. Errors in general practice: development of an error classification and pilot study of a method for detecting errors. *Qual Saf Health Care* 2003;12:443–7.
20. Kennedy AG, Littenberg B, Senders JW. Using nurses and office staff to report prescribing errors in primary care. *Int J Qual Health Care* 2008;20:238–45.
21. West DR, Pace WD, Dickinson LM, et al. Relationship between patient harm and reported medical errors in primary care: a report from the ASIPS Collaborative. In: Henriksen K, Battles JB, Keyes MA, Grady ML (eds). *Advances in patient safety: new directions and alternative approaches*. Vol. 1: Assessment. Rockville, MD: Agency for Healthcare Research and Quality; 2008.
22. Elder NC, Vonder Meulen M, Cassedy A. The identification of medical errors by family physicians during outpatient visits. *Ann Fam Med* 2004;2:125–9.
23. Woods DM, Thomas EJ, Holl JL, Weiss KB, Brennan TA. Ambulatory care adverse events and preventable adverse events leading to a hospital admission. *Qual Saf Health Care* 2007;16:127–31.
24. Aspden P, Corrigan JM, Wolcott J, et al. *Patient safety: achieving a new standard for care*. Washington, DC: National Academies Press; 2004.
25. Elder NC, Vonder Meulen M, Cassedy A. The identification of medical errors by family physicians during outpatient visits. *Ann Fam Med* 2004;2:125–9.
26. Dovey SM, Meyers DS, Phillips RL Jr, et al. A preliminary taxonomy of medical errors in family practice. *Qual Saf Health Care* 2002;11:233–8.
27. Bates DW, Cohen M, Leape LL, Overhage JM, Shabot MM, Sheridan T. Reducing the frequency of errors in medicine using information technology. *J Am Med Inform Assoc* 2001;8:299–308.
28. Harrington L, Kennerly D, Johnson C. Safety issues related to the electronic medical record (EMR): synthesis of the literature from the last decade, 2000–2009. *J Healthc Manag* 2011;56:31–44.
29. Nanji KC, Rothschild JM, Salzberg C, et al. Errors associated with outpatient computerized prescribing system. *J Am Med Inform Assoc* 2011;18:767–73.
31. Simmons D, Mick J, Graves K, Martin S. 26,000 Close call reports: lessons from the University of Texas Close Call Reporting System. In: Henriksen K, Battles JB, Keyes MA, Grady ML (eds). *Advances in patient safety: new directions and alternative approaches*. Vol. 1: Assessment. Rockville, MD: Agency for Healthcare Research and Quality; 2008.
32. Phillips RL, Dovey SM, Graham D, Elder NC, Hickner JM. Learning from different lenses: reports of medical errors in primary care by clinicians, staff, and patients: a Project of the American Academy of Family Physicians National Research Network. *J Patient Saf* 2006;2:140–6.

