

EDITORS' NOTE

Outcomes of Health System Structures, Highly Pertinent Clinical Information, Idea Stimulators, Clinical Reviews, and Prediction Tools: *JABFM* Exemplified

Marjorie A. Bowman, MD, MPA, Anne Victoria Neale, PhD, MPH, and
Dean A. Seebusen, MD, MPH

This issue exemplifies the types of articles that *JABFM* publishes to advance family medicine. We have articles on the implications of health system organizational structures. Three of these are international articles at the level of the national health system (1 from China) and systematic local health interventions (1 from Canada and 1 from Netherlands). Inside the United States, where there are more family physicians, there is less obesity, and designation as a Patient Centered Medical Home is related to increased rates of colorectal cancer screening. Review articles on common clinical topics discuss treatments that are changing (acne in pregnancy) or lack consensus (distal radial fractures). We have articles on making life easier in the office, such as for predicting Vitamin D levels, osteoporosis, and pre-diabetes in normal weight adults. There are articles to raise awareness of the “newest” testing or treatments, that is, auditory brainstem implants. “Reminder” articles highlight known entities that need to be reinforced to prevent over-/underdiagnosis or treatment, for example, “cotton fever.” Another article discusses the increased risk for postoperative complications with sleep apnea. We also provide “thought” pieces, in this case about the terminology we are using to extend our concept of patient-centered medical homes. (J Am Board Fam Med 2016;29:171–172.)

Systems of care make a difference. In the United States, areas with more family physicians per population have lower rates of obesity after controlling for other factors.¹ Patient-centered medical homes (PCMHs) are associated with increased rates of colorectal cancer screening compared with another targeted intervention.² Articles from international sources also help us to understand how the structures of health systems contribute to health and illness. We highlight an article herein on a lack of primary care clinicians in the Chinese health care system³ and 2 articles on health care–based systems of care for the frail elderly. The first, from Canada,⁴ found that instituting a specific care system in long-term care facilities reduced trips to the emergency department by one third. The second, from the Netherlands, found that a specific team-based intervention for frail elderly in their homes did not

lead to improvements in overall function or mortality.⁵

For our thought piece this month, Howard et al⁶ note that many practices undergoing transformation are using different terminology than are associated nationally with the PCMH initiative. These differences suggest future directions or areas that are inadequately addressed through the classic PCMH model.

Highly pertinent clinical articles reflect on common clinical problems and lack of evidence or disagreements about best treatments. For distal radius closed fractures, there is clearly insufficient evidence on the “best” treatment. These fractures, without complicating features, can be handled without surgery,⁷ thus permitting treatment in the family medicine setting. We have a clinical review article about treating acne during pregnancy.⁸ Whether or not family physicians provide prenatal care, it is common to be asked to treat medical problems that occur during pregnancy. Given fam-

Conflict of interest: The authors are editors of the *JABFM*.

ily physicians' substantial experience treating acne, doing so for pregnant women would clearly be within the realm of usual care.

While all our *JABFM* articles are chosen to provide new information to advance the care of patients in family medicine, we also choose articles that could be considered as reminders—such as one about an entity that is not easily recognized, or not well known: reducing unnecessary testing, or the wrong treatment. As an example, “cotton fever”⁹ is an entity that is apparently well known among intravenous drug users (including the patient in this report) but not by many physicians. With this diagnosis, less treatment is better, yet treatment of the underlying drug abuse is paramount. A second reminder article shows us that preoperative diagnosis of preexisting sleep disorders could lead to a more appropriate level of postoperative monitoring and treatment.¹⁰

In the category of new information, did you know that deaf children who fail treatment with a cochlear implant now have a new option? A new technology called an “auditory brainstem implant” offers the potential for some hearing¹¹—yet another avenue to improve lives.

We have two articles that help family physicians decide when testing or treatment is necessary – by predicting vitamin D levels, and predicting osteoporosis. For vitamin D, all that is needed is body mass index and ethnicity; refer to the useful graph in the article by Weishaar et al.¹² For osteoporosis, consider the differences between the commonly available screening tools.¹³ Mainous et al¹⁴ report on lower grip strength, an easy test that predicts diabetes, even in normal-weight adults; however, most readers are probably unaware of this connection. In addition, prediabetes is underdiagnosed.¹⁵ Next time you shake a patient's hand, think about how strong they are!

References

1. Gaglioti A, Petterson S, Bazemore A, Phillips RJ. Access to primary care in US counties is associated with lower obesity rates. *J Am Board Fam Med* 2016;29:182–90.
2. Green BB, Anderson ML, Chubak J, et al. Colorectal cancer screening rates increased after exposure to the patient-centered medical home. *J Am Board Fam Med* 2016; 29:191–200.
3. Wu D, Lam TP. Underuse of primary care in China: the scale, causes, and solutions. *J Am Board Fam Med* 2016;29:240–7.
4. Marshall EG, Clarke B, Burge F, Varatharasan N, Archibald G, Andrew MK. Improving continuity of care reduces emergency department visits by long-term care residents. *J Am Board Fam Med* 2016;29:201–8.
5. Ruikes FGH, Zuidema SU, Akkermans RP, Assendelft WJJ, Schers HJ, Koopmans RTCM. Multi-component program to reduce functional decline in frail elderly people: a cluster controlled trial. *J Am Board Fam Med* 2016;29:209–17.
6. Howard J, Etz RS, Crocker JB, et al. Maximizing the patient-centered medical home (PCMH) by choosing words wisely. *J Am Board Fam Med* 2016;29: 248–53.
7. Bruce KK, Merenstein DJ, Narvaez MV, et al. Lack of agreement on distal radius fracture treatment. *J Am Board Fam Med* 2016;29:218–25.
8. Chien AL, Qi J, Rainer B, Sachs DL, Helfrich YR. Treatment of acne in pregnancy. *J Am Board Fam Med* 2016;29:254–62.
9. Zerr AM, Ku K, Kara A. Cotton fever: a condition self-diagnosed by IV drug users. *J Am Board Fam Med* 2016;29:276–9.
10. Wolfe RM, Pomerantz J, Miller DE, Weiss-Coleman R, Solomonides T. Obstructive sleep apnea: preoperative screening and postoperative care. *J Am Board Fam Med* 2016;29:263–75.
11. Shah P, Kozin ED, Kaplan AB, Lee DJ. Pediatric auditory brainstem implant surgery: a new option for auditory habilitation in congenital deafness? *J Am Board Fam Med* 2016;29:286–8.
12. Weishaar T, Rajan S, Keller B. Probability of vitamin D deficiency by body weight and race/ethnicity. *J Am Board Fam Med* 2016;29:226–32.
13. Pecina JL, Romanovsky L, Merry SP, Kennel KA, Thacher TD. Comparison of clinical risk tools for predicting osteoporosis in women ages 50–64. *J Am Board Fam Med* 2016;29:233–9.
14. Mainous AG, Tanner RJ, Anton SD, Jo A. Low grip strength and prediabetes in normal-weight adults. *J Am Board Fam Med* 2016;29:280–2.
15. Mainous AG, Tanner RJ, Baker R. Prediabetes diagnosis and treatment in primary care. *J Am Board Fam Med* 2016;29:283–5.