Correspondence

Re: Primary Care Patients' Willingness to Participate in Comprehensive Weight Loss Programs: From the WWAMI Region Practice and Research Network

To the Editor: The article by Cole et al¹ about primary care patients' willingness to participate in comprehensive weight loss programs is progressive in addressing a major health problem, yet some areas need improvement. One objective of this study was to determine patient characteristics associated with willingness to participate in these programs. The investigators failed to include 2 important factors in their survey: income and educational level. According to the Centers for Disease Control and Prevention, (1) among Mexican American and non-Hispanic black men, those with higher incomes are more likely to be obese than those with lower incomes, (2) women with higher incomes are less likely to be obese than women with lower incomes, and (3) women with college degrees are less likely to be obese than women with lower educational levels.²

Another issue is the single delivery method of the survey, that is, article format. Investigators may have lost a population of patients who may not be able to read or write well (eg, less educated people, older adults) as a result of the lack of assistance in reading the survey. They also may have lost those who are more technologically advanced. Also, clinical staff offered the surveys to patients, which may have made patients feel obliged to take the survey. Some patients may have felt that the quality of their clinical care would be affected by not participating in the clinic-offered survey.

Next, the investigators aimed to identify potential facilitators and barriers to participation in comprehensive weight loss programs, but identified only the facilitators. They asked patients to mark the top 3 of 8 listed potential factors, yet these factors were all positive and did not identify barriers to participation.

In the discussion, the investigators explained that the Patient Protection and Affordable Care Act requires insurance companies to provide coverage for obesity treatment. This study included participants who were overweight, obese, and extremely obese. Yet, the sample included participants who were considered "at risk" but not obese. Therefore, this group should be excluded from the analysis in order for the results to pertain to patients who qualify for obesity treatment.

While this article was an advancement in the understanding of primary care patients' willingness to participate in comprehensive weight loss programs, all associated factors in the health outcomes were not assessed. These factors are necessary to tailor parsimonious and

appropriate comprehensive weight loss programs for primary care patients.

Denny Fe Garcia Agana, MPH
Department of Epidemiology
University of Florida College of Public
Health and Health Professions
H. James Free Center for Primary Care
Education & Innovation
University of Florida College of Medicine
Department of Community Health and Family Medicine
University of Florida College of Medicine
Gainesville, FL
dfgagana@ufl.edu

To see this article online, please go to: <http://jabfm.org/content/30/2/264.full>.

References

1. Cole AM, Keppel GA, Andrilla HA, Cox CM, Baldwin LM; WWAMI (Washington, Wyoming, Alaska, Montana, and Idaho) Region Practice and Research Network (WPRN) Patient Preferences for Weight Loss in Primary Care Development Group; The WPRN Practice Champions. Primary care patients' willingness to participate in comprehensive weight loss programs: from the WWAMI Region Practice and Research Network. *J Am Board Fam Med* 2016;29:572–80.
2. Centers for Disease Control and Prevention. Adult obesity facts. Available from <https://www.cdc.gov/obesity/data/adult.html>. Updated September 1, 2016. Accessed October 27, 2016.

doi: 10.3122/jabfm.2017.02.160353

The above letter was referred to the author of the article in question, who offers the following reply.

Response: Re: Primary Care Patients' Willingness to Participate in Comprehensive Weight Loss Programs: From the WWAMI Region Practice and Research Network

To the Editor: We appreciate the thoughtful comments from Fe Garcia Agana with regard to our recent article.¹ Fe Garcia Agana notes that our instrument did not assess patient income or education level as variables associated with reported willingness to participate in comprehensive weight loss programs. We agree that these patient factors are associated with risk of obesity within racial and ethnic groups, and may be important in predicting reported willingness to participate in comprehensive weight loss programs. We also acknowledge that offering the questionnaire only in written format may have limited or favored participation for certain groups.

In our project, we used a card study methodology, an established method for collecting observational data in

practice-based research networks.² Card studies are, by definition, brief and limited in scope.² Our study was developed and conducted using participatory methods.³ Thus, the method of administration and selected list of variables were chosen by the participating primary care practice champions to maximize simplicity during administration and minimize impact on clinical workflow. Finally, we considered excluding overweight and nearly overweight adults from our response sample, but sensitivity analysis without their responses found no changes in our primary outcomes; thus we chose to leave them in the sample.

Allison Cole, MD, MPH
Department of Family Medicine
University of Washington, Seattle
WWAMI Region Practice and Research Network
Institute of Translational Health Sciences, Seattle, WA
acole2@u.washington.edu

To see this article online, please go to: <http://jabfm.org/content/30/2/264.full>.

References

1. Cole AM, Keppel GA, Andrilla HA, Cox CM, Baldwin LM. Primary care patients' willingness to participate in comprehensive weight loss programs: from the WWAMI Region Practice and Research Network. *J Am Board Fam Med* 2016;29:572–80.
2. Westfall JM, Zittleman L, Staton EW, et al. Card studies for observational research in practice. *Ann Fam Med* 2011;9:63–8.
3. Cole A, Keppel GA, Linares A, et al. Evaluating the development, implementation and dissemination of a multisite card study in the WWAMI Region Practice and Research Network. *Clin Transl Sci* 2015;8:764–9.

doi: 10.3122/jabfm.2017.02.170016

Re: Presenting Signs of Multiple Myeloma and the Effect of Diagnostic Delay on the Prognosis

To the Editor: The article by Goldschmidt et al¹ addresses an important issue of the impact of early diagnosis in the outcome of patients with multiple myeloma. The authors mention that some have advocated use of serum-free light-chain assay (SFLCA) for “screening.” SFLCA has been promoted for diagnosing, determining the prognosis, and monitoring of monoclonal gammopathies.² However, empirical evidence suggests a far more limited role for SFLCA. Serum protein electrophoresis and serum immunofixation electrophoresis are the gold standards for diagnosis;³ these two alone are sufficient to diagnose about 95% cases. Patients with light-chain gammopathy can be detected by urine protein electrophoresis and urine immunofixation electrophoresis. Among patients without monoclonal gammopathy, the κ -to- λ ratio is abnormal in >35%, and the false-positive rate is about 55% in patients with polyclonal hypergammaglobulinemia.⁴ In monoclonal gammopathy there is an overall 27% false-negative κ -to- λ ratio. The false-negative rate is up to 67% for patients with monoclonal gammopathy of undeter-

mined significance.⁵ SFLCA and κ -to- λ ratio have virtually no role in the diagnosis of monoclonal gammopathy, as an abnormal κ -to- λ ratio is not diagnostic of monoclonal gammopathy and a normal κ -to- λ ratio does not exclude monoclonal gammopathy.⁶

Gurmukh Singh, MD, PhD, MBA
Department of Pathology
Medical College of Georgia at Augusta University, Augusta
gurmukhsinghmdphd@yahoo.com

To see this article online, please go to: <http://jabfm.org/content/30/2/265.full>.

References

1. Goldschmidt N, Zamir L, Poperno A, Kahan NR, Paltiel O. Presenting signs of multiple myeloma and the effect of diagnostic delay on the prognosis. *J Am Board Fam Med* 2016;29:702–9.
2. Bradwell A. Serum free light chain analysis plus Hevlylite. 7th ed. Birmingham, UK: Binding Site Group Ltd; 2015.
3. Rajkumar SV, Dimopoulos MA, Palumbo A, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol* 2014;15:e538–48.
4. Singh G. Serum free light chain assay and κ/λ ratio performance in patients without monoclonal gammopathies: high false-positive rate. *Am J Clin Pathol* 2016;146:207–14.
5. Kyle RA, Buadi F, Rajkumar SV. Management of monoclonal gammopathy of undetermined significance (MGUS) and smoldering multiple myeloma (SMM). *Oncology (Williston Park)* 2011;25:578–86.
6. Singh G. Serum free light chain assay and κ/λ ratio: performance in patients with monoclonal gammopathy-high false negative rate for κ/λ ratio. *J Clin Med Res* 2017;9:46–57.

doi: 10.3122/jabfm.2017.02.160356

The above letter was referred to the author of the article in question, who offers the following reply.

Response: Re: Presenting Signs of Multiple Myeloma and the Effect of Diagnostic Delay on the Prognosis

To the Editor: We thank Dr. Gurmukh Singh for his response. We are not advocating screening for multiple myeloma using a serum-free light-chain assay, and we agree with Dr. Gurmukh Singh that no evidence exists for the efficacy of serum-free light-chain testing in asymptomatic individuals. However, we suggest that this might be a worthwhile diagnostic test for patients with unexplained back pain and other “red flag” signs or symptoms, in whom multiple myeloma is suspected.

Neta Goldschmidt, MD
Department of Hematology
Hadassah Ein Kerem
Jerusalem, Israel
neta@hadassah.org.il

doi: 10.3122/jabfm.2017.02.170017

To see this article online, please go to: <http://jabfm.org/content/30/2/265.full>.