

RESEARCH LETTER

The Effects of Telephone Visits and Rurality on Veterans Perceptions of Access to Primary Care

Michelle Lampman, PhD, Susan Stockdale, PhD, Peter J. Kaboli, MD, MS, Erin Jaske, MPH, G. Blake Wood, MS, Walter L. Clinton, PhD, and Greg Stewart, PhD

Introduction: The objectives of this study were to examine if self-reported access to primary care is associated with actual patient wait times and use of telephone visits, and to assess whether this relationship differs by rural residence.

Methods: This study used 2016 administrative data from 994 primary care clinics within the Veterans Health Administration. Multiple-linear regression was used to examine relationships between patient perceptions of access and average actual patient wait time, use of telephone visits, and rural residence. Average panel size, clinic type, and panel severity were included as model covariates with cross-product terms for actual wait time, telephone use, and rurality to test for interactions.

Results: This study found patient perceptions of access aggregated at the clinic level to be conditional on the relationship between use of telephone visits, actual patient wait times, and rural residence. As actual wait time for routine appointments increases, Veterans served by clinics with a higher percent of rural Veterans perceive telephone visits more positively.

Discussion: These findings contribute to our understanding of factors associated with patient perceptions of access by highlighting complex interrelationships between strategies intended to improve access to care and how they can have differing impacts on perceptions among those living in rural or urban locations. (J Am Board Fam Med 2019;32:749–751.)

Keywords: Appointments and Schedules, Rural Health, Self Report, United States Department of Veterans Affairs, Veterans Health

Rural residents have identified several barriers to accessing care including distance, access to transportation, and availability of services.¹ The use of

telephone visits is 1 strategy being promoted to enhance access to primary care that does not require in-person visits.^{2,3} A recent study from the Veterans Health Administration (VHA) found that the use of telephone communication is strongly associated with higher patient perceptions of access.⁴ To our knowledge, no studies have specifically examined rural patients' perceptions of access associated with telephone visits, especially for clinics with longer wait times for in-person visits. Our objective was to examine whether self-reported access to primary care is associated with actual clinic

the national evaluation team of PCAT. Funding for PCAT is provided by the VA Office of Primary Care. The views expressed are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs.

Conflict of interest: none declared.

Corresponding author: Michelle Lampman, PhD, Division of Health Care Policy and Research, Department of Health Sciences Research, Mayo Clinic, 200 First Street SW, Rochester, MN 55905 (E-mail: Lampman.Michelle@mayo.edu).

This article was externally peer reviewed.

Submitted 4 February 2019; revised 23 May 2019; accepted 24 May 2019.

From VISN 23 Patient Aligned Care Team Demonstration Lab, Iowa City VA Healthcare System, Iowa City, IA (ML, GS); Department of Health Sciences Research, Mayo Clinic, Rochester, MN (ML); Health Services Research and Development Center for the Study of Healthcare Innovation, Implementation, and Policy, VA Greater Los Angeles Healthcare System Sepulveda, CA (SS); Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, CA (SS); Center for Access Delivery Research & Evaluation (CADRE), Iowa City VA Healthcare System, Iowa City, IA (PJK, GS); Department of Internal Medicine, The University of Iowa Carver College of Medicine, Iowa City, IA (PJK); VA Puget Sound Healthcare System, Seattle, WA (EJ, GBW, WLC); Kaiser Permanente Washington Health Research Institute, Seattle, WA (WLC); University of Iowa-Tippie College of Business, Iowa City, IA (GS).

Funding: This work was undertaken as part of the Veterans Administration's Primary Care Analytics Team (PCAT), supporting and evaluating VA's transition to a patient-centered medical home. Data for this report were developed by

wait times and use of telephone visits, and to assess whether this relationship differs between rural and urban settings.

Methods

In an observational study, we analyzed 2016 VHA administrative data from 994 VHA primary care clinics in the United States, representing data from 160,447 Veterans. Multiple-linear regression with clinics as the unit of analysis was used to examine relationships between patient perceptions of access (ie, percent of Veterans assigned to a clinic who reported they “always” received an appointment for routine primary care as soon as they needed) and 3 variables of interest: clinic wait time, use of telephone visits, and rurality defined by the proportion of patients classified as rural using the Rural-Urban Commuting Area codes corresponding with their home address.⁵ Average panel size, clinic type, and panel severity were included as model covariates with cross-product terms for clinic wait time, use of telephone visits, and rurality to test for interactions. This work is part of a larger quality improvement effort at the VHA and is not considered research activity subject to institutional review board review or waiver.

Results

As shown in Table 1, on average 52% of the Veterans assigned to these clinics reported that they were “Always” able to schedule an appointment for a checkup or routine care with their provider as soon as they needed. Average clinic performance related to access indicated that veterans wait about 4 days for routine care, and telephone visits accounted for 28% of all primary care encounters. The mean proportion of rural patients among these clinics was 53%. Results from our regression analyses (Table 1) indicate a significant 3-way interaction between clinic wait time, use of telephone visits, and rurality, indicating that perceptions differed depending on proportion of rural Veterans and use of telephone visits ($\beta = 0.071$, $P < .05$). We illustrate this interaction in Figure 1. As clinic wait time for routine appointments increases, Veterans served by clinics with a higher percent of rural Veterans perceive greater use of telephone visits as improving their access to primary care, but Veterans served by clinics with a lower percent of

Table 1. Clinic Characteristics and Multivariable Analyses of Patient Perceptions of Access to Routine Care (n = 994)

Clinic Characteristics	Mean (SD)
Patient perception of access to routine care (%)	51.9 (13.7)
Use of telephone visits (% of visits completed by telephone)	27.8 (11.5)
Clinic wait time (average patient wait time in days)	3.9 (3.2)
Rurality (% of rural patients)	52.9 (37.2)
Multivariable analyses of patient perceptions of access to routine care*	B
Clinic rurality	0.046
Clinic wait time	-0.009
Use of telephone visits	0.118
Clinic wait time × rurality (2-way interaction)	-0.012
Rurality × use of telephone visits (2-way interaction)	-0.151
Clinic wait time × use of telephone visits (2-way interaction)	-0.041
Clinic wait time × rurality × use of telephone visits (3-way interaction)	0.071 [†]

*Model adjusted for clinic type (outpatient clinic vs. medical center), average panel size, and panel severity (average comorbidity score).

[†] $P < .05$.


SD, standard deviation.

rural Veterans perceive greater use of telephone visits as inhibiting their access to primary care.

Discussion

This study found patient perceptions of access aggregated at the clinic level to be conditional on the relationship between use of telephone visits, clinic wait times, and rural residence. As actual wait time for routine appointments increases, Veterans served by clinics with a higher percent of rural Veterans perceive telephone visits more positively. While not focused specifically on rural, another study conducted in the VHA found favorable perceptions of telephone visits for routine care not requiring a physical from both Veterans and primary care providers.³ Relatedly, McKinstry et al⁶ found that telephone consultations were often used by rural patients and staff to maintain more regular contact and to avoid having to travel. They also reported that, unlike urban patients, rural patients were more likely to use telephone consultations to determine whether an in-person visit was necessary.⁶ Although more research is needed, these

Figure 1. Comparison of Interaction between Clinic Wait Time and Use of Telephone Visits among Urban and Rural Clinics.



findings contribute to our understanding of factors associated with patient perceptions of access by highlighting complex interrelationships between strategies intended to improve access to care and how they can have differing impacts on perceptions among those living in rural or urban locations.

To see this article online, please go to: <http://jabfm.org/content/32/5/749.full>.

References

1. Douthit N, Kiv S, Dwolatzky T, Biswas S. Exposing some important barriers to health care access in the rural USA. *Public Health* 2015;129:611–20.
2. Fortney JC, Burgess JF Jr, Bosworth HB, Booth BM, Kaboli PJ. A re-conceptualization of access for 21st century healthcare. *J Gen Intern Med* 2011;26:639–47.
3. Sperber NR, King HA, Steinhauer K, Ammarell N, Danus S, Powers BJ. Scheduled telephone visits in the veterans health administration patient-centered medical home. *BMC Health Serv Res* 2014;14:145.
4. Brunner J, Chuang E, Washington DL, et al. Patient-rated access to needed care: patient-centered medical home principles intertwined. *Women's Health Issues* 2018;28:165–71.
5. Johnson V, Wong E, Lampman M, et al. Comparing patient-centered medical home implementation in urban and rural VHA clinics: Results from the Patient Aligned Care Team Initiative. *J Ambul Care Manage* 2018;41:47–57.
6. McKinstry B, Watson P, Pinnock H, Heaney D, Sheikh A. Telephone consulting in primary care: a triangulated qualitative study of patients and providers. *Br J Gen Pract* 2009;59:e209–e218.