

Subject Index to Volume 32, 2019

- Academic medical centers**, yoga mentions in the electronic health record, 32(6):790–800
- Access to health care**
housing insecurity and, 32(4):521–530
team-based breastfeeding support, 32(6):818–826
- Accountable care organizations**, serving deprived communities, savings rate and, 32(6):913–922
- Acquired immunodeficiency syndrome (AIDS)**
Canadian HIV care settings as PC-MHs, 32(2):158–167
text messages for HIV testing in primary care, 32(3):408–412
- Acupuncture therapy**, training, physician burnout and, 32(2):259–263
- Addiction medicine**
opioid management, systems-based improvements to, 32(5):715–723
opioid prescribing
intervention in chronic noncancer pain, 32(4):559–566
pain-depression dyad and, 32(4):614–618
regulation on, in an urban safety-net, 32(3):362–374
opioid use disorder, treatment initiation for, 32(5):724–731
- Adolescence**, identifying substance use using frequency questions, 32(4):550–558
- Advance care planning**
billing code use, 32(6):827–834
conversations, 32(1):108–114, 32(6):835–846
outreach, portal-based, 32(6):925–930
- Advance directives**
adopting, in primary care, 32(2):168–179
portal-based advance care planning outreach, 32(6):925–930
- Adverse decision making**, optimal prescribing in patients with dementia, 32(3):383–391
- Altruism**, parent participation in child health research, 32(5):685–694
- Alzheimer disease**, disease-modifying treatments for, 32(6):931–940
- Ambulatory care**, retail clinic use among Medicare beneficiaries, 32(4):531–538
- American Academy of Family Physicians (AAFP)**, values and policies of, 32(6):948–950
- American Board of Family Medicine (ABFM)**
certification examination, validating, 32(6):876–882
data collection method for tracking specialty, 32(1):89–95
election of new officers and board members, 32(4):643–645
role in continuing education for family physicians, 32(5):756–758
self-assessment modules, physician opinions about, 32(1):79–88
- Amyloid β -peptides**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Analgesics**, natural history of back pain in older adults, 32(6):781–789
- Antidepressants**, enhanced shared decision making for depression, 32(1):6–9
- Antiviral agents**
hepatitis C and the role of primary care, 32(3):428–430
rapid point of care test for influenza, 32(2):226–233
- Appointments and schedules**
alternative payment impact on, 32(4):539–549
veterans perceptions of access to primary care, 32(5):749–751
- Asthma**, Colorado asthma toolkit implementation, 32(1):37–49
- Atherosclerosis**, suboptimal statin use for secondary prevention, 32(6):807–817
- Atopic dermatitis**, childhood, in primary care, 32(2):191–200
- Attitudes**, prediabetes at an academic practice, 32(4):505–512
- Automobile driving**, safety, discussions about, 32(4):607–613
- Back pain**, natural history of, in older adults, 32(6):781–789
- Behavior therapy**, translating self-management support into primary care, 32(3):341–352
- Behavioral medicine**
depression treatment for Spanish and English speakers, 32(6):904–912
outpatient follow-up and hospital readmissions, 32(3):353–361
translating self-management support into primary care, 32(3):341–352
walk-in psychiatric model for integrated care, 32(4):481–489
yoga mentions in the electronic health record, 32(6):790–800
- Behavioral risk factors**, housing insecurity and access to care, 32(4):521–530
- Beverages**, sugar-sweetened, intake, rural populations, 32(4):601–606
- Biomarkers**
dietary inflammatory index and diabetes severity, 32(6):801–806
disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Blood donors**, blood donation rates in the United States 1999–2016, 32(5):746–748
- Blood glucose**, levels, awareness of prediabetes status and subsequent health behavior, 32(1):20–27
- Blood pressure**, measurement, sources of error in, 32(5):732–738
- Blood transfusion**, blood donation rates in the United States 1999–2016, 32(5):746–748
- Body composition**, patient BMI knowledge in a rural population, 32(3):413–417
- Body mass index**
dietary inflammatory index and diabetes severity, 32(6):801–806
knee pain and osteoarthritis among overweight women, 32(4):575–584
knowledge, patient, in a rural population, 32(3):413–417
lifestyle intervention programs for diabetes, 32(4):596–600
sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Body weight**, awareness of prediabetes status and subsequent health behavior, 32(1):20–27
- Boston, Massachusetts**, social and behavioral determinants of health screening, 32(3):297–306
- Breastfeeding**, team-based support, 32(6):818–826
- Burnout, professional**
acupuncture training and, 32(2):259–263
clinic capacity and social needs, 32(1):69–78
medication management, impact on providers' work-life, 32(4):462–473
- California**
outpatient follow-up and hospital readmissions, 32(3):353–361
practice characteristics and patient self-management support, 32(3):329–340
- Canada**
doctor-patient relationships and men's depression treatment, 32(1):13–19
eConsult service implementation, 32(2):146–157
HIV care settings as PCMHs, 32(2):158–167
palliative health care for the homeless, 32(6):858–867
- Cannabis**, identifying substance use using frequency questions, 32(4):550–558
- Cardiovascular disease**
quality improvement measures by type of practice, 32(3):398–407
suboptimal statin use for secondary prevention, 32(6):807–817
veterans' reporting of, 32(1):50–57
- Centers for Medicare and Medicaid Services (US)**, accountable care savings in deprived communities, 32(6):913–922
- Certification**
ABM's data collection method for tracking specialty, 32(1):89–95
physician opinions about ABFM self-assessment modules, 32(1):79–88

- validating the ABFM certification examination, 32(6):876–882
- Child health**
- high-risk medication without documented contraception, 32(4):474–480
 - parent participation in child health research, 32(5):685–694
 - patient-centered research priorities, 32(5):674–684
 - research, parent participation in, 32(5):685–694
- Chronic disease**
- Canadian HIV care settings as PC-MHs, 32(2):158–167
 - housing insecurity and access to care, 32(4):521–530
 - knee pain and osteoarthritis among overweight women, 32(4):575–584
 - lifestyle intervention programs for diabetes, 32(4):596–600
 - management, yoga mentions in the electronic health record, 32(6):790–800
 - opioid management in the safety-net, 32(3):375–382
 - practice characteristics and patient self-management support, 32(3):329–340
 - preexisting conditions and the Affordable Care Act, 32(6):883–889
 - retail clinic use among Medicare beneficiaries, 32(4):531–538
 - social risk stratification in primary care, 32(4):585–595
 - suboptimal statin use for secondary prevention, 32(6):807–817
 - training vulnerable patients to use a patient portal, 32(2):248–258
 - translating self-management support into primary care, 32(3):341–352
 - visit entropy associated with diabetic control outcome, 32(5):739–745
- Chronic pain**
- management, opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
 - opioid management in the safety-net, 32(3):375–382
 - systems-based improvements to opioid management, 32(5):715–723
- Cigarette smoking**, electronic cigarettes and smoking reduction, 32(4):567–574
- Clinical decision making**, depression treatment for Spanish and English speakers, 32(6):904–912
- Codeine-combination**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Cognitive dysfunction**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Cohort studies**
- discussions about driving safety, 32(4):607–613
 - patient BMI knowledge in a rural population, 32(3):413–417
 - prediabetes at an academic practice, 32(4):505–512
 - preexisting conditions and the Affordable Care Act, 32(6):883–889
 - socioeconomics and hospitalization of veterans, 32(6):890–903
- Colon cancer**, screening, messaging for, 32(3):307–317, 32(3):318–328
- Colonoscopy**, colorectal cancer screening in a FQHC, 32(2):180–190
- Colorado**
- asthma toolkit implementation, 32(1):37–49
 - developing patient and community advisory groups, 32(5):663–673
 - implementation of patient-team partnership, 32(4):490–504
 - patient-centered research priorities, 32(5):674–684
 - practice characteristics and patient self-management support, 32(3):329–340
 - process for setting research priorities, 32(5):655–662
 - social risk stratification in primary care, 32(4):585–595
- Colorectal cancer**, screening endoscopic services, 32(4):454–456 in a FQHC, 32(2):180–190 messaging for, 32(3):307–317, 32(3):318–328
- Commentary**
- Commentary Response for “LARC Provision by Family Physicians: Low But on the Rise,” 32(1):4–5
 - Community Pharmacists as Partners in Reducing Suicide Risk, 32(6):763–767
 - Declining Presence of Family Physicians in Hospital-Based Care: A Major Concern or Totally Makes Sense?, 32(6):768–770
 - Endoscopic Services in the United States: By Whom, for What, and Why?, 32(4):454–456
 - Is It Time to Prioritize Diabetes Prevention in Practice?, 32(4):457–459
 - New Allopathic Medical Schools and Family Physicians, 32(5):651–652
 - No Magic Pill: A Prescription for Enhanced Shared Decision-Making for Depression Treatment, 32(1):6–9
 - Pawn and King Play: A Stalemate on Rural Emergency Medicine Staffing, 32(3):292–294
 - Practice-Based Research Today: A Changing Primary Care Landscape Requires Changes in Practice-Based Research Network (PBRN) Research, 32(5):647–650
 - Response for “LARC Provision by Family Physicians: Low But on the Rise,” 32(1):4–5
 - Stuck in Graduate Medical Education Traffic? Teaching Health Centers Are Family Medicine’s High-Occupancy-Vehicle Lane, 32(2):130–133
- Communication**
- discussions about driving safety, 32(4):607–613
 - patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
- Communication barriers**, a Hispanic health story from Texas, 32(1):115–117
- Community-acquired infections**, pneumonia, ruling out, 32(2):234–247
- Community health centers**
- Latino-based colorectal cancer screening messaging, 32(3):307–317
 - messaging for colorectal cancer screening, 32(3):318–328
 - preexisting conditions and the Affordable Care Act, 32(6):883–889
 - social and behavioral determinants of health screening, 32(3):297–306
- Community health nursing**, team-based breastfeeding support, 32(6):818–826
- Community medicine**
- developing patient and community advisory groups, 32(5):663–673
 - high-risk medication without documented contraception, 32(4):474–480
 - process for setting research priorities, 32(5):655–662
 - role of pharmacists in reducing suicide risk, 32(6):763–767
 - social and behavioral determinants of health screening, 32(3):297–306
 - team-based breastfeeding support, 32(6):818–826
- Community networks**
- developing patient and community advisory groups, 32(5):663–673
 - process for setting research priorities, 32(5):655–662
- Comorbidity**
- heart failure with preserved ejection fraction, 32(3):424–427
 - optimal prescribing in patients with dementia, 32(3):383–391
 - socioeconomics and hospitalization of veterans, 32(6):890–903
- Comprehensive medication management (CMM)**, impact on providers’ work-life, 32(4):462–473
- Continuity of care**, retail clinic use among Medicare beneficiaries, 32(4):531–538
- Continuity of patient care**
- declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
 - role of pharmacists in reducing suicide risk, 32(6):763–767
 - team-based breastfeeding support, 32(6):818–826
 - visit entropy associated with diabetic control outcome, 32(5):739–745
- Contraception**
- broken Nexplanon rods, 32(2):269–271

- high-risk medication without documented contraception, 32(4):474–480
- LARC provision by family physicians, 32(1):10–12
- Contraceptive devices**, broken Nexplanon rods, 32(2):269–271
- Correspondence**
- better doctor-patient relationships are associated with men choosing more active depression treatment, 32(4):640–641
- the complex interpretation and management of zika virus test results, 32(2):280–281
- family medicine and obstetrics: let us stop pretending, 32(2):279–280
- how evolving United States payment models influence primary care and its impact on the quadruple aim: the need for health equity, 32(1):118–119
- if we do not ask, they will not tell: screening for urinary and fecal incontinence by primary care providers, 32(1):119–120
- impact of a rapid point of care test for influenza on guideline consistent care and antibiotic use, 32(4):641–642
- impact of medical scribes in primary care on productivity, face-to-face time, and patient comfort, 32(1):120
- physician burnout and higher clinic capacity to address patients' social needs, 32(5):752–753
- a reflective case study in family medicine advance care planning conversations, 32(4):639–640
- signs and symptoms that rule out community-acquired pneumonia in outpatient adults: a systematic review and meta-Analysis, 32(5):753
- timely outpatient follow-up is associated with fewer hospital readmissions among patients with behavioral health conditions, 32(5):754–755
- Counseling**
- on fitness and strengthening, 32(1):103–107
- high-risk medication without documented contraception, 32(4):474–480
- practice characteristics and patient self-management support, 32(3):329–340
- team-based breastfeeding support, 32(6):818–826
- translating self-management support into primary care, 32(3):341–352
- Cross-cultural comparison**, depression treatment for Spanish and English speakers, 32(6):904–912
- Cross-sectional analysis**, quality improvement measures by type of practice, 32(3):398–407
- Cross-sectional studies**
- accountable care savings in deprived communities, 32(6):913–922
- benign skin lesions and TADA, 32(1):96–102
- dietary inflammatory index and diabetes severity, 32(6):801–806
- discussions about driving safety, 32(4):607–613
- family physicians providing e-visits, 32(6):868–875
- implementation of patient-team partnership, 32(4):490–504
- patient BMI knowledge in a rural population, 32(3):413–417
- Cultural competency**, hair care barrier to physical activity, 32(6):944–947
- Current procedural terminology**, advance care planning billing code use, 32(6):827–834
- Data warehousing**, advance care planning billing code use, 32(6):827–834
- Decision making**
- AAFP and politics, 32(6):948–950
- adverse, optimal prescribing in patients with dementia, 32(3):383–391
- depression treatment for Spanish and English speakers, 32(6):904–912
- enhanced shared, for depression, 32(1):6–9
- Delivery of health care**
- Canadian HIV care settings as PC-MHs, 32(2):158–167
- emergency department high utilizers, 32(2):264–268
- family physicians providing e-visits, 32(6):868–875
- palliative health care for the homeless, 32(6):858–867
- quality improvement measures by type of practice, 32(3):398–407
- team-based breastfeeding support, 32(6):818–826
- Dementia**
- disease-modifying treatments for Alzheimer disease, 32(6):931–940
- patients with, optimal prescribing in, 32(3):383–391
- Demography**, emergency department high utilizers, 32(2):264–268
- Depersonalization**, acupuncture training and physician burnout, 32(2):259–263
- Depression**
- enhanced shared decision making for, 32(1):6–9
- opioid prescribing and the pain-depression dyad, 32(4):614–618
- treatment, men's, and doctor-patient relationships, 32(1):13–19
- treatment of, for Spanish and English speakers, 32(6):904–912
- Dermatofibroma**, benign skin lesions and TADA, 32(1):96–102
- Dermatology**, drug-induced hyperpigmentation, 32(4):628–638
- Dermoscopy**, benign skin lesions and TADA, 32(1):96–102
- Diabetes mellitus**
- basal insulin in primary care, 32(3):431–447
- low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- monitoring the metabolic effects of antipsychotics, 32(3):418–423
- practice characteristics and patient self-management support, 32(3):329–340
- prevention, prioritizing, 32(4):457–459
- screening and treatment during office visits, 32(2):209–217
- severity, dietary inflammatory index and, 32(6):801–806
- translating self-management support into primary care, 32(3):341–352
- Diagnostic imaging**, early imaging for back pain in older adults, 32(6):773–780
- Diagnostic tests**
- disease-modifying treatments for Alzheimer disease, 32(6):931–940
- rapid point of care test for influenza, 32(2):226–233
- Diet**, dietary inflammatory index and diabetes severity, 32(6):801–806
- Disease management**
- practice characteristics and patient self-management support, 32(3):329–340
- social risk stratification in primary care, 32(4):585–595
- translating self-management support into primary care, 32(3):341–352
- Disease progression**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Doctor-patient relations**
- advance care planning conversations, 32(6):835–846
- discussions about driving safety, 32(4):607–613
- men's depression treatment and, 32(1):13–19
- patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
- portal-based advance care planning outreach, 32(6):925–930
- Drug abuse**, psychotherapeutic medication misuse screening, 32(2):272–278
- Drug overdose**, opioid prescribing and the pain-depression dyad, 32(4):614–618
- Drug reaction**, optimal prescribing in patients with dementia, 32(3):383–391
- Early detection of cancer**
- colorectal cancer screening in a FQHC, 32(2):180–190
- Latino-based colorectal cancer screening messaging, 32(3):307–317
- messaging for colorectal cancer screening, 32(3):318–328
- Editorial Office News and Notes**
- Content Usage and the Most Frequently Read Articles of 2018, 32(3):288–291

- Peer Reviewers for the Journal of the American Board of Family Medicine in 2018, 32(2):126–129
- Editor's Notes**
- Conversations, Communication, and Counseling Are Key Family Medicine Tactics to Improve Patient Health, 32(1):1–3
- Family Medicine: Data Driven Practice with Emphasis on Underserved Patients, 32(3):285–287
- Implementing Practice Changes in Family Medicine to Enhance Care and Prevent Disease Progression, 32(4):451–453
- Multiple Research Methodologies Can Advance the Science of Family Medicine, 32(2):123–125
- New Research on Back Pain, Diet and Diabetes, Advanced Care Planning, and Other Issues Frequently Seen in Family Medicine, 32(6):759–762
- Education**
- new allopathic medical schools train fewer family physicians, 32(5):653–654
- teaching health centers, 32(2):130–133
- Educational measurement**, validating the ABFM certification examination, 32(6):876–882
- Electronic cigarettes**, smoking reduction and, 32(4):567–574
- Electronic health records**
- advance care planning billing code use, 32(6):827–834
- eConsult service implementation in a new health region, 32(2):146–157
- portal-based advance care planning outreach, 32(6):925–930
- preexisting conditions and the Affordable Care Act, 32(6):883–889
- quality of the after-visit summary (AVS), 32(1):65–68
- social risk stratification in primary care, 32(4):585–595
- training vulnerable patients to use a patient portal, 32(2):248–258
- yoga mentions in, 32(6):790–800
- Electronic nicotine delivery systems**, smoking reduction and, 32(4):567–574
- Emergency care**, rural, family physicians' contributions, 32(3):295–296
- Emergency departments**
- high utilizers, 32(2):264–268
- staffing, rural hospitals, 32(3):292–294
- Emergency medicine**, staffing, rural hospitals, 32(3):292–294
- End of life care**
- adopting advance directives in primary care, 32(2):168–179
- portal-based advance care planning outreach, 32(6):925–930
- Endoscopic services**, colorectal cancer screening, 32(4):454–456
- Endoscopy**, endoscopic care by family physicians, 32(4):460–461
- Energy intake**, sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Entropy**, visit entropy associated with diabetic control outcome, 32(5):739–745
- Ethical theory**, advance care planning conversations, 32(1):108–114
- Evidence-based medicine**
- early imaging for back pain in older adults, 32(6):773–780
- implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Examination blueprint**, validating the ABFM certification examination, 32(6):876–882
- Exercise**
- counseling on fitness and strengthening, 32(1):103–107
- hair care barrier to physical activity, 32(6):944–947
- Family medicine**, reflections, Hispanic health story from Texas, 32(1):115–117
- Family physicians**
- ABFM's data collection method for tracking specialty, 32(1):89–95
- acupuncture training and physician burnout, 32(2):259–263
- advance care planning conversations, 32(1):108–114
- allopathic medical schools and, 32(5):653–654
- benign skin lesions and TADA, 32(1):96–102
- burnout, clinic capacity, and social needs, 32(1):69–78
- conversations on Twitter about diversity in medicine, 32(1):28–36
- declining presence of, in hospital-based care, 32(6):768–770, 32(6):771–772
- doctor-patient relationships and men's depression treatment, 32(1):13–19
- in emergent and urgent care, 32(3):295–296
- endoscopic care by family physicians, 32(4):460–461
- LARC provision by, 32(1):10–12
- opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
- patient BMI knowledge in a rural population, 32(3):413–417
- physician opinions about ABFM self-assessment modules, 32(1):79–88
- policies of AAFP and, 32(6):948–950
- provider perspectives of patient experiences in imaging, 32(3):392–397
- providing e-visits, 32(6):868–875
- reporting specialty status in research, 32(6):941–943
- systems-based improvements to opioid management, 32(5):715–723
- text messages for HIV testing in primary care, 32(3):408–412
- trained in teaching health centers, 32(2):134–135
- validating the ABFM certification examination, 32(6):876–882
- Fetal hemoglobin**, low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- Firearms**, values and policies of AAFP, 32(6):948–950
- Focus groups**
- palliative health care for the homeless, 32(6):858–867
- patient-centered research priorities, 32(5):674–684
- social and behavioral determinants of health screening, 32(3):297–306
- systems-based improvements to opioid management, 32(5):715–723
- Follow-up studies**, natural history of back pain in older adults, 32(6):781–789
- General practitioners**
- natural history of back pain in older adults, 32(6):781–789
- reporting specialty status in research, 32(6):941–943
- validating the ABFM certification examination, 32(6):876–882
- Geographic health care financing**, accountable care savings in deprived communities, 32(6):913–922
- Geography**, endoscopic care by family physicians, 32(4):460–461
- Geriatrics**
- early imaging for back pain in older adults, 32(6):773–780
- optimal prescribing in patients with dementia, 32(3):383–391
- Glucose**, low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- Glycohemoglobin A**
- low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- translating self-management support into primary care, 32(3):341–352
- Graduate medical education**, teaching health centers, 32(2):130–133, 32(2):134–135
- Grounded theory**, opioid management in the safety-net, 32(3):375–382
- Habits**, hair care barrier to physical activity, 32(6):944–947
- Hair care**, as barrier to physical activity, 32(6):944–947
- Harm reduction**
- end of life services for the homeless, 32(6):847–857
- palliative health care for the homeless, 32(6):858–867
- Health behavior**
- and awareness of prediabetes status, 32(1):20–27
- hair care barrier to physical activity, 32(6):944–947
- sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Health care costs**, physician characteristics and low value care spending, 32(2):218–225

- Health care disparities**
end of life services for the homeless, 32(6):847–857
social and behavioral determinants of health screening, 32(3):297–306
team-based breastfeeding support, 32(6):818–826
treatment initiation for opioid use disorder, 32(5):724–731
- Health care reform**, preexisting conditions and the Affordable Care Act, 32(6):883–889
- Health care surveys**
advance care planning conversations, 32(6):835–846
diabetes screening and treatment during office visits, 32(2):209–217
opioid prescribing and the pain-depression dyad, 32(4):614–618
validating the ABFM certification examination, 32(6):876–882
- Health care systems**, alternative payment impact on visits and scheduling, 32(4):539–549
- Health care use**, outpatient follow-up and hospital readmissions, 32(3):353–361
- Health equity**
accountable care savings in deprived communities, 32(6):913–922
end of life services for the homeless, 32(6):847–857
- Health expenditures**, social risk stratification in primary care, 32(4):585–595
- Health information systems**, accountable care savings in deprived communities, 32(6):913–922
- Health insurance**
alternative payment impact on visits and scheduling, 32(4):539–549
values and policies of AAFP, 32(6):948–950
- Health literacy**
patient BMI knowledge in a rural population, 32(3):413–417
training vulnerable patients to use a patient portal, 32(2):248–258
- Health personnel**, LARC provision by family physicians, 32(1):10–12
- Health policy**
advance care planning billing code use, 32(6):827–834
alternative payment impact on visits and scheduling, 32(4):539–549
declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
endoscopic care by family physicians, 32(4):460–461
preexisting conditions and the Affordable Care Act, 32(6):883–889
- Health promotion**
team-based breastfeeding support, 32(6):818–826
translating self-management support into primary care, 32(3):341–352
- Health services**, alternative payment impact on visits and scheduling, 32(4):539–549
- Health services research**, laboratory safety toolkit in primary care, 32(2):136–145
- Heart failure**, with preserved ejection fraction, 32(3):424–427
- Hematology**, low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- Hemoglobinopathies** low hemoglobin A1c and, 32(6):923–924
- Hepatitis C**, expanding the role of primary care and, 32(3):428–430
- Hispanic Americans**
depression treatment for Spanish and English speakers, 32(6):904–912
a Hispanic health story from Texas, 32(1):115–117
Latino-based colorectal cancer screening messaging, 32(3):307–317
messaging for colorectal cancer screening, 32(3):318–328
- Homeless persons**
end of life services for, 32(6):847–857
housing insecurity and access to care, 32(4):521–530
palliative health care for, 32(6):858–867
- Hospital medicine**, declining presence of family physicians, 32(6):768–770, 32(6):771–772
- Hospitalists**, blood donation rates in the United States 1999–2016, 32(5):746–748
- Hospitalization**
outpatient follow-up and hospital readmissions, 32(3):353–361
patients frequently readmitted versus other admitted patients, 32(1):58–64
of veterans, socioeconomics, 32(6):890–903
- House calls**, advance care planning conversations, 32(6):835–846
- Housing**
insecurity, access to care and, 32(4):521–530
issues, socioeconomics and hospitalization of veterans, 32(6):890–903
palliative health care for the homeless, 32(6):858–867
- Hydrocodone**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Hydrocodone-combination**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Hypercholesterolemia**, suboptimal statin use for secondary prevention, 32(6):807–817
- Hyperglycemia**, basal insulin in primary care, 32(3):431–447
- Hyperlipidemia**, suboptimal statin use for secondary prevention, 32(6):807–817
- Hyperpigmentation**, drug-induced, 32(4):628–638
- Hypertension**, sources of error in blood pressure measurement, 32(5):732–738
- Hypoglycemia**, basal insulin in primary care, 32(3):431–447
- Hypoglycemic drugs**, diabetes screening and treatment during office visits, 32(2):209–217
- Immigrants**, Hispanic health story from Texas, 32(1):115–117
- Implementation**, laboratory safety toolkit in primary care, 32(2):136–145
- Incentive reimbursement**, advance care planning billing code use, 32(6):827–834
- Infant health**, high-risk medication without documented contraception, 32(4):474–480
- Inflammation**, dietary inflammatory index and diabetes severity, 32(6):801–806
- Influenza**, rapid point of care test for, 32(2):226–233
- Information technology**
practice characteristics and patient self-management support, 32(3):329–340
social risk stratification in primary care, 32(4):585–595
training vulnerable patients to use a patient portal, 32(2):248–258
- Inpatients**, declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
- Insulin**, basal, in primary care, 32(3):431–447
- Insurance**, accountable care savings in deprived communities, 32(6):913–922
- Insurance coverage**, preexisting conditions and the Affordable Care Act, 32(6):883–889
- Integrated care**, walk-in psychiatric model for, 32(4):481–489
- Integrative medicine**, acupuncture training and physician burnout, 32(2):259–263
- Interdisciplinary research**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Internet**, portal-based advance care planning outreach, 32(6):925–930
- Internship and residency**, reporting specialty status in research, 32(6):941–943
- Iowa**, advance care planning billing code use, 32(6):827–834
- Job satisfaction**, medication management improves providers' work-life, 32(4):462–473
- Journal impact factor**, reporting specialty status in research, 32(6):941–943
- Journal of the American Board of Family Medicine (JABFM)**
content usage and the most frequently read articles of 2018, 32(3):288–291
peer reviewers for, 32(2):126–129

- Korea**, electronic cigarettes and smoking reduction, 32(4):567–574
- Lactation**, team-based breastfeeding support, 32(6):818–826
- Latino immigrants**, Hispanic health story from Texas, 32(1):115–117
- Leadership**, declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
- Lifestyle**
awareness of prediabetes status and subsequent health behavior, 32(1):20–27
hair care barrier to physical activity, 32(6):944–947
practice characteristics and patient self-management support, 32(3):329–340
prediabetes at an academic practice, 32(4):505–512
translating self-management support into primary care, 32(3):341–352
yoga mentions in the electronic health record, 32(6):790–800
- Linear models**, housing insecurity and access to care, 32(4):521–530
- Linguistic issues**, Hispanic health story from Texas, 32(1):115–117
- Lisinopril**, high-risk medication without documented contraception, 32(4):474–480
- Living wills**, adopting advance directives in primary care, 32(2):168–179
- Logistic models**
depression treatment for Spanish and English speakers, 32(6):904–912
dietary inflammatory index and diabetes severity, 32(6):801–806
discussions about driving safety, 32(4):607–613
family physicians providing e-visits, 32(6):868–875
high-risk medication without documented contraception, 32(4):474–480
housing insecurity and access to care, 32(4):521–530
opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
suboptimal statin use for secondary prevention, 32(6):807–817
- Logistic regression**
monitoring the metabolic effects of antipsychotics, 32(3):418–423
outpatient follow-up and hospital readmissions, 32(3):353–361
- Long-acting reversible contraception (LARC)**, provision by family physicians, 32(1):10–12
- Low back pain**, early imaging for back pain in older adults, 32(6):773–780
- Low-income population**, text messages for HIV testing in primary care, 32(3):408–412
- Major depressive disorder**, enhanced shared decision making for depression, 32(1):6–9
- Meaningful use**, quality of the after-visit summary (AVS), 32(1):65–68
- Medicaid**
advance care planning billing code use, 32(6):827–834
alternative payment impact on visits and scheduling, 32(4):539–549
depression treatment for Spanish and English speakers, 32(6):904–912
emergency department high utilizers, 32(2):264–268
preexisting conditions and the Affordable Care Act, 32(6):883–889
walk-in psychiatric model for integrated care, 32(4):481–489
- Medical education**
declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
validating the ABFM certification examination, 32(6):876–882
- Medical geography** accountable care savings in deprived communities, 32(6):913–922
- Medical informatics**
social risk stratification in primary care, 32(4):585–595
yoga mentions in the electronic health record, 32(6):790–800
- Medical schools**, new allopathic medical schools train fewer family physicians, 32(5):653–654
- Medical societies**, conversations on Twitter about diversity in medicine, 32(1):28–36
- Medically uninsured**
alternative payment impact on visits and scheduling, 32(4):539–549
outpatient follow-up and hospital readmissions, 32(3):353–361
preexisting conditions and the Affordable Care Act, 32(6):883–889
- Medicare**
early imaging for back pain in older adults, 32(6):773–780
emergency department high utilizers, 32(2):264–268
physician characteristics and low value care spending, 32(2):218–225
retail clinic use among Medicare beneficiaries, 32(4):531–538
- Meditation**, yoga mentions in the electronic health record, 32(6):790–800
- MEDLINE**, reporting specialty status in research, 32(6):941–943
- Mental disorders**, monitoring the metabolic effects of antipsychotics, 32(3):418–423
- Mental health**
depression treatment for Spanish and English speakers, 32(6):904–912
housing insecurity and access to care, 32(4):521–530
monitoring the metabolic effects of antipsychotics, 32(3):418–423
patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
role of pharmacists in reducing suicide risk, 32(6):763–767
treatment initiation for opioid use disorder, 32(5):724–731
- Mental health services**, Canadian HIV care settings as PCMHs, 32(2):158–167
- Mental status and dementia tests**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Meta-analysis**, ruling out community-acquired pneumonia, 32(2):234–247
- Metabolic diseases**, monitoring the metabolic effects of antipsychotics, 32(3):418–423
- Metformin**, diabetes screening and treatment during office visits, 32(2):209–217
- Military personnel veterans'** reporting of cardiovascular disease, 32(1):50–57
- Minority health**
hair care barrier to physical activity, 32(6):944–947
opioid management in the safety-net, 32(3):375–382
walk-in psychiatric model for integrated care, 32(4):481–489
- Morphine milligram equivalency (MME)**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Motivation**
colorectal cancer screening in a FQHC, 32(2):180–190
parent participation in child health research, 32(5):685–694
- Multimorbidity**, advance care planning conversations, 32(6):835–846
- Multiple chronic conditions**, advance care planning conversations, 32(6):835–846
- Multivariate analysis**, housing insecurity and access to care, 32(4):521–530
- Narcotics**, opioid prescribing and the pain-depression dyad, 32(4):614–618
- Nebraska**, sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Needs assessment**
establishing a PBRN stakeholder engagement infrastructure, 32(5):695–704
team-based breastfeeding support, 32(6):818–826
- Netherlands**
knee pain and osteoarthritis among overweight women, 32(4):575–584
natural history of back pain in older adults, 32(6):781–789
- New Mexico**, implementation of patient-team partnership, 32(4):490–504

- Newborns**, team-based breastfeeding support, 32(6):818–826
- Nicotine chewing gum**, smoking reduction and, 32(4):567–574
- No-show patients**, alternative payment impact on visits and scheduling, 32(4):539–549
- Northwestern United States**, treatment initiation for opioid use disorder, 32(5):724–731
- Nurse practitioners**
 provider perspectives of patient experiences in imaging, 32(3):392–397
 retail clinic use among Medicare beneficiaries, 32(4):531–538
- Nutrition surveys**
 blood donation rates in the United States 1999–2016, 32(5):746–748
 dietary inflammatory index and diabetes severity, 32(6):801–806
- Obesity**
 hair care barrier to physical activity, 32(6):944–947
 knee pain and osteoarthritis among overweight women, 32(4):575–584
 lifestyle intervention programs for diabetes, 32(4):596–600
 patient BMI knowledge in a rural population, 32(3):413–417
 practice characteristics and patient self-management support, 32(3):329–340
 translating self-management support into primary care, 32(3):341–352
- Occult blood**
 colorectal cancer screening in a FQHC, 32(2):180–190
 Latino-based colorectal cancer screening messaging, 32(3):307–317
- Occupational burnout**, medication management improves providers' work-life, 32(4):462–473
- Oklahoma**, adopting advance directives in primary care, 32(2):168–179
- Ondansetron**, high-risk medication without documented contraception, 32(4):474–480
- Ontario**, end of life services for the homeless, 32(6):847–857
- Opioid prescription**
 intervention in chronic noncancer pain, 32(4):559–566
 pain-depression dyad and, 32(4):614–618
 regulation on, in an urban safety-net, 32(3):362–374
- Opioid-related disorders**
 opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
 treatment initiation for opioid use disorder, 32(5):724–731
- Opioids**
 early imaging for back pain in older adults, 32(6):773–780
 opioid management in the safety-net, 32(3):375–382
- systems-based improvements to opioid management, 32(5):715–723
- Oregon**
 alternative payment impact on visits and scheduling, 32(4):539–549
 social risk stratification in primary care, 32(4):585–595
- Organizational innovation**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Organizations**, establishing a PBRN stakeholder engagement infrastructure, 32(5):695–704
- Osteoarthritis**, knee pain and osteoarthritis among overweight women, 32(4):575–584
- Outcomes assessment**
 Colorado asthma toolkit implementation, 32(1):37–49
 housing insecurity and access to care, 32(4):521–530
 patient-centered research priorities, 32(5):674–684
- Outcomes research**, early imaging for back pain in older adults, 32(6):773–780
- Outpatients**
 emergency department high utilizers, 32(2):264–268
 follow-up and hospital readmissions, 32(3):353–361
 ruling out community-acquired pneumonia, 32(2):234–247
- Overweight**, lifestyle intervention programs for diabetes, 32(4):596–600
- Ownership**, quality improvement measures by type of practice, 32(3):398–407
- Pain**
 knee pain and osteoarthritis among overweight women, 32(4):575–584
 opioid prescribing and the pain-depression dyad, 32(4):614–618
- Pain management**, early imaging for back pain in older adults, 32(6):773–780
- Palliative care**
 advance care planning billing code use, 32(6):827–834
 advance care planning conversations, 32(6):835–846
 for the homeless, 32(6):858–867
- Parents**
 lifestyle intervention programs for diabetes, 32(4):596–600
 parent participation in child health research, 32(5):685–694
 patient-centered research priorities, 32(5):674–684
- Partnership practice**, implementation of patient-team partnership, 32(4):490–504
- Patient activation**, text messages for HIV testing in primary care, 32(3):408–412
- Patient care**
 patient-centered research priorities, 32(5):674–684
- patients frequently readmitted versus other admitted patients, 32(1):58–64
 physician characteristics and low value care spending, 32(2):218–225
- Patient care team**
 depression treatment for Spanish and English speakers, 32(6):904–912
 implementation of patient-team partnership, 32(4):490–504
 practice characteristics and patient self-management support, 32(3):329–340
- Patient-centered care**
 adopting advance directives in primary care, 32(2):168–179
 advance care planning conversations, 32(6):835–846
 burnout, clinic capacity, and social needs, 32(1):69–78
 Canadian HIV care settings as PC-MHs, 32(2):158–167
 hair care barrier to physical activity, 32(6):944–947
 implementation of patient-team partnership, 32(4):490–504
 patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
 practice characteristics and patient self-management support, 32(3):329–340
 research priorities, 32(5):674–684
 retail clinic use among Medicare beneficiaries, 32(4):531–538
 socioeconomic and hospitalization of veterans, 32(6):890–903
 translating self-management support into primary care, 32(3):341–352
 visit entropy associated with diabetic control outcome, 32(5):739–745
- Patient compliance**, depression treatment for Spanish and English speakers, 32(6):904–912
- Patient discharge**
 family medicine transitional care service line, 32(4):619–627
 patients frequently readmitted versus other admitted patients, 32(1):58–64
- Patient health questionnaire**, depression treatment for Spanish and English speakers, 32(6):904–912
- Patient outcome assessment**, patient-centered research priorities, 32(5):674–684
- Patient participation**
 developing patient and community advisory groups, 32(5):663–673
 portal-based advance care planning outreach, 32(6):925–930
 process for setting research priorities, 32(5):655–662
- Patient positioning**, sources of error in blood pressure measurement, 32(5):732–738
- Patient preference**, adopting advance directives in primary care, 32(2):168–179
- Patient Protection and Affordable Care Act (ACA)**

- alternative payment impact on visits and scheduling, 32(4):539–549
preexisting conditions and, 32(6):883–889
- Patient readmission**
family medicine transitional care service line, 32(4):619–627
outpatient follow-up and hospital readmissions, 32(3):353–361
patients frequently readmitted versus other admitted patients, 32(1):58–64
- Patient safety**
laboratory safety toolkit in primary care, 32(2):136–145
opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
- Patient satisfaction**, patient characteristics and making requests, 32(2):201–208
- Patient transfer**, family medicine transitional care service line, 32(4):619–627
- Pay for performance**, declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
- Pediatric obesity**, lifestyle intervention programs for diabetes, 32(4):596–600
- Personal satisfaction**, eConsult service implementation in a new health region, 32(2):146–157
- Personality**, patient characteristics and making requests, 32(2):201–208
- Pharmacists**
community, role in reducing suicide risk, 32(6):763–767
medication management improves providers' work-life, 32(4):462–473
- Pharmacoepidemiology**, high-risk medication without documented contraception, 32(4):474–480
- Pharmacology**, drug-induced hyperpigmentation, 32(4):628–638
- Physical fitness**, counseling on fitness and strengthening, 32(1):103–107
- Physician assistants**, provider perspectives of patient experiences in imaging, 32(3):392–397
- Physician-patient relations**
advance care planning conversations, 32(6):835–846
discussions about driving safety, 32(4):607–613
men's depression treatment and, 32(1):13–19
patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
portal-based advance care planning outreach, 32(6):925–930
- Pneumonia**, community-acquired, ruling out, 32(2):234–247
- Point-of-care systems**, prediabetes at an academic practice, 32(4):505–512
- Point-of-care testing**, rapid point of care test for influenza, 32(2):226–233
- Politics**, AAFP values and policies, 32(6):948–950
- Polymerase chain reaction**, rapid point of care test for influenza, 32(2):226–233
- Polypharmacy**, optimal prescribing in patients with dementia, 32(3):383–391
- Population health**
accountable care savings in deprived communities, 32(6):913–922
blood donation rates in the United States 1999–2016, 32(5):746–748
socioeconomics and hospitalization of veterans, 32(6):890–903
- Potentially inappropriate medications**, optimal prescribing in patients with dementia, 32(3):383–391
- Practice-based research**
developing patient and community advisory groups, 32(5):663–673
establishing a PBRN stakeholder engagement infrastructure, 32(5):695–704
parent participation in child health research, 32(5):685–694
patient-centered research priorities, 32(5):674–684
process for setting research priorities, 32(5):655–662
team-based breastfeeding support, 32(6):818–826
- Practice management**
drug-induced hyperpigmentation, 32(4):628–638
team-based breastfeeding support, 32(6):818–826
- Preconception care**, high-risk medication without documented contraception, 32(4):474–480
- Prediabetes**
diabetes screening and treatment during office visits, 32(2):209–217
prediabetes at an academic practice, 32(4):505–512
status, awareness of, subsequent health behavior and, 32(1):20–27
- Prediabetic state**, lifestyle intervention programs for diabetes, 32(4):596–600
- Preexisting condition coverage**, Affordable Care Act and, 32(6):883–889
- Pregnancy**, high-risk medication without documented contraception, 32(4):474–480
- Prehypertension**, sources of error in blood pressure measurement, 32(5):732–738
- Premature mortality**, counseling on fitness and strengthening, 32(1):103–107
- Prenatal care**, high-risk medication without documented contraception, 32(4):474–480
- Prescriptions**, opioid prescribing and the pain-depression dyad, 32(4):614–618
- Prevalence**
childhood atopic dermatitis in primary care, 32(2):191–200
diabetes screening and treatment during office visits, 32(2):209–217
psychotherapeutic medication misuse screening, 32(2):272–278
veterans' reporting of cardiovascular disease, 32(1):50–57
- Preventive medicine**
suboptimal statin use for secondary prevention, 32(6):807–817
team-based breastfeeding support, 32(6):818–826
- Primary care physicians**
accountable care savings in deprived communities, 32(6):913–922
advance care planning conversations, 32(6):835–846
basal insulin in primary care, 32(3):431–447
family medicine transitional care service line, 32(4):619–627
hepatitis C and expanding the role of primary care, 32(3):428–430
low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
patient characteristics and making requests, 32(2):201–208
physician characteristics and low value care spending, 32(2):218–225
providing e-visits, 32(6):868–875
retail clinic use among Medicare beneficiaries, 32(4):531–538
- Primary health care**
accountable care savings in deprived communities, 32(6):913–922
adopting advance directives in, 32(2):168–179
alternative payment impact on visits and scheduling, 32(4):539–549
basal insulin in, 32(3):431–447
burnout, clinic capacity, and social needs, 32(1):69–78
Canadian HIV care settings as PC-MHs, 32(2):158–167
childhood atopic dermatitis in, 32(2):191–200
Colorado asthma toolkit implementation, 32(1):37–49
colorectal cancer screening in a FQHC, 32(2):180–190
counseling on fitness and strengthening, 32(1):103–107
depression treatment for Spanish and English speakers, 32(6):904–912
developing patient and community advisory groups, 32(5):663–673
disease-modifying treatments for Alzheimer disease, 32(6):931–940
eConsult service implementation in a new health region, 32(2):146–157
emergency department high utilizers, 32(2):264–268
enhanced shared decision making for depression, 32(1):6–9
hair care barrier to physical activity, 32(6):944–947

- high-risk medication without documented contraception, 32(4):474–480
- housing insecurity and access to care, 32(4):521–530
- implementation of patient-team partnership, 32(4):490–504
- implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- knee pain and osteoarthritis among overweight women, 32(4):575–584
- laboratory safety toolkit in, 32(2):136–145
- LARC provision by family physicians, 32(1):10–12
- low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- medication management improves providers' work-life, 32(4):462–473
- monitoring the metabolic effects of antipsychotics, 32(3):418–423
- natural history of back pain in older adults, 32(6):781–789
- new allopathic medical schools train fewer family physicians, 32(5):653–654
- opioid management in the safety-net, 32(3):375–382
- opioid prescribing and the pain-depression dyad, 32(4):614–618
- opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
- optimal prescribing in patients with dementia, 32(3):383–391
- parent participation in child health research, 32(5):685–694
- patient BMI knowledge in a rural population, 32(3):413–417
- patient-centered research priorities, 32(5):674–684
- patient characteristics and making requests, 32(2):201–208
- patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
- physicians trained in teaching health centers, 32(2):134–135
- portal-based advance care planning outreach, 32(6):925–930
- practice characteristics and patient self-management support, 32(3):329–340
- prediabetes at an academic practice, 32(4):505–512
- process for setting research priorities, 32(5):655–662
- provider perspectives of patient experiences in imaging, 32(3):392–397
- psychotherapeutic medication misuse screening, 32(2):272–278
- quality improvement measures by type of practice, 32(3):398–407
- quality of the after-visit summary (AVS), 32(1):65–68
- reporting specialty status in research, 32(6):941–943
- retail clinic use among Medicare beneficiaries, 32(4):531–538
- role of pharmacists in reducing suicide risk, 32(6):763–767
- social and behavioral determinants of health screening, 32(3):297–306
- social risk stratification in primary care, 32(4):585–595
- socioeconomics and hospitalization of veterans, 32(6):890–903
- sugar-sweetened beverage intake, rural populations, 32(4):601–606
- systems-based improvements to opioid management, 32(5):715–723
- team-based breastfeeding support, 32(6):818–826
- text messages for HIV testing in, 32(3):408–412
- training vulnerable patients to use a patient portal, 32(2):248–258
- translating self-management support into primary care, 32(3):341–352
- treatment initiation for opioid use disorder, 32(5):724–731
- visit entropy associated with diabetic control outcome, 32(5):739–745
- walk-in psychiatric model for integrated care, 32(4):481–489
- yoga mentions in the electronic health record, 32(6):790–800
- Process measures**, quality improvement measures by type of practice, 32(3):398–407
- Progestins**, broken Nexplanon® rods, 32(2):269–271
- Proof of concept study**, text messages for HIV testing in primary care, 32(3):408–412
- Prospective studies**
natural history of back pain in older adults, 32(6):781–789
ruling out community-acquired pneumonia, 32(2):234–247
- Psychiatry**
monitoring the metabolic effects of antipsychotics, 32(3):418–423
walk-in psychiatric model for integrated care, 32(4):481–489
- Psychometrics**, validating the ABFM certification examination, 32(6):876–882
- Psychotherapy**
depression treatment for Spanish and English speakers, 32(6):904–912
doctor-patient relationships and men's depression treatment, 32(1):13–19
- Psychotropic drugs**, psychotherapeutic medication misuse screening, 32(2):272–278
- Public health**
emergency department high utilizers, 32(2):264–268
monitoring the metabolic effects of antipsychotics, 32(3):418–423
opioid prescribing and the pain-depression dyad, 32(4):614–618
role of pharmacists in reducing suicide risk, 32(6):763–767
social and behavioral determinants of health screening, 32(3):297–306
- Qualitative research**
depression treatment for Spanish and English speakers, 32(6):904–912
laboratory safety toolkit in primary care, 32(2):136–145
optimal prescribing in patients with dementia, 32(3):383–391
palliative health care for the homeless, 32(6):858–867
patient-centered research priorities, 32(5):674–684
provider perspectives of patient experiences in imaging, 32(3):392–397
social and behavioral determinants of health screening, 32(3):297–306
social risk stratification in primary care, 32(4):585–595
systems-based improvements to opioid management, 32(5):715–723
- Quality improvement**
after-visit summaries, 32(1):65–68
family medicine transitional care service line, 32(4):619–627
implementation of patient-team partnership, 32(4):490–504
implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
laboratory safety toolkit in primary care, 32(2):136–145
measures, by type of practice, 32(3):398–407
opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
portal-based advance care planning outreach, 32(6):925–930
practice characteristics and patient self-management support, 32(3):329–340
systems-based improvements to opioid management, 32(5):715–723
- Quality of life**
adopting advance directives in primary care, 32(2):168–179
natural history of back pain in older adults, 32(6):781–789
- Racism**, conversations on Twitter about diversity in medicine, 32(1):28–36
- Radiology**, early imaging for back pain in older adults, 32(6):773–780
- Referral and consultation**
depression treatment for Spanish and English speakers, 32(6):904–912
eConsult service implementation in a new health region, 32(2):146–157
patient characteristics and making requests, 32(2):201–208
- Registries**, implementation of patient-team partnership, 32(4):490–504
- Reproductive health**, high-risk medication without documented contraception, 32(4):474–480
- Rescheduling**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Research report**, reporting specialty status in research, 32(6):941–943

- Residency**
 declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
 physicians trained in teaching health centers, 32(2):134–135
- Resource allocation**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Retail clinics**, use among Medicare beneficiaries, 32(4):531–538
- Retrospective studies**
 opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
 outpatient follow-up and hospital readmissions, 32(3):353–361
 visit entropy associated with diabetic control outcome, 32(5):739–745
- Rheumatology**, knee pain and osteoarthritis among overweight women, 32(4):575–584
- Risk adjustment**, social risk stratification in primary care, 32(4):585–595
- Risk assessment**
 dietary inflammatory index and diabetes severity, 32(6):801–806
 discussions about driving safety, 32(4):607–613
 optimal prescribing in patients with dementia, 32(3):383–391
- Risk factors**, patient BMI knowledge in a rural population, 32(3):413–417
- Risk-taking**, housing insecurity and access to care, 32(4):521–530
- Rural health**
 emergency medicine staffing, 32(3):292–294
 endoscopic care by family physicians, 32(4):460–461
 family physicians in emergent and urgent care, 32(3):295–296
 patient BMI knowledge in a rural population, 32(3):413–417
 veterans perceptions of access to primary care, 32(5):749–751
- Rural population**
 patient BMI knowledge in, 32(3):413–417
 sugar-sweetened beverage intake, 32(4):601–606
- Screening**
 colorectal cancer
 endoscopic services, 32(4):454–456
 in a FQHC, 32(2):180–190
 Latino-based messaging, 32(3):307–317
 health, social and behavioral determinants of, 32(3):297–306
- Seborrhic keratosis**, benign skin lesions and TADA, 32(1):96–102
- Secondary prevention**, suboptimal statin use for secondary prevention, 32(6):807–817
- Self-assessment**, physician opinions about ABFM self-assessment modules, 32(1):79–88
- Self care**
 practice characteristics and patient self-management support, 32(3):329–340
 translating self-management support into primary care, 32(3):341–352
- Self-management**, support, translating into primary care, 32(3):341–352
- Self report**
 acupuncture training and physician burnout, 32(2):259–263
 sugar-sweetened beverage intake, rural populations, 32(4):601–606
 veterans perceptions of access to primary care, 32(5):749–751
- Sensitivity and specificity**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Sexism**, conversations on Twitter about diversity in medicine, 32(1):28–36
- Shared decision making**, provider perspectives of patient experiences in imaging, 32(3):392–397
- Sigmoidoscopy**, colorectal cancer screening in a FQHC, 32(2):180–190
- Skin cancer**, benign skin lesions and TADA, 32(1):96–102
- Skin care**, childhood atopic dermatitis in primary care, 32(2):191–200
- Smoking cessation**, electronic cigarettes and, 32(4):567–574
- Smoking reduction**, electronic cigarettes and, 32(4):567–574
- Social determinants of health**
 hair care barrier to physical activity, 32(6):944–947
 housing insecurity and access to care, 32(4):521–530
 patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
 social and behavioral determinants of health screening, 32(3):297–306
 socioeconomic and hospitalization of veterans, 32(6):890–903
- Social media**, conversations on Twitter about diversity in medicine, 32(1):28–36
- Social work**, end of life services for the homeless, 32(6):847–857
- Socioeconomic status**, hospitalization of veterans and, 32(6):890–903
- Spain**, drug-induced hyperpigmentation, 32(4):628–638
- Spanish-speaking patients**
 depression treatment for, 32(6):904–912
 Hispanic health story from Texas, 32(1):115–117
- Special communications**
 Does My Patient with Multiple Comorbidities Have Heart Failure with Preserved Ejection Fraction, and Does it Matter?, 32(3):424–427
 Exercise Is Medicine: Primary Care Counseling on Aerobic Fitness and Muscle Strengthening, 32(1):103–107
 The Potential Emergence of Disease-Modifying Treatments for Alzheimer Disease: The Role of Primary Care in Managing the Patient Journey, 32(6):931–940
- Specialization**, reporting specialty status in research, 32(6):941–943
- Standard of care**
 depression treatment for Spanish and English speakers, 32(6):904–912
 Hispanic health story from Texas, 32(1):115–117
- Statins**, suboptimal use, for secondary prevention, 32(6):807–817
- Street drugs**, identifying substance use using frequency questions, 32(4):550–558
- Stroke volume**, heart failure with preserved ejection fraction, 32(3):424–427
- Students**, new allopathic medical schools train fewer family physicians, 32(5):653–654
- Substance abuse detection**, psychotherapeutic medication misuse screening, 32(2):272–278
- Substance-related disorders**
 identifying substance use using frequency questions, 32(4):550–558
 opioid management in the safety-net, 32(3):375–382
 palliative health care for the homeless, 32(6):858–867
- Sugars**, sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Suicidal ideation**, role of pharmacists in reducing suicide risk, 32(6):763–767
- Suicide**, role of pharmacists in reducing risk, 32(6):763–767
- Surveillance systems**, housing insecurity and access to care, 32(4):521–530
- Surveys and questionnaires**
 ABFM's data collection method for tracking specialty, 32(1):89–95
 accountable care savings in deprived communities, 32(6):913–922
 doctor-patient relationships and men's depression treatment, 32(1):13–19
 end of life services for the homeless, 32(6):847–857
 family physicians providing e-visits, 32(6):868–875
 hair care barrier to physical activity, 32(6):944–947
 identifying substance use using frequency questions, 32(4):550–558
 implementation of patient-team partnership, 32(4):490–504
 lifestyle intervention programs for diabetes, 32(4):596–600
 monitoring the metabolic effects of antipsychotics, 32(3):418–423
 natural history of back pain in older adults, 32(6):781–789
 patient BMI knowledge in a rural population, 32(3):413–417
 patient-centered research priorities, 32(5):674–684
 physician opinions about ABFM self-assessment modules, 32(1):79–88

- prediabetes at an academic practice, 32(4):505–512
- social risk stratification in primary care, 32(4):585–595
- suboptimal statin use for secondary prevention, 32(6):807–817
- team-based breastfeeding support, 32(6):818–826
- veterans' reporting of cardiovascular disease, 32(1):50–57
- Sustained virologic response**, hepatitis C and expanding the role of primary care, 32(3):428–430
- Sweetening agents**, sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Systematic review**, ruling out community-acquired pneumonia, 32(2):234–247
- Teaching health centers**, 32(2):130–133, 32(2):134–135
- Teaching hospitals** advance care planning billing code use, 32(6):827–834
- Telemedicine**
- family physicians providing e-visits, 32(6):868–875
 - text messages for HIV testing in primary care, 32(3):408–412
 - training vulnerable patients to use a patient portal, 32(2):248–258
- Telephone**, visit entropy associated with diabetic control outcome, 32(5):739–745
- Teratogens**, high-risk medication without documented contraception, 32(4):474–480
- Terminal care**
- advance care planning conversations, 32(6):835–846
 - end of life services for the homeless, 32(6):847–857
 - palliative health care for the homeless, 32(6):858–867
- Tertiary care centers**, advance care planning billing code use, 32(6):827–834
- Texas**, Hispanic health story from, 32(1):115–117
- Text messaging**
- for colorectal cancer screening, 32(3):307–317, 32(3):318–328
 - for HIV testing in primary care, 32(3):408–412
- Tobacco**, identifying substance use using frequency questions, 32(4):550–558
- Tobacco products**, electronic cigarettes and smoking reduction, 32(4):567–574
- Tramadol**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Transitional care**, family medicine transitional care service line, 32(4):619–627
- Triage amalgamated dermoscopic algorithm (TADA)**, and benign skin lesions, 32(1):96–102
- Twitter**, conversations about diversity in medicine, 32(1):28–36
- Type 2 diabetes mellitus**
- basal insulin in primary care, 32(3):431–447
 - lifestyle intervention programs for diabetes, 32(4):596–600
 - practice characteristics and patient self-management support, 32(3):329–340
 - prediabetes at an academic practice, 32(4):505–512
 - translating self-management support into primary care, 32(3):341–352
 - visit entropy associated with diabetic control outcome, 32(5):739–745
- Underserved populations**, walk-in psychiatric model for integrated care, 32(4):481–489
- United States Agency for Health care Research and Quality**, laboratory safety toolkit in primary care, 32(2):136–145
- United States Department of Veterans Affairs**
- socioeconomics and hospitalization of veterans, 32(6):890–903
 - veterans perceptions of access to primary care, 32(5):749–751
- Urban areas**, urgent care, family physicians' contributions, 32(3):295–296
- Urgent care**, urban, family physicians' contributions, 32(3):295–296
- Validity**, ABFM certification examination, 32(6):876–882
- Veterans**
- hospitalization of, socioeconomics and, 32(6):890–903
 - perceptions of access to primary care, 32(5):749–751
 - reporting of cardiovascular disease, 32(1):50–57
- Violence**, values and policies of AAFP, 32(6):948–950
- Virginia**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Vulnerable populations**
- colorectal cancer screening in a FQHC, 32(2):180–190
 - monitoring the metabolic effects of antipsychotics, 32(3):418–423
 - opioid management in the safety-net, 32(3):375–382
- Washington**, socioeconomics and hospitalization of veterans, 32(6):890–903
- Weight gain**, basal insulin in primary care, 32(3):431–447
- Women's health**
- hair care barrier to physical activity, 32(6):944–947
 - LARC provision by family physicians, 32(1):10–12
 - team-based breastfeeding support, 32(6):818–826
- Workflow**
- portal-based advance care planning outreach, 32(6):925–930
 - social and behavioral determinants of health screening, 32(3):297–306
 - social risk stratification in primary care, 32(4):585–595
- Workload**, medication management improves providers' work-life, 32(4):462–473
- Yoga**, mentions in the electronic health record, 32(6):790–800