

# Subject Index to Volume 32, 2019

- Academic medical centers**, yoga mentions in the electronic health record, 32(6):790–800
- Access to health care**  
housing insecurity and, 32(4):521–530  
team-based breastfeeding support, 32(6):818–826
- Accountable care organizations**, serving deprived communities, savings rate and, 32(6):913–922
- Acquired immunodeficiency syndrome (AIDS)**  
Canadian HIV care settings as PCMHs, 32(2):158–167  
text messages for HIV testing in primary care, 32(3):408–412
- Acupuncture therapy**, training, physician burnout and, 32(2):259–263
- Addiction medicine**  
opioid management, systems-based improvements to, 32(5):715–723  
opioid prescribing  
intervention in chronic noncancer pain, 32(4):559–566  
pain-depression dyad and, 32(4):614–618  
regulation on, in an urban safety-net, 32(3):362–374  
opioid use disorder, treatment initiation for, 32(5):724–731
- Adolescence**, identifying substance use using frequency questions, 32(4):550–558
- Advance care planning**  
billing code use, 32(6):827–834  
conversations, 32(1):108–114, 32(6):835–846  
outreach, portal-based, 32(6):925–930
- Advance directives**  
adopting, in primary care, 32(2):168–179  
portal-based advance care planning outreach, 32(6):925–930
- Adverse decision making**, optimal prescribing in patients with dementia, 32(3):383–391
- Altruism**, parent participation in child health research, 32(5):685–694
- Alzheimer disease**, disease-modifying treatments for, 32(6):931–940
- Ambulatory care**, retail clinic use among Medicare beneficiaries, 32(4):531–538
- American Academy of Family Physicians (AAFP)**, values and policies of, 32(6):948–950
- American Board of Family Medicine (ABFM)**  
certification examination, validating, 32(6):876–882  
data collection method for tracking specialty, 32(1):89–95  
election of new officers and board members, 32(4):643–645  
role in continuing education for family physicians, 32(5):756–758  
self-assessment modules, physician opinions about, 32(1):79–88
- Amyloid  $\beta$ -peptides**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Analgesics**, natural history of back pain in older adults, 32(6):781–789
- Antidepressants**, enhanced shared decision making for depression, 32(1):6–9
- Antiviral agents**  
hepatitis C and the role of primary care, 32(3):428–430  
rapid point of care test for influenza, 32(2):226–233
- Appointments and schedules**  
alternative payment impact on, 32(4):539–549  
veterans perceptions of access to primary care, 32(5):749–751
- Asthma**, Colorado asthma toolkit implementation, 32(1):37–49
- Atherosclerosis**, suboptimal statin use for secondary prevention, 32(6):807–817
- Atopic dermatitis**, childhood, in primary care, 32(2):191–200
- Attitudes**, prediabetes at an academic practice, 32(4):505–512
- Automobile driving**, safety, discussions about, 32(4):607–613
- Back pain**, natural history of, in older adults, 32(6):781–789
- Behavior therapy**, translating self-management support into primary care, 32(3):341–352
- Behavioral medicine**  
depression treatment for Spanish and English speakers, 32(6):904–912  
outpatient follow-up and hospital readmissions, 32(3):353–361  
translating self-management support into primary care, 32(3):341–352  
walk-in psychiatric model for integrated care, 32(4):481–489  
yoga mentions in the electronic health record, 32(6):790–800
- Behavioral risk factors**, housing insecurity and access to care, 32(4):521–530
- Beverages**, sugar-sweetened, intake, rural populations, 32(4):601–606
- Biomarkers**  
dietary inflammatory index and diabetes severity, 32(6):801–806  
disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Blood donors**, blood donation rates in the United States 1999–2016, 32(5):746–748
- Blood glucose**, levels, awareness of prediabetes status and subsequent health behavior, 32(1):20–27
- Blood pressure**, measurement, sources of error in, 32(5):732–738
- Blood transfusion**, blood donation rates in the United States 1999–2016, 32(5):746–748
- Body composition**, patient BMI knowledge in a rural population, 32(3):413–417
- Body mass index**  
dietary inflammatory index and diabetes severity, 32(6):801–806  
knee pain and osteoarthritis among overweight women, 32(4):575–584  
knowledge, patient, in a rural population, 32(3):413–417  
lifestyle intervention programs for diabetes, 32(4):596–600  
sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Body weight**, awareness of prediabetes status and subsequent health behavior, 32(1):20–27
- Boston, Massachusetts**, social and behavioral determinants of health screening, 32(3):297–306
- Breastfeeding**, team-based support, 32(6):818–826
- Burnout, professional**  
acupuncture training and, 32(2):259–263  
clinic capacity and social needs, 32(1):69–78  
medication management, impact on providers' work-life, 32(4):462–473
- California**  
outpatient follow-up and hospital readmissions, 32(3):353–361  
practice characteristics and patient self-management support, 32(3):329–340
- Canada**  
doctor-patient relationships and men's depression treatment, 32(1):13–19  
eConsult service implementation, 32(2):146–157  
HIV care settings as PCMHs, 32(2):158–167  
palliative health care for the homeless, 32(6):858–867
- Cannabis**, identifying substance use using frequency questions, 32(4):550–558
- Cardiovascular disease**  
quality improvement measures by type of practice, 32(3):398–407  
suboptimal statin use for secondary prevention, 32(6):807–817  
veterans' reporting of, 32(1):50–57
- Centers for Medicare and Medicaid Services (US)**, accountable care savings in deprived communities, 32(6):913–922
- Certification**  
ABM's data collection method for tracking specialty, 32(1):89–95  
physician opinions about ABFM self-assessment modules, 32(1):79–88

validating the ABFM certification examination, 32(6):876–882

### Child health

high-risk medication without documented contraception, 32(4):474–480  
parent participation in child health research, 32(5):685–694  
patient-centered research priorities, 32(5):674–684  
research, parent participation in, 32(5):685–694

### Chronic disease

Canadian HIV care settings as PC-MHs, 32(2):158–167  
housing insecurity and access to care, 32(4):521–530  
knee pain and osteoarthritis among overweight women, 32(4):575–584  
lifestyle intervention programs for diabetes, 32(4):596–600  
management, yoga mentions in the electronic health record, 32(6):790–800  
opioid management in the safety-net, 32(3):375–382  
practice characteristics and patient self-management support, 32(3):329–340  
preexisting conditions and the Affordable Care Act, 32(6):883–889  
retail clinic use among Medicare beneficiaries, 32(4):531–538  
social risk stratification in primary care, 32(4):585–595  
suboptimal statin use for secondary prevention, 32(6):807–817  
training vulnerable patients to use a patient portal, 32(2):248–258  
translating self-management support into primary care, 32(3):341–352  
visit entropy associated with diabetic control outcome, 32(5):739–745

### Chronic pain

management, opioid prescribing intervention in chronic noncancer pain, 32(4):559–566  
opioid management in the safety-net, 32(3):375–382  
systems-based improvements to opioid management, 32(5):715–723

**Cigarette smoking**, electronic cigarettes and smoking reduction, 32(4):567–574

**Clinical decision making**, depression treatment for Spanish and English speakers, 32(6):904–912

**Codeine-combination**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374

**Cognitive dysfunction**, disease-modifying treatments for Alzheimer disease, 32(6):931–940

### Cohort studies

discussions about driving safety, 32(4):607–613  
patient BMI knowledge in a rural population, 32(3):413–417  
prediabetes at an academic practice, 32(4):505–512

preexisting conditions and the Affordable Care Act, 32(6):883–889  
socioeconomics and hospitalization of veterans, 32(6):890–903

**Colon cancer**, screening, messaging for, 32(3):307–317, 32(3):318–328

**Colonoscopy**, colorectal cancer screening in a FQHC, 32(2):180–190

### Colorado

asthma toolkit implementation, 32(1):37–49  
developing patient and community advisory groups, 32(5):663–673  
implementation of patient-team partnership, 32(4):490–504  
patient-centered research priorities, 32(5):674–684  
practice characteristics and patient self-management support, 32(3):329–340  
process for setting research priorities, 32(5):655–662  
social risk stratification in primary care, 32(4):585–595

**Colorectal cancer**, screening endoscopic services, 32(4):454–456 in a FQHC, 32(2):180–190  
messaging for, 32(3):307–317, 32(3):318–328

### Commentary

Commentary Response for “LARC Provision by Family Physicians: Low But on the Rise,” 32(1):4–5  
Community Pharmacists as Partners in Reducing Suicide Risk, 32(6):763–767  
Declining Presence of Family Physicians in Hospital-Based Care: A Major Concern or Totally Makes Sense?, 32(6):768–770  
Endoscopic Services in the United States: By Whom, for What, and Why?, 32(4):454–456  
Is It Time to Prioritize Diabetes Prevention in Practice?, 32(4):457–459  
New Allopathic Medical Schools and Family Physicians, 32(5):651–652  
No Magic Pill: A Prescription for Enhanced Shared Decision-Making for Depression Treatment, 32(1):6–9  
Pawn and King Play: A Stalemate on Rural Emergency Medicine Staffing, 32(3):292–294  
Practice-Based Research Today: A Changing Primary Care Landscape Requires Changes in Practice-Based Research Network (PBRN) Research, 32(5):647–650  
Response for “LARC Provision by Family Physicians: Low But on the Rise,” 32(1):4–5  
Stuck in Graduate Medical Education Traffic? Teaching Health Centers Are Family Medicine’s High-Occupancy-Vehicle Lane, 32(2):130–133

### Communication

discussions about driving safety, 32(4):607–613  
patient-defined visit priorities: psychosocial versus medical, 32(4):513–520

**Communication barriers**, a Hispanic health story from Texas, 32(1):115–117

**Community-acquired infections**, pneumonia, ruling out, 32(2):234–247

### Community health centers

Latino-based colorectal cancer screening messaging, 32(3):307–317  
messaging for colorectal cancer screening, 32(3):318–328  
preexisting conditions and the Affordable Care Act, 32(6):883–889  
social and behavioral determinants of health screening, 32(3):297–306

**Community health nursing**, team-based breastfeeding support, 32(6):818–826

### Community medicine

developing patient and community advisory groups, 32(5):663–673  
high-risk medication without documented contraception, 32(4):474–480  
process for setting research priorities, 32(5):655–662  
role of pharmacists in reducing suicide risk, 32(6):763–767  
social and behavioral determinants of health screening, 32(3):297–306  
team-based breastfeeding support, 32(6):818–826

### Community networks

developing patient and community advisory groups, 32(5):663–673  
process for setting research priorities, 32(5):655–662

### Comorbidity

heart failure with preserved ejection fraction, 32(3):424–427  
optimal prescribing in patients with dementia, 32(3):383–391  
socioeconomics and hospitalization of veterans, 32(6):890–903

**Comprehensive medication management (CMM)**, impact on providers’ work-life, 32(4):462–473

**Continuity of care**, retail clinic use among Medicare beneficiaries, 32(4):531–538

### Continuity of patient care

declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772  
role of pharmacists in reducing suicide risk, 32(6):763–767  
team-based breastfeeding support, 32(6):818–826  
visit entropy associated with diabetic control outcome, 32(5):739–745

### Contraception

broken Nexplanon rods, 32(2):269–271

- high-risk medication without documented contraception, 32(4):474–480
- LARC provision by family physicians, 32(1):10–12
- Contraceptive devices**, broken Nexplanon rods, 32(2):269–271
- Correspondence**
- better doctor-patient relationships are associated with men choosing more active depression treatment, 32(4):640–641
  - the complex interpretation and management of zika virus test results, 32(2):280–281
  - family medicine and obstetrics: let us stop pretending, 32(2):279–280
  - how evolving United States payment models influence primary care and its impact on the quadruple aim: the need for health equity, 32(1):118–119
  - if we do not ask, they will not tell: screening for urinary and fecal incontinence by primary care providers, 32(1):119–120
  - impact of a rapid point of care test for influenza on guideline consistent care and antibiotic use, 32(4):641–642
  - impact of medical scribes in primary care on productivity, face-to-face time, and patient comfort, 32(1):120
  - physician burnout and higher clinic capacity to address patients' social needs, 32(5):752–753
  - a reflective case study in family medicine advance care planning conversations, 32(4):639–640
  - signs and symptoms that rule out community-acquired pneumonia in outpatient adults: a systematic review and meta-Analysis, 32(5):753
  - timely outpatient follow-up is associated with fewer hospital readmissions among patients with behavioral health conditions, 32(5):754–755
- Counseling**
- on fitness and strengthening, 32(1):103–107
  - high-risk medication without documented contraception, 32(4):474–480
  - practice characteristics and patient self-management support, 32(3):329–340
  - team-based breastfeeding support, 32(6):818–826
  - translating self-management support into primary care, 32(3):341–352
- Cross-cultural comparison**, depression treatment for Spanish and English speakers, 32(6):904–912
- Cross-sectional analysis**, quality improvement measures by type of practice, 32(3):398–407
- Cross-sectional studies**
- accountable care savings in deprived communities, 32(6):913–922
  - benign skin lesions and TADA, 32(1):96–102
  - dietary inflammatory index and diabetes severity, 32(6):801–806
  - discussions about driving safety, 32(4):607–613
  - family physicians providing e-visits, 32(6):868–875
  - implementation of patient-team partnership, 32(4):490–504
  - patient BMI knowledge in a rural population, 32(3):413–417
- Cultural competency**, hair care barrier to physical activity, 32(6):944–947
- Current procedural terminology**, advance care planning billing code use, 32(6):827–834
- Data warehousing**, advance care planning billing code use, 32(6):827–834
- Decision making**
- AAFP and politics, 32(6):948–950
  - adverse, optimal prescribing in patients with dementia, 32(3):383–391
  - depression treatment for Spanish and English speakers, 32(6):904–912
  - enhanced shared, for depression, 32(1):6–9
- Delivery of health care**
- Canadian HIV care settings as PC-MHs, 32(2):158–167
  - emergency department high utilizers, 32(2):264–268
  - family physicians providing e-visits, 32(6):868–875
  - palliative health care for the homeless, 32(6):858–867
  - quality improvement measures by type of practice, 32(3):398–407
  - team-based breastfeeding support, 32(6):818–826
- Dementia**
- disease-modifying treatments for Alzheimer disease, 32(6):931–940
  - patients with, optimal prescribing in, 32(3):383–391
- Demography**, emergency department high utilizers, 32(2):264–268
- Depersonalization**, acupuncture training and physician burnout, 32(2):259–263
- Depression**
- enhanced shared decision making for, 32(1):6–9
  - opioid prescribing and the pain-depression dyad, 32(4):614–618
  - treatment, men's, and doctor-patient relationships, 32(1):13–19
  - treatment of, for Spanish and English speakers, 32(6):904–912
- Dermatofibroma**, benign skin lesions and TADA, 32(1):96–102
- Dermatology**, drug-induced hyperpigmentation, 32(4):628–638
- Dermoscopy**, benign skin lesions and TADA, 32(1):96–102
- Diabetes mellitus**
- basal insulin in primary care, 32(3):431–447
  - low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
  - monitoring the metabolic effects of antipsychotics, 32(3):418–423
  - practice characteristics and patient self-management support, 32(3):329–340
  - prevention, prioritizing, 32(4):457–459
  - screening and treatment during office visits, 32(2):209–217
  - severity, dietary inflammatory index and, 32(6):801–806
  - translating self-management support into primary care, 32(3):341–352
- Diagnostic imaging**, early imaging for back pain in older adults, 32(6):773–780
- Diagnostic tests**
- disease-modifying treatments for Alzheimer disease, 32(6):931–940
  - rapid point of care test for influenza, 32(2):226–233
- Diet**, dietary inflammatory index and diabetes severity, 32(6):801–806
- Disease management**
- practice characteristics and patient self-management support, 32(3):329–340
  - social risk stratification in primary care, 32(4):585–595
  - translating self-management support into primary care, 32(3):341–352
- Disease progression**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Doctor-patient relations**
- advance care planning conversations, 32(6):835–846
  - discussions about driving safety, 32(4):607–613
  - men's depression treatment and, 32(1):13–19
  - patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
  - portal-based advance care planning outreach, 32(6):925–930
- Drug abuse**, psychotherapeutic medication misuse screening, 32(2):272–278
- Drug overdose**, opioid prescribing and the pain-depression dyad, 32(4):614–618
- Drug reaction**, optimal prescribing in patients with dementia, 32(3):383–391
- Early detection of cancer**
- colorectal cancer screening in a FQHC, 32(2):180–190
  - Latino-based colorectal cancer screening messaging, 32(3):307–317
  - messaging for colorectal cancer screening, 32(3):318–328
- Editorial Office News and Notes**
- Content Usage and the Most Frequently Read Articles of 2018, 32(3):288–291



Peer Reviewers for the Journal of the American Board of Family Medicine in 2018, 32(2):126–129

## Editor's Notes

Conversations, Communication, and Counseling Are Key Family Medicine Tactics to Improve Patient Health, 32(1):1–3

Family Medicine: Data Driven Practice with Emphasis on Underserved Patients, 32(3):285–287

Implementing Practice Changes in Family Medicine to Enhance Care and Prevent Disease Progression, 32(4):451–453

Multiple Research Methodologies Can Advance the Science of Family Medicine, 32(2):123–125

New Research on Back Pain, Diet and Diabetes, Advanced Care Planning, and Other Issues Frequently Seen in Family Medicine, 32(6):759–762

## Education

new allopathic medical schools train fewer family physicians, 32(5):653–654

teaching health centers, 32(2):130–133

**Educational measurement**, validating the ABFM certification examination, 32(6):876–882

**Electronic cigarettes**, smoking reduction and, 32(4):567–574

## Electronic health records

advance care planning billing code use, 32(6):827–834

eConsult service implementation in a new health region, 32(2):146–157

portal-based advance care planning outreach, 32(6):925–930

preexisting conditions and the Affordable Care Act, 32(6):883–889

quality of the after-visit summary (AVS), 32(1):65–68

social risk stratification in primary care, 32(4):585–595

training vulnerable patients to use a patient portal, 32(2):248–258

yoga mentions in, 32(6):790–800

**Electronic nicotine delivery systems**, smoking reduction and, 32(4):567–574

**Emergency care**, rural, family physicians' contributions, 32(3):295–296

## Emergency departments

high utilizers, 32(2):264–268

staffing, rural hospitals, 32(3):292–294

**Emergency medicine**, staffing, rural hospitals, 32(3):292–294

## End of life care

adopting advance directives in primary care, 32(2):168–179

portal-based advance care planning outreach, 32(6):925–930

**Endoscopic services**, colorectal cancer screening, 32(4):454–456

**Endoscopy**, endoscopic care by family physicians, 32(4):460–461

**Energy intake**, sugar-sweetened beverage intake, rural populations, 32(4):601–606

**Entropy**, visit entropy associated with diabetic control outcome, 32(5):739–745

**Ethical theory**, advance care planning conversations, 32(1):108–114

## Evidence-based medicine

early imaging for back pain in older adults, 32(6):773–780

implementing EvidenceNow to improve cardiovascular care, 32(5):705–714

**Examination blueprint**, validating the ABFM certification examination, 32(6):876–882

## Exercise

counseling on fitness and strengthening, 32(1):103–107

hair care barrier to physical activity, 32(6):944–947

**Family medicine**, reflections, Hispanic health story from Texas, 32(1):115–117

## Family physicians

ABFM's data collection method for tracking specialty, 32(1):89–95

acupuncture training and physician burnout, 32(2):259–263

advance care planning conversations, 32(1):108–114

allopathic medical schools and, 32(5):653–654

benign skin lesions and TADA, 32(1):96–102

burnout, clinic capacity, and social needs, 32(1):69–78

conversations on Twitter about diversity in medicine, 32(1):28–36

declining presence of, in hospital-based care, 32(6):768–770, 32(6):771–772

doctor-patient relationships and men's depression treatment, 32(1):13–19

in emergent and urgent care, 32(3):295–296

endoscopic care by family physicians, 32(4):460–461

LARC provision by, 32(1):10–12

opioid prescribing intervention in chronic noncancer pain, 32(4):559–566

patient BMI knowledge in a rural population, 32(3):413–417

physician opinions about ABFM self-assessment modules, 32(1):79–88

policies of AAFP and, 32(6):948–950

provider perspectives of patient experiences in imaging, 32(3):392–397

providing e-visits, 32(6):868–875

reporting specialty status in research, 32(6):941–943

systems-based improvements to opioid management, 32(5):715–723

text messages for HIV testing in primary care, 32(3):408–412

trained in teaching health centers, 32(2):134–135

validating the ABFM certification examination, 32(6):876–882

**Fetal hemoglobin**, low hemoglobin A1c and hemoglobinopathy, 32(6):923–924

**Firearms**, values and policies of AAFP, 32(6):948–950

## Focus groups

palliative health care for the homeless, 32(6):858–867

patient-centered research priorities, 32(5):674–684

social and behavioral determinants of health screening, 32(3):297–306

systems-based improvements to opioid management, 32(5):715–723

**Follow-up studies**, natural history of back pain in older adults, 32(6):781–789

## General practitioners

natural history of back pain in older adults, 32(6):781–789

reporting specialty status in research, 32(6):941–943

validating the ABFM certification examination, 32(6):876–882

## Geographic health care financing

accountable care savings in deprived communities, 32(6):913–922

**Geography**, endoscopic care by family physicians, 32(4):460–461

## Geriatrics

early imaging for back pain in older adults, 32(6):773–780

optimal prescribing in patients with dementia, 32(3):383–391

**Glucose**, low hemoglobin A1c and hemoglobinopathy, 32(6):923–924

## Glycohemoglobin A

low hemoglobin A1c and hemoglobinopathy, 32(6):923–924

translating self-management support into primary care, 32(3):341–352

**Graduate medical education**, teaching health centers, 32(2):130–133, 32(2):134–135

**Grounded theory**, opioid management in the safety-net, 32(3):375–382

**Habits**, hair care barrier to physical activity, 32(6):944–947

**Hair care**, as barrier to physical activity, 32(6):944–947

## Harm reduction

end of life services for the homeless, 32(6):847–857

palliative health care for the homeless, 32(6):858–867

## Health behavior

and awareness of prediabetes status, 32(1):20–27

hair care barrier to physical activity, 32(6):944–947

sugar-sweetened beverage intake, rural populations, 32(4):601–606

**Health care costs**, physician characteristics and low value care spending, 32(2):218–225

- Health care disparities**  
end of life services for the homeless, 32(6):847–857  
social and behavioral determinants of health screening, 32(3):297–306  
team-based breastfeeding support, 32(6):818–826  
treatment initiation for opioid use disorder, 32(5):724–731
- Health care reform**, preexisting conditions and the Affordable Care Act, 32(6):883–889
- Health care surveys**  
advance care planning conversations, 32(6):835–846  
diabetes screening and treatment during office visits, 32(2):209–217  
opioid prescribing and the pain-depression dyad, 32(4):614–618  
validating the ABFM certification examination, 32(6):876–882
- Health care systems**, alternative payment impact on visits and scheduling, 32(4):539–549
- Health care use**, outpatient follow-up and hospital readmissions, 32(3):353–361
- Health equity**  
accountable care savings in deprived communities, 32(6):913–922  
end of life services for the homeless, 32(6):847–857
- Health expenditures**, social risk stratification in primary care, 32(4):585–595
- Health information systems**, accountable care savings in deprived communities, 32(6):913–922
- Health insurance**  
alternative payment impact on visits and scheduling, 32(4):539–549  
values and policies of AAFP, 32(6):948–950
- Health literacy**  
patient BMI knowledge in a rural population, 32(3):413–417  
training vulnerable patients to use a patient portal, 32(2):248–258
- Health personnel**, LARC provision by family physicians, 32(1):10–12
- Health policy**  
advance care planning billing code use, 32(6):827–834  
alternative payment impact on visits and scheduling, 32(4):539–549  
declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772  
endoscopic care by family physicians, 32(4):460–461  
preexisting conditions and the Affordable Care Act, 32(6):883–889
- Health promotion**  
team-based breastfeeding support, 32(6):818–826  
translating self-management support into primary care, 32(3):341–352
- Health services**, alternative payment impact on visits and scheduling, 32(4):539–549
- Health services research**, laboratory safety toolkit in primary care, 32(2):136–145
- Heart failure**, with preserved ejection fraction, 32(3):424–427
- Hematology**, low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- Hemoglobinopathies** low hemoglobin A1c and, 32(6):923–924
- Hepatitis C**, expanding the role of primary care and, 32(3):428–430
- Hispanic Americans**  
depression treatment for Spanish and English speakers, 32(6):904–912  
a Hispanic health story from Texas, 32(1):115–117  
Latino-based colorectal cancer screening messaging, 32(3):307–317  
messaging for colorectal cancer screening, 32(3):318–328
- Homeless persons**  
end of life services for, 32(6):847–857  
housing insecurity and access to care, 32(4):521–530  
palliative health care for, 32(6):858–867
- Hospital medicine**, declining presence of family physicians, 32(6):768–770, 32(6):771–772
- Hospitalists**, blood donation rates in the United States 1999–2016, 32(5):746–748
- Hospitalization**  
outpatient follow-up and hospital readmissions, 32(3):353–361  
patients frequently readmitted versus other admitted patients, 32(1):58–64  
of veterans, socioeconomic, 32(6):890–903
- House calls**, advance care planning conversations, 32(6):835–846
- Housing**  
insecurity, access to care and, 32(4):521–530  
issues, socioeconomic and hospitalization of veterans, 32(6):890–903  
palliative health care for the homeless, 32(6):858–867
- Hydrocodone**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Hydrocodone-combination**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Hypercholesterolemia**, suboptimal statin use for secondary prevention, 32(6):807–817
- Hyperglycemia**, basal insulin in primary care, 32(3):431–447
- Hyperlipidemia**, suboptimal statin use for secondary prevention, 32(6):807–817
- Hyperpigmentation**, drug-induced, 32(4):628–638
- Hypertension**, sources of error in blood pressure measurement, 32(5):732–738
- Hypoglycemia**, basal insulin in primary care, 32(3):431–447
- Hypoglycemic drugs**, diabetes screening and treatment during office visits, 32(2):209–217
- Immigrants**, Hispanic health story from Texas, 32(1):115–117
- Implementation**, laboratory safety toolkit in primary care, 32(2):136–145
- Incentive reimbursement**, advance care planning billing code use, 32(6):827–834
- Infant health**, high-risk medication without documented contraception, 32(4):474–480
- Inflammation**, dietary inflammatory index and diabetes severity, 32(6):801–806
- Influenza**, rapid point of care test for, 32(2):226–233
- Information technology**  
practice characteristics and patient self-management support, 32(3):329–340  
social risk stratification in primary care, 32(4):585–595  
training vulnerable patients to use a patient portal, 32(2):248–258
- Inpatients**, declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
- Insulin**, basal, in primary care, 32(3):431–447
- Insurance**, accountable care savings in deprived communities, 32(6):913–922
- Insurance coverage**, preexisting conditions and the Affordable Care Act, 32(6):883–889
- Integrated care**, walk-in psychiatric model for, 32(4):481–489
- Integrative medicine**, acupuncture training and physician burnout, 32(2):259–263
- Interdisciplinary research**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Internet**, portal-based advance care planning outreach, 32(6):925–930
- Internship and residency**, reporting specialty status in research, 32(6):941–943
- Iowa**, advance care planning billing code use, 32(6):827–834
- Job satisfaction**, medication management improves providers' work-life, 32(4):462–473
- Journal impact factor**, reporting specialty status in research, 32(6):941–943
- Journal of the American Board of Family Medicine (JABFM)**  
content usage and the most frequently read articles of 2018, 32(3):288–291  
peer reviewers for, 32(2):126–129

- Korea**, electronic cigarettes and smoking reduction, 32(4):567–574
- Lactation**, team-based breastfeeding support, 32(6):818–826
- Latino immigrants**, Hispanic health story from Texas, 32(1):115–117
- Leadership**, declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
- Lifestyle**  
awareness of prediabetes status and subsequent health behavior, 32(1):20–27  
hair care barrier to physical activity, 32(6):944–947  
practice characteristics and patient self-management support, 32(3):329–340  
prediabetes at an academic practice, 32(4):505–512  
translating self-management support into primary care, 32(3):341–352  
yoga mentions in the electronic health record, 32(6):790–800
- Linear models**, housing insecurity and access to care, 32(4):521–530
- Linguistic issues**, Hispanic health story from Texas, 32(1):115–117
- Lisinopril**, high-risk medication without documented contraception, 32(4):474–480
- Living wills**, adopting advance directives in primary care, 32(2):168–179
- Logistic models**  
depression treatment for Spanish and English speakers, 32(6):904–912  
dietary inflammatory index and diabetes severity, 32(6):801–806  
discussions about driving safety, 32(4):607–613  
family physicians providing e-visits, 32(6):868–875  
high-risk medication without documented contraception, 32(4):474–480  
housing insecurity and access to care, 32(4):521–530  
opioid prescribing intervention in chronic noncancer pain, 32(4):559–566  
suboptimal statin use for secondary prevention, 32(6):807–817
- Logistic regression**  
monitoring the metabolic effects of antipsychotics, 32(3):418–423  
outpatient follow-up and hospital readmissions, 32(3):353–361
- Long-acting reversible contraception (LARC)**, provision by family physicians, 32(1):10–12
- Low back pain**, early imaging for back pain in older adults, 32(6):773–780
- Low-income population**, text messages for HIV testing in primary care, 32(3):408–412
- Major depressive disorder**, enhanced shared decision making for depression, 32(1):6–9
- Meaningful use**, quality of the after-visit summary (AVS), 32(1):65–68
- Medicaid**  
advance care planning billing code use, 32(6):827–834  
alternative payment impact on visits and scheduling, 32(4):539–549  
depression treatment for Spanish and English speakers, 32(6):904–912  
emergency department high utilizers, 32(2):264–268  
preexisting conditions and the Affordable Care Act, 32(6):883–889  
walk-in psychiatric model for integrated care, 32(4):481–489
- Medical education**  
declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772  
validating the ABFM certification examination, 32(6):876–882
- Medical geography** accountable care savings in deprived communities, 32(6):913–922
- Medical informatics**  
social risk stratification in primary care, 32(4):585–595  
yoga mentions in the electronic health record, 32(6):790–800
- Medical schools**, new allopathic medical schools train fewer family physicians, 32(5):653–654
- Medical societies**, conversations on Twitter about diversity in medicine, 32(1):28–36
- Medically uninsured**  
alternative payment impact on visits and scheduling, 32(4):539–549  
outpatient follow-up and hospital readmissions, 32(3):353–361  
preexisting conditions and the Affordable Care Act, 32(6):883–889
- Medicare**  
early imaging for back pain in older adults, 32(6):773–780  
emergency department high utilizers, 32(2):264–268  
physician characteristics and low value care spending, 32(2):218–225  
retail clinic use among Medicare beneficiaries, 32(4):531–538
- Meditation**, yoga mentions in the electronic health record, 32(6):790–800
- MEDLINE**, reporting specialty status in research, 32(6):941–943
- Mental disorders**, monitoring the metabolic effects of antipsychotics, 32(3):418–423
- Mental health**  
depression treatment for Spanish and English speakers, 32(6):904–912  
housing insecurity and access to care, 32(4):521–530  
monitoring the metabolic effects of antipsychotics, 32(3):418–423
- patient-defined visit priorities: psychosocial versus medical, 32(4):513–520  
role of pharmacists in reducing suicide risk, 32(6):763–767  
treatment initiation for opioid use disorder, 32(5):724–731
- Mental health services**, Canadian HIV care settings as PCMHs, 32(2):158–167
- Mental status and dementia tests**, disease-modifying treatments for Alzheimer disease, 32(6):931–940
- Meta-analysis**, ruling out community-acquired pneumonia, 32(2):234–247
- Metabolic diseases**, monitoring the metabolic effects of antipsychotics, 32(3):418–423
- Metformin**, diabetes screening and treatment during office visits, 32(2):209–217
- Military personnel** veterans' reporting of cardiovascular disease, 32(1):50–57
- Minority health**  
hair care barrier to physical activity, 32(6):944–947  
opioid management in the safety-net, 32(3):375–382  
walk-in psychiatric model for integrated care, 32(4):481–489
- Morphine milligram equivalency (MME)**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Motivation**  
colorectal cancer screening in a FQHC, 32(2):180–190  
parent participation in child health research, 32(5):685–694
- Multimorbidity**, advance care planning conversations, 32(6):835–846
- Multiple chronic conditions**, advance care planning conversations, 32(6):835–846
- Multivariate analysis**, housing insecurity and access to care, 32(4):521–530
- Narcotics**, opioid prescribing and the pain-depression dyad, 32(4):614–618
- Nebraska**, sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Needs assessment**  
establishing a PBRN stakeholder engagement infrastructure, 32(5):695–704  
team-based breastfeeding support, 32(6):818–826
- Netherlands**  
knee pain and osteoarthritis among overweight women, 32(4):575–584  
natural history of back pain in older adults, 32(6):781–789
- New Mexico**, implementation of patient-team partnership, 32(4):490–504



- Newborns**, team-based breastfeeding support, 32(6):818–826
- Nicotine chewing gum**, smoking reduction and, 32(4):567–574
- No-show patients**, alternative payment impact on visits and scheduling, 32(4):539–549
- Northwestern United States**, treatment initiation for opioid use disorder, 32(5):724–731
- Nurse practitioners**  
 provider perspectives of patient experiences in imaging, 32(3):392–397  
 retail clinic use among Medicare beneficiaries, 32(4):531–538
- Nutrition surveys**  
 blood donation rates in the United States 1999–2016, 32(5):746–748  
 dietary inflammatory index and diabetes severity, 32(6):801–806
- Obesity**  
 hair care barrier to physical activity, 32(6):944–947  
 knee pain and osteoarthritis among overweight women, 32(4):575–584  
 lifestyle intervention programs for diabetes, 32(4):596–600  
 patient BMI knowledge in a rural population, 32(3):413–417  
 practice characteristics and patient self-management support, 32(3):329–340  
 translating self-management support into primary care, 32(3):341–352
- Occult blood**  
 colorectal cancer screening in a FQHC, 32(2):180–190  
 Latino-based colorectal cancer screening messaging, 32(3):307–317
- Occupational burnout**, medication management improves providers' work-life, 32(4):462–473
- Oklahoma**, adopting advance directives in primary care, 32(2):168–179
- Ondansetron**, high-risk medication without documented contraception, 32(4):474–480
- Ontario**, end of life services for the homeless, 32(6):847–857
- Opioid prescription**  
 intervention in chronic noncancer pain, 32(4):559–566  
 pain-depression dyad and, 32(4):614–618  
 regulation on, in an urban safety-net, 32(3):362–374
- Opioid-related disorders**  
 opioid prescribing intervention in chronic noncancer pain, 32(4):559–566  
 treatment initiation for opioid use disorder, 32(5):724–731
- Opioids**  
 early imaging for back pain in older adults, 32(6):773–780  
 opioid management in the safety-net, 32(3):375–382
- systems-based improvements to opioid management, 32(5):715–723
- Oregon**  
 alternative payment impact on visits and scheduling, 32(4):539–549  
 social risk stratification in primary care, 32(4):585–595
- Organizational innovation**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Organizations**, establishing a PBRN stakeholder engagement infrastructure, 32(5):695–704
- Osteoarthritis**, knee pain and osteoarthritis among overweight women, 32(4):575–584
- Outcomes assessment**  
 Colorado asthma toolkit implementation, 32(1):37–49  
 housing insecurity and access to care, 32(4):521–530  
 patient-centered research priorities, 32(5):674–684
- Outcomes research**, early imaging for back pain in older adults, 32(6):773–780
- Outpatients**  
 emergency department high utilizers, 32(2):264–268  
 follow-up and hospital readmissions, 32(3):353–361  
 ruling out community-acquired pneumonia, 32(2):234–247
- Overweight**, lifestyle intervention programs for diabetes, 32(4):596–600
- Ownership**, quality improvement measures by type of practice, 32(3):398–407
- Pain**  
 knee pain and osteoarthritis among overweight women, 32(4):575–584  
 opioid prescribing and the pain-depression dyad, 32(4):614–618
- Pain management**, early imaging for back pain in older adults, 32(6):773–780
- Palliative care**  
 advance care planning billing code use, 32(6):827–834  
 advance care planning conversations, 32(6):835–846  
 for the homeless, 32(6):858–867
- Parents**  
 lifestyle intervention programs for diabetes, 32(4):596–600  
 parent participation in child health research, 32(5):685–694  
 patient-centered research priorities, 32(5):674–684
- Partnership practice**, implementation of patient-team partnership, 32(4):490–504
- Patient activation**, text messages for HIV testing in primary care, 32(3):408–412
- Patient care**  
 patient-centered research priorities, 32(5):674–684
- patients frequently readmitted versus other admitted patients, 32(1):58–64
- physician characteristics and low value care spending, 32(2):218–225
- Patient care team**  
 depression treatment for Spanish and English speakers, 32(6):904–912  
 implementation of patient-team partnership, 32(4):490–504  
 practice characteristics and patient self-management support, 32(3):329–340
- Patient-centered care**  
 adopting advance directives in primary care, 32(2):168–179  
 advance care planning conversations, 32(6):835–846  
 burnout, clinic capacity, and social needs, 32(1):69–78  
 Canadian HIV care settings as PC-MHs, 32(2):158–167  
 hair care barrier to physical activity, 32(6):944–947  
 implementation of patient-team partnership, 32(4):490–504  
 patient-defined visit priorities: psychosocial versus medical, 32(4):513–520  
 practice characteristics and patient self-management support, 32(3):329–340  
 research priorities, 32(5):674–684  
 retail clinic use among Medicare beneficiaries, 32(4):531–538  
 socioeconomic and hospitalization of veterans, 32(6):890–903  
 translating self-management support into primary care, 32(3):341–352  
 visit entropy associated with diabetic control outcome, 32(5):739–745
- Patient compliance**, depression treatment for Spanish and English speakers, 32(6):904–912
- Patient discharge**  
 family medicine transitional care service line, 32(4):619–627  
 patients frequently readmitted versus other admitted patients, 32(1):58–64
- Patient health questionnaire**, depression treatment for Spanish and English speakers, 32(6):904–912
- Patient outcome assessment**, patient-centered research priorities, 32(5):674–684
- Patient participation**  
 developing patient and community advisory groups, 32(5):663–673  
 portal-based advance care planning outreach, 32(6):925–930  
 process for setting research priorities, 32(5):655–662
- Patient positioning**, sources of error in blood pressure measurement, 32(5):732–738
- Patient preference**, adopting advance directives in primary care, 32(2):168–179
- Patient Protection and Affordable Care Act (ACA)**

- alternative payment impact on visits and scheduling, 32(4):539–549  
preexisting conditions and, 32(6): 883–889
- Patient readmission**  
family medicine transitional care service line, 32(4):619–627  
outpatient follow-up and hospital readmissions, 32(3):353–361  
patients frequently readmitted versus other admitted patients, 32(1): 58–64
- Patient safety**  
laboratory safety toolkit in primary care, 32(2):136–145  
opioid prescribing intervention in chronic noncancer pain, 32(4): 559–566
- Patient satisfaction**, patient characteristics and making requests, 32(2): 201–208
- Patient transfer**, family medicine transitional care service line, 32(4): 619–627
- Pay for performance**, declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772
- Pediatric obesity**, lifestyle intervention programs for diabetes, 32(4): 596–600
- Personal satisfaction**, eConsult service implementation in a new health region, 32(2):146–157
- Personality**, patient characteristics and making requests, 32(2):201–208
- Pharmacists**  
community, role in reducing suicide risk, 32(6):763–767  
medication management improves providers' work-life, 32(4):462–473
- Pharmacoepidemiology**, high-risk medication without documented contraception, 32(4):474–480
- Pharmacology**, drug-induced hyperpigmentation, 32(4):628–638
- Physical fitness**, counseling on fitness and strengthening, 32(1):103–107
- Physician assistants**, provider perspectives of patient experiences in imaging, 32(3):392–397
- Physician-patient relations**  
advance care planning conversations, 32(6):835–846  
discussions about driving safety, 32(4):607–613  
men's depression treatment and, 32(1):13–19  
patient-defined visit priorities: psychosocial versus medical, 32(4): 513–520  
portal-based advance care planning outreach, 32(6):925–930
- Pneumonia**, community-acquired, ruling out, 32(2):234–247
- Point-of-care systems**, prediabetes at an academic practice, 32(4):505–512
- Point-of-care testing**, rapid point of care test for influenza, 32(2): 226–233
- Politics**, AAFP values and policies, 32(6):948–950
- Polymerase chain reaction**, rapid point of care test for influenza, 32(2):226–233
- Polypharmacy**, optimal prescribing in patients with dementia, 32(3): 383–391
- Population health**  
accountable care savings in deprived communities, 32(6):913–922  
blood donation rates in the United States 1999–2016, 32(5):746–748  
socioeconomics and hospitalization of veterans, 32(6):890–903
- Potentially inappropriate medications**, optimal prescribing in patients with dementia, 32(3): 383–391
- Practice-based research**  
developing patient and community advisory groups, 32(5):663–673  
establishing a PBRN stakeholder engagement infrastructure, 32(5): 695–704  
parent participation in child health research, 32(5):685–694  
patient-centered research priorities, 32(5):674–684  
process for setting research priorities, 32(5):655–662  
team-based breastfeeding support, 32(6):818–826
- Practice management**  
drug-induced hyperpigmentation, 32(4):628–638  
team-based breastfeeding support, 32(6):818–826
- Preconception care**, high-risk medication without documented contraception, 32(4):474–480
- Prediabetes**  
diabetes screening and treatment during office visits, 32(2):209–217  
prediabetes at an academic practice, 32(4):505–512  
status, awareness of, subsequent health behavior and, 32(1):20–27
- Prediabetic state**, lifestyle intervention programs for diabetes, 32(4): 596–600
- Preexisting condition coverage**, Affordable Care Act and, 32(6): 883–889
- Pregnancy**, high-risk medication without documented contraception, 32(4):474–480
- Prehypertension**, sources of error in blood pressure measurement, 32(5):732–738
- Premature mortality**, counseling on fitness and strengthening, 32(1): 103–107
- Prenatal care**, high-risk medication without documented contraception, 32(4):474–480
- Prescriptions**, opioid prescribing and the pain-depression dyad, 32(4): 614–618
- Prevalence**  
childhood atopic dermatitis in primary care, 32(2):191–200  
diabetes screening and treatment during office visits, 32(2):209–217  
psychotherapeutic medication misuse screening, 32(2):272–278  
veterans' reporting of cardiovascular disease, 32(1):50–57
- Preventive medicine**  
suboptimal statin use for secondary prevention, 32(6):807–817  
team-based breastfeeding support, 32(6):818–826
- Primary care physicians**  
accountable care savings in deprived communities, 32(6):913–922  
advance care planning conversations, 32(6):835–846  
basal insulin in primary care, 32(3): 431–447  
family medicine transitional care service line, 32(4):619–627  
hepatitis C and expanding the role of primary care, 32(3):428–430  
low hemoglobin A1c and hemoglobinopathy, 32(6):923–924  
patient characteristics and making requests, 32(2):201–208  
physician characteristics and low value care spending, 32(2):218–225  
providing e-visits, 32(6):868–875  
retail clinic use among Medicare beneficiaries, 32(4):531–538
- Primary health care**  
accountable care savings in deprived communities, 32(6):913–922  
adopting advance directives in, 32(2): 168–179  
alternative payment impact on visits and scheduling, 32(4):539–549  
basal insulin in, 32(3):431–447  
burnout, clinic capacity, and social needs, 32(1):69–78  
Canadian HIV care settings as PC-MHs, 32(2):158–167  
childhood atopic dermatitis in, 32(2): 191–200  
Colorado asthma toolkit implementation, 32(1):37–49  
colorectal cancer screening in a FQHC, 32(2):180–190  
counseling on fitness and strengthening, 32(1):103–107  
depression treatment for Spanish and English speakers, 32(6):904–912  
developing patient and community advisory groups, 32(5):663–673  
disease-modifying treatments for Alzheimer disease, 32(6):931–940  
eConsult service implementation in a new health region, 32(2):146–157  
emergency department high utilizers, 32(2):264–268  
enhanced shared decision making for depression, 32(1):6–9  
hair care barrier to physical activity, 32(6):944–947



- high-risk medication without documented contraception, 32(4):474–480
- housing insecurity and access to care, 32(4):521–530
- implementation of patient-team partnership, 32(4):490–504
- implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- knee pain and osteoarthritis among overweight women, 32(4):575–584
- laboratory safety toolkit in, 32(2):136–145
- LARC provision by family physicians, 32(1):10–12
- low hemoglobin A1c and hemoglobinopathy, 32(6):923–924
- medication management improves providers' work-life, 32(4):462–473
- monitoring the metabolic effects of antipsychotics, 32(3):418–423
- natural history of back pain in older adults, 32(6):781–789
- new allopathic medical schools train fewer family physicians, 32(5):653–654
- opioid management in the safety-net, 32(3):375–382
- opioid prescribing and the pain-depression dyad, 32(4):614–618
- opioid prescribing intervention in chronic noncancer pain, 32(4):559–566
- optimal prescribing in patients with dementia, 32(3):383–391
- parent participation in child health research, 32(5):685–694
- patient BMI knowledge in a rural population, 32(3):413–417
- patient-centered research priorities, 32(5):674–684
- patient characteristics and making requests, 32(2):201–208
- patient-defined visit priorities: psychosocial versus medical, 32(4):513–520
- physicians trained in teaching health centers, 32(2):134–135
- portal-based advance care planning outreach, 32(6):925–930
- practice characteristics and patient self-management support, 32(3):329–340
- prediabetes at an academic practice, 32(4):505–512
- process for setting research priorities, 32(5):655–662
- provider perspectives of patient experiences in imaging, 32(3):392–397
- psychotherapeutic medication misuse screening, 32(2):272–278
- quality improvement measures by type of practice, 32(3):398–407
- quality of the after-visit summary (AVS), 32(1):65–68
- reporting specialty status in research, 32(6):941–943
- retail clinic use among Medicare beneficiaries, 32(4):531–538
- role of pharmacists in reducing suicide risk, 32(6):763–767
- social and behavioral determinants of health screening, 32(3):297–306
- social risk stratification in primary care, 32(4):585–595
- socioeconomics and hospitalization of veterans, 32(6):890–903
- sugar-sweetened beverage intake, rural populations, 32(4):601–606
- systems-based improvements to opioid management, 32(5):715–723
- team-based breastfeeding support, 32(6):818–826
- text messages for HIV testing in, 32(3):408–412
- training vulnerable patients to use a patient portal, 32(2):248–258
- translating self-management support into primary care, 32(3):341–352
- treatment initiation for opioid use disorder, 32(5):724–731
- visit entropy associated with diabetic control outcome, 32(5):739–745
- walk-in psychiatric model for integrated care, 32(4):481–489
- yoga mentions in the electronic health record, 32(6):790–800
- Process measures**, quality improvement measures by type of practice, 32(3):398–407
- Progestins**, broken Nexplanon® rods, 32(2):269–271
- Proof of concept study**, text messages for HIV testing in primary care, 32(3):408–412
- Prospective studies**  
natural history of back pain in older adults, 32(6):781–789  
ruling out community-acquired pneumonia, 32(2):234–247
- Psychiatry**  
monitoring the metabolic effects of antipsychotics, 32(3):418–423  
walk-in psychiatric model for integrated care, 32(4):481–489
- Psychometrics**, validating the ABFM certification examination, 32(6):876–882
- Psychotherapy**  
depression treatment for Spanish and English speakers, 32(6):904–912  
doctor-patient relationships and men's depression treatment, 32(1):13–19
- Psychotropic drugs**, psychotherapeutic medication misuse screening, 32(2):272–278
- Public health**  
emergency department high utilizers, 32(2):264–268  
monitoring the metabolic effects of antipsychotics, 32(3):418–423  
opioid prescribing and the pain-depression dyad, 32(4):614–618  
role of pharmacists in reducing suicide risk, 32(6):763–767  
social and behavioral determinants of health screening, 32(3):297–306
- Qualitative research**  
depression treatment for Spanish and English speakers, 32(6):904–912  
laboratory safety toolkit in primary care, 32(2):136–145  
optimal prescribing in patients with dementia, 32(3):383–391  
palliative health care for the homeless, 32(6):858–867  
patient-centered research priorities, 32(5):674–684  
provider perspectives of patient experiences in imaging, 32(3):392–397  
social and behavioral determinants of health screening, 32(3):297–306  
social risk stratification in primary care, 32(4):585–595  
systems-based improvements to opioid management, 32(5):715–723
- Quality improvement**  
after-visit summaries, 32(1):65–68  
family medicine transitional care service line, 32(4):619–627  
implementation of patient-team partnership, 32(4):490–504  
implementing EvidenceNow to improve cardiovascular care, 32(5):705–714  
laboratory safety toolkit in primary care, 32(2):136–145  
measures, by type of practice, 32(3):398–407  
opioid prescribing intervention in chronic noncancer pain, 32(4):559–566  
portal-based advance care planning outreach, 32(6):925–930  
practice characteristics and patient self-management support, 32(3):329–340  
systems-based improvements to opioid management, 32(5):715–723
- Quality of life**  
adopting advance directives in primary care, 32(2):168–179  
natural history of back pain in older adults, 32(6):781–789
- Racism**, conversations on Twitter about diversity in medicine, 32(1):28–36
- Radiology**, early imaging for back pain in older adults, 32(6):773–780
- Referral and consultation**  
depression treatment for Spanish and English speakers, 32(6):904–912  
eConsult service implementation in a new health region, 32(2):146–157  
patient characteristics and making requests, 32(2):201–208
- Registries**, implementation of patient-team partnership, 32(4):490–504
- Reproductive health**, high-risk medication without documented contraception, 32(4):474–480
- Rescheduling**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Research report**, reporting specialty status in research, 32(6):941–943

**Residency**

declining presence of family physicians in hospital-based care, 32(6):768–770, 32(6):771–772  
physicians trained in teaching health centers, 32(2):134–135

**Resource allocation**, implementing

EvidenceNow to improve cardiovascular care, 32(5):705–714

**Retail clinics**, use among Medicare

beneficiaries, 32(4):531–538

**Retrospective studies**

opioid prescribing intervention in chronic noncancer pain, 32(4):559–566  
outpatient follow-up and hospital readmissions, 32(3):353–361  
visit entropy associated with diabetic control outcome, 32(5):739–745

**Rheumatology**, knee pain and osteoarthritis among overweight

women, 32(4):575–584

**Risk adjustment**, social risk stratification in primary care, 32(4):585–595**Risk assessment**

dietary inflammatory index and diabetes severity, 32(6):801–806  
discussions about driving safety, 32(4):607–613  
optimal prescribing in patients with dementia, 32(3):383–391

**Risk factors**, patient BMI knowledge in a rural population, 32(3):413–417**Risk-taking**, housing insecurity and access to care, 32(4):521–530**Rural health**

emergency medicine staffing, 32(3):292–294  
endoscopic care by family physicians, 32(4):460–461  
family physicians in emergent and urgent care, 32(3):295–296  
patient BMI knowledge in a rural population, 32(3):413–417  
veterans perceptions of access to primary care, 32(5):749–751

**Rural population**

patient BMI knowledge in, 32(3):413–417  
sugar-sweetened beverage intake, 32(4):601–606

**Screening**

colorectal cancer  
endoscopic services, 32(4):454–456  
in a FQHC, 32(2):180–190  
Latino-based messaging, 32(3):307–317  
health, social and behavioral determinants of, 32(3):297–306

**Seborrheic keratosis**, benign skin lesions and TADA, 32(1):96–102**Secondary prevention**, suboptimal statin use for secondary prevention, 32(6):807–817**Self-assessment**, physician opinions about ABFM self-assessment modules, 32(1):79–88**Self care**

practice characteristics and patient self-management support, 32(3):329–340

translating self-management support into primary care, 32(3):341–352

**Self-management**, support, translating into primary care, 32(3):341–352**Self report**

acupuncture training and physician burnout, 32(2):259–263  
sugar-sweetened beverage intake, rural populations, 32(4):601–606  
veterans perceptions of access to primary care, 32(5):749–751

**Sensitivity and specificity**, disease-modifying treatments for Alzheimer disease, 32(6):931–940**Sexism**, conversations on Twitter about diversity in medicine, 32(1):28–36**Shared decision making**, provider perspectives of patient experiences in imaging, 32(3):392–397**Sigmoidoscopy**, colorectal cancer screening in a FQHC, 32(2):180–190**Skin cancer**, benign skin lesions and TADA, 32(1):96–102**Skin care**, childhood atopic dermatitis in primary care, 32(2):191–200**Smoking cessation**, electronic cigarettes and, 32(4):567–574**Smoking reduction**, electronic cigarettes and, 32(4):567–574**Social determinants of health**

hair care barrier to physical activity, 32(6):944–947  
housing insecurity and access to care, 32(4):521–530  
patient-defined visit priorities: psychosocial versus medical, 32(4):513–520  
social and behavioral determinants of health screening, 32(3):297–306  
socioeconomics and hospitalization of veterans, 32(6):890–903

**Social media**, conversations on Twitter about diversity in medicine, 32(1):28–36**Social work**, end of life services for the homeless, 32(6):847–857**Socioeconomic status**, hospitalization of veterans and, 32(6):890–903**Spain**, drug-induced hyperpigmentation, 32(4):628–638**Spanish-speaking patients**

depression treatment for, 32(6):904–912  
Hispanic health story from Texas, 32(1):115–117

**Special communications**

Does My Patient with Multiple Comorbidities Have Heart Failure with Preserved Ejection Fraction, and Does it Matter?, 32(3):424–427  
Exercise Is Medicine: Primary Care Counseling on Aerobic Fitness and Muscle Strengthening, 32(1):103–107

The Potential Emergence of Disease-Modifying Treatments for

Alzheimer Disease: The Role of Primary Care in Managing the Patient Journey, 32(6):931–940

**Specialization**, reporting specialty status in research, 32(6):941–943**Standard of care**

depression treatment for Spanish and English speakers, 32(6):904–912  
Hispanic health story from Texas, 32(1):115–117

**Statins**, suboptimal use, for secondary prevention, 32(6):807–817**Street drugs**, identifying substance use using frequency questions, 32(4):550–558**Stroke volume**, heart failure with preserved ejection fraction, 32(3):424–427**Students**, new allopathic medical schools train fewer family physicians, 32(5):653–654**Substance abuse detection**, psychotherapeutic medication misuse screening, 32(2):272–278**Substance-related disorders**

identifying substance use using frequency questions, 32(4):550–558  
opioid management in the safety-net, 32(3):375–382

palliative health care for the homeless, 32(6):858–867

**Sugars**, sugar-sweetened beverage intake, rural populations, 32(4):601–606**Suicidal ideation**, role of pharmacists in reducing suicide risk, 32(6):763–767**Suicide**, role of pharmacists in reducing risk, 32(6):763–767**Surveillance systems**, housing insecurity and access to care, 32(4):521–530**Surveys and questionnaires**

ABFM's data collection method for tracking specialty, 32(1):89–95  
accountable care savings in deprived communities, 32(6):913–922  
doctor-patient relationships and men's depression treatment, 32(1):13–19  
end of life services for the homeless, 32(6):847–857  
family physicians providing e-visits, 32(6):868–875  
hair care barrier to physical activity, 32(6):944–947  
identifying substance use using frequency questions, 32(4):550–558  
implementation of patient-team partnership, 32(4):490–504  
lifestyle intervention programs for diabetes, 32(4):596–600  
monitoring the metabolic effects of antipsychotics, 32(3):418–423  
natural history of back pain in older adults, 32(6):781–789  
patient BMI knowledge in a rural population, 32(3):413–417  
patient-centered research priorities, 32(5):674–684  
physician opinions about ABFM self-assessment modules, 32(1):79–88

- prediabetes at an academic practice, 32(4):505–512
- social risk stratification in primary care, 32(4):585–595
- suboptimal statin use for secondary prevention, 32(6):807–817
- team-based breastfeeding support, 32(6):818–826
- veterans' reporting of cardiovascular disease, 32(1):50–57
- Sustained virologic response**, hepatitis C and expanding the role of primary care, 32(3):428–430
- Sweetening agents**, sugar-sweetened beverage intake, rural populations, 32(4):601–606
- Systematic review**, ruling out community-acquired pneumonia, 32(2):234–247
- Teaching health centers**, 32(2):130–133, 32(2):134–135
- Teaching hospitals** advance care planning billing code use, 32(6):827–834
- Telemedicine**
  - family physicians providing e-visits, 32(6):868–875
  - text messages for HIV testing in primary care, 32(3):408–412
  - training vulnerable patients to use a patient portal, 32(2):248–258
- Telephone**, visit entropy associated with diabetic control outcome, 32(5):739–745
- Teratogens**, high-risk medication without documented contraception, 32(4):474–480
- Terminal care**
  - advance care planning conversations, 32(6):835–846
  - end of life services for the homeless, 32(6):847–857
  - palliative health care for the homeless, 32(6):858–867
- Tertiary care centers**, advance care planning billing code use, 32(6):827–834
- Texas**, Hispanic health story from, 32(1):115–117
- Text messaging**
  - for colorectal cancer screening, 32(3):307–317, 32(3):318–328
  - for HIV testing in primary care, 32(3):408–412
- Tobacco**, identifying substance use using frequency questions, 32(4):550–558
- Tobacco products**, electronic cigarettes and smoking reduction, 32(4):567–574
- Tramadol**, regulation on opioid prescribing in an urban safety-net, 32(3):362–374
- Transitional care**, family medicine transitional care service line, 32(4):619–627
- Triage amalgamated dermoscopic algorithm (TADA)**, and benign skin lesions, 32(1):96–102
- Twitter**, conversations about diversity in medicine, 32(1):28–36
- Type 2 diabetes mellitus**
  - basal insulin in primary care, 32(3):431–447
  - lifestyle intervention programs for diabetes, 32(4):596–600
  - practice characteristics and patient self-management support, 32(3):329–340
  - prediabetes at an academic practice, 32(4):505–512
  - translating self-management support into primary care, 32(3):341–352
  - visit entropy associated with diabetic control outcome, 32(5):739–745
- Underserved populations**, walk-in psychiatric model for integrated care, 32(4):481–489
- United States Agency for Health care Research and Quality**, laboratory safety toolkit in primary care, 32(2):136–145
- United States Department of Veterans Affairs**
  - socioeconomics and hospitalization of veterans, 32(6):890–903
  - veterans perceptions of access to primary care, 32(5):749–751
- Urban areas**, urgent care, family physicians' contributions, 32(3):295–296
- Urgent care**, urban, family physicians' contributions, 32(3):295–296
- Validity**, ABFM certification examination, 32(6):876–882
- Veterans**
  - hospitalization of, socioeconomics and, 32(6):890–903
  - perceptions of access to primary care, 32(5):749–751
  - reporting of cardiovascular disease, 32(1):50–57
- Violence**, values and policies of AAFP, 32(6):948–950
- Virginia**, implementing EvidenceNow to improve cardiovascular care, 32(5):705–714
- Vulnerable populations**
  - colorectal cancer screening in a FQHC, 32(2):180–190
  - monitoring the metabolic effects of antipsychotics, 32(3):418–423
  - opioid management in the safety-net, 32(3):375–382
- Washington**, socioeconomics and hospitalization of veterans, 32(6):890–903
- Weight gain**, basal insulin in primary care, 32(3):431–447
- Women's health**
  - hair care barrier to physical activity, 32(6):944–947
  - LARC provision by family physicians, 32(1):10–12
  - team-based breastfeeding support, 32(6):818–826
- Workflow**
  - portal-based advance care planning outreach, 32(6):925–930
  - social and behavioral determinants of health screening, 32(3):297–306
  - social risk stratification in primary care, 32(4):585–595
- Workload**, medication management improves providers' work-life, 32(4):462–473
- Yoga**, mentions in the electronic health record, 32(6):790–800