

BRIEF REPORT

Practice Patterns of Family Physicians With and Without Addiction Medicine Board Certification

Sebastian T. Tong, MD, MPH, Zachary J. Morgan, MS,
Andrew W. Bazemore, MD, MPH, Aimee R. Eden, PhD, MPH,
Ruchi M. Fitzgerald, MD, and Lars E. Peterson, MD, PhD

Background: The American Board of Medical Specialties recognized addiction medicine (ADM) as a subspecialty in 2016, which was timely given the recent rise in substance use disorder (SUD). The impact of this dual board opportunity on Family Medicine has not been described. Our study enumerates and characterizes physicians dually certified in Family Medicine and ADM.

Methods: We linked American Board of Medical Specialties data from March 2020 on physicians dually boarded in Family Medicine and ADM to responses on demographic and scope of practice questions in the American Board of Family Medicine (ABFM) National Graduate Survey and Family Medicine Certification Examination Registration Questionnaire.

Results: Of current ABFM Diplomates, 0.53% (492/93,269) are also boarded in ADM. Based on survey responses from a subset of dually certified physicians, those who are dually certified are more likely to practice in federally qualified health centers and to hold a faculty position. Dually certified physicians are more likely to provide HIV/AIDS and hepatitis C management and are as likely as non-dually certified physicians to provide newborn care, obstetric deliveries, inpatient adult medicine care, and intensive care.

Discussion: While only a small proportion of family physicians carry dual ADM board certification, those that do disproportionately serve vulnerable populations while retaining broad scope of care. Further work is needed to examine whether SUD treatment access could be addressed by implementing models that support dually certified physicians in consultative and educational efforts that would amplify their impact across the primary care workforce. (J Am Board Fam Med 2021;34:814–819.)

Keywords: Addiction Medicine, Attitude of Health Personnel, Certification, Family Medicine, Family Physicians, Outcome Measures, Primary Health Care, Referral and Consultation, Scope of Practice, Social Determinants of Health, Social Support, Socioeconomic Factors, Specialty Boards, Substance-Related Disorders, Surveys and Questionnaires, Workforce

Introduction

Drug overdose deaths in the United States have continued to increase, after a brief decline in 2018,

with a rise of 4.8% in 2019 over the past year to 70,980 deaths according to provisional data.¹ An estimated 20.4 million individuals aged 12 or older had a substance use disorder (SUD) in 2019, but only 12.2% of those who needed SUD treatment received it.² Early data suggest that SUD and drug overdose death rates have increased substantially while access to treatment has declined during the COVID-19 pandemic.^{3–5}

This article was externally peer reviewed.
Submitted 31 August 2020; revised 15 December 2020; accepted 16 December 2020.

From the Agency for Healthcare Research and Quality, Rockville, MD (STT); American Board of Family Medicine, Lexington, KY (ZJM, AWB, ARE, LEP); Departments of Family Medicine, and Psychiatry and Behavioral Sciences, Rush University, Chicago, IL (RMF); Department of Family and Community Medicine, College of Medicine, University of Kentucky, Lexington, KY (LEP).

Funding: Dr. Tong was supported by the American Board of Family Medicine Foundation as a Visiting Scholar.

Conflict of interest: Dr. Tong is employed by the Agency for Healthcare Research and Quality but contributed to this article in his personal capacity. The opinions expressed are the author's own and do not reflect the view of the Agency for Healthcare Research and Quality, the Department of

Health and Human Services, or the Federal government. Drs. Bazemore, Eden and Peterson, and Mr. Morgan are employed by the American Board of Family Medicine.

Corresponding author: Sebastian Tong, MD, MPH, Agency for Healthcare Research and Quality, 5600 Fishers Lane, Mail Stop 06E53A, Rockville, MD, 20857, Phone: 301-427-1627 (E-mail: sebastian.tong@ahrq.hhs.gov).

While evidence suggests that primary care physicians can successfully treat SUD,^{6,7} a recent study at a single institution reported that only 20% felt prepared to care for patients with SUD.⁸ A prior national survey in 2000 reported that less than 20% of primary care physicians felt prepared to identify SUD, and more than 50% of patients with SUD felt that their primary care physician did not address their SUD.⁹ Furthermore, only 28.6% of family medicine residencies reported having a dedicated addiction medicine (ADM) curricula with a lack of faculty expertise listed as the top reason why many programs did not offer training in ADM.¹⁰ Studies have also shown that few family physicians feel prepared to prescribe or are currently prescribing buprenorphine, which is the predominant medication used in office-based settings for opioid use disorder.¹¹⁻¹⁴

With a goal of increasing the workforce to address SUD, ADM became an American Board of Medical Specialties (ABMS)-recognized subspecialty in 2016 under the American Board of Preventive Medicine.^{15,16} Any physician with an ABMS-recognized primary board certification is eligible to apply with appropriate fellowship training or, before 2025, sufficient addiction-focused practice experience.^{17,18} However, the current need for SUD treatment services far surpasses the supply of ADM specialists.² Physicians who are dually certified in a primary care specialty and ADM could play an important role in educating other primary care physicians about ADM and serving as a consultant for complicated patients.^{19,20}

Little is known about family physicians (FPs) who hold a secondary board certification in ADM. We sought to describe physicians' demographic, practice, and scope of care characteristics dually certified in ADM and family medicine.

Methods

We obtained a list of FPs who are certified in ADM as of March 2020 from the ABMS. Using their ABMS identification code, we linked these physicians to FPs in the American Board of Family Medicine (ABFM) database. We then linked physicians who were dually certified in family medicine and ADM to their responses in the 2016–2019 ABFM National Graduate Survey (NGS) and the 2017–2019 ABFM Family Medicine Certification Examination Registration Questionnaire (CERQ),

if available. The NGS is administered annually to all ABFM Diplomates who graduated from residency 3 years prior. The CERQ is completed by FPs seeking to continue their certification 3 to 4 months before the examination date as a required component of registration. Both surveys asked questions about practice characteristics and scope of care. Detailed information about the NGS and CERQ can be found elsewhere.^{21,22} Physician demographic data were obtained from ABFM administrative data.

After excluding physicians, not in direct patient care, we produced descriptive statistics and performed χ^2 tests and, where appropriate, Fisher exact tests for associations between ADM certification and personal and practice characteristics. SAS Version 9.4 (Cary, NC) was used for analyses. The American Academy of Family Physicians Institutional Review Board approved this study.

Results

Overall, 0.53% or 492 of 93,269 ABFM Diplomates are also certified in ADM. We were able to match NGS or CERQ responses for 186 of these physicians (37.8%). Of NGS respondents, from which there was a response rate ranging from 67–68%, 0.53% (47/8,863) are dually certified. Of CERQ respondents, from which there was a response rate of 100%, 0.56% (139/24,991) are dually certified.

FPs who are ADM board certified were more likely to practice in a federally qualified health center (for NGS: 28.9% ADM certified vs. 11.7% not ADM certified, $P=.003$; for CERQ: 8.6% vs. 5.9%, $P<.001$) and in principal practices with multiple specialties (for NGS: 40.0% vs. 23.3%, $P=.030$; for CERQ: 39.3% vs. 21.7%, $P<.001$). From the NGS, core/salaried faculty FPs were more likely to be certified in ADM (23.4% vs. 12.0%, $P=.004$), although this difference was not significant for CERQ respondents. No statistically significant differences were found in age, gender, degree type (MD vs. DO), training site, or race/ethnicity (Table 1).

In terms of scope of practice, FPs certified in ADM who responded to the NGS were more likely than those not certified to provide buprenorphine (93.6% vs. 10.1%, $P<.001$), pharmacologic management of HIV/AIDS (31.9% vs. 17.8%, $P=.012$) and pharmacologic management

Table 1. Personal and Practice Characteristics of Family Physicians With and Without Addiction Medicine Board Certification From 2017–2019 ABFM Continuing Certification Examination Registration Questionnaire and 2016–2019 National Graduate Survey

Characteristics	National Graduate Survey 2016–2019			Continuing Certification 2017–2019		
	ADM Certified n (%)	Not ADM Certified n (%)	<i>P</i> value*	ADM Certified n (%)	Not ADM Certified n (%)	<i>P</i> value*
Total	47	8,816		139	24,991	
Age, y						
Under 40	40 (85.1)	7,712 (87.5)	.144 [†]	11 (7.9)	2,255 (9.0)	.295
40–49	4 (8.5)	939 (10.7)		57 (41.0)	9,338 (37.4)	
50–59	3 (6.4)	147 (1.7)		50 (36.0)	8,084 (32.3)	
60 and over	0 (0.0)	18 (0.2)		21 (15.1)	5,313 (21.3)	
Sex						
Male	23 (48.9)	3,880 (44.0)	.498	81 (58.3)	14,288 (57.2)	.794
Female	24 (51.1)	4,936 (56.0)		58 (41.7)	10,702 (42.8)	
IMG						
Yes	8 (17.0)	2,960 (33.6)	.016	26 (18.7)	5,588 (22.4)	.299
No	39 (83.0)	5,856 (66.4)		113 (81.3)	19,378 (77.6)	
Race ^{††}						
Asian	–	–		14 (10.1)	3,744 (15.0)	.221
Black or African American	–	–		5 (3.6)	1,459 (5.8)	
White	–	–		110 (79.1)	17,985 (72.0)	
Other	–	–		10 (7.2)	1,803 (7.2)	
Ethnicity [‡]						
Hispanic or Latino	–	–		11 (7.9)	1,692 (6.8)	.593
Not Hispanic or Latino	–	–		128 (92.1)	23,299 (93.2)	
Faculty at medical school or residency [§]						
Yes, core/salaried	10 (24.4)	812 (12.1)	.004	14 (10.1)	2,312 (9.3)	.106
Yes, volunteer/clinical	13 (31.7)	1,354 (20.1)		38 (27.3)	5,106 (20.4)	
No	18 (43.9)	4,563 (67.8)		87 (62.6)	17,572 (70.3)	
Principal practice size						
Solo practice	1 (2.2)	263 (3.6)	.439	29 (22.7)	2,544 (11.9)	<.001
2–5 providers	18 (40.0)	2,622 (35.6)		46 (35.9)	7,146 (33.5)	
6–20 providers	20 (44.4)	2,795 (38.0)		33 (25.8)	6,541 (30.7)	
>20 providers	6 (13.3)	1,675 (22.8)		20 (15.6)	5,097 (23.9)	
Principal practice site						
Hospital/health system owned medical practice	9 (20.0)	2,856 (38.8)	.003	21 (16.4)	7,219 (33.8)	<.001
Independently owned	7 (15.6)	1,082 (14.7)		56 (43.8)	6,813 (31.9)	
Managed care/HMO	2 (4.4)	532 (7.2)		7 (5.5)	1,245 (5.8)	
Academic health center/faculty practice	8 (17.8)	813 (11.0)		6 (4.7)	1,558 (7.3)	
FQHC or look-alike	13 (28.9)	858 (11.7)		11 (8.6)	1,256 (5.9)	
Other	6 (13.3)	1,223 (16.6)		27 (21.1)	3,238 (15.2)	
Specialty mix of principal practice						
Family medicine only	16 (35.6)	3,213 (43.7)	.030	34 (38.2)	10,193 (51.5)	<.001
Primary care specialty mix	11 (24.4)	2,430 (33.0)		20 (22.5)	5,310 (26.8)	
Multiple specialties	18 (40.0)	1,713 (23.3)		35 (39.3)	4,292 (21.7)	

Continued

Table 1. Continued

Characteristics	National Graduate Survey 2016–2019			Continuing Certification 2017–2019		
	ADM Certified n (%)	Not ADM Certified n (%)	<i>P</i> value*	ADM Certified n (%)	Not ADM Certified n (%)	<i>P</i> value*
Rurality						
Urban	44 (95.7)	7,178 (84.6)	.038	122 (87.8)	20,924 (84.5)	.289
Rural	2 (4.3)	1,310 (15.4)		17 (12.2)	3,836 (15.5)	

Abbreviations: ABFM, American Board of Family Medicine; ADM, addiction medicine; HMO, health maintenance organization; FQHC, federally qualified health center, IMG, International Medical Graduate.

*Unless otherwise noted, χ^2 test was to calculate the *P* value.

†Fisher exact test was used to calculate this *P* value.

‡Race and ethnicity data were not available for National Graduate Survey respondents.

2016 National Graduate Survey responses were excluded because the question about faculty differed from the 2017–2019 National Graduate Survey.

of hepatitis C (48.9% vs. 19.3%, $P < .001$). For CERQ respondents, this difference was observed for buprenorphine provision and pharmacologic management of hepatitis C. For both surveys, there were no differences in the provision of obstetric deliveries and other hospital-based care (including newborn, pediatric, adult and intensive care). FPs certified in ADM were less likely to provide pediatric outpatient care (Table 2).

Discussion

Despite the ongoing and worsening substance use disorder crisis in the United States,⁴ FPs who are dually certified in ADM are few in number. Dually certified physicians were more likely to practice in academic settings or federally qualified health centers. In addition, dually certified physician respondents to the NGS survey were more likely to hold a faculty position, although this finding was not statistically significant for

Table 2. Scope of Practice of Family Physicians With and Without Addiction Medicine Board Certification From 2017–2019 ABFM Continuing Certification Examination Registration Questionnaire and 2016–2019 National Graduate Survey

Characteristics	National Graduate Survey 2016–2019			Continuing Certification 2017–2019		
	ADM Certified n (%)	Not ADM Certified n (%)	<i>P</i> value*	ADM Certified n (%)	Not ADM Certified n (%)	<i>P</i> value*
Prenatal care	†			18 (12.9)	2,805 (11.2)	.521
Delivering babies	7 (14.9)	1,194 (13.7)	.806	12 (8.6)	1,637 (6.6)	.323
Newborn hospital care	10 (21.3)	2,101 (24.0)	.665	14 (10.1)	3,169 (12.7)	.356
Pediatric hospital care (not newborn)	8 (17.0)	1,714 (19.6)	.661	8 (5.8)	2,286 (9.1)	.166
Pediatric outpatient care	28 (59.6)	6,704 (76.5)	.006	43 (30.9)	13,812 (55.3)	<.001
Adult inpatient medicine	19 (40.4)	3,445 (39.4)	.885	41 (29.5)	6,079 (24.3)	.157
Intensive care/ICU-CCU	8 (17.0)	1,953 (22.3)	.386	11 (7.9)	2,316 (9.3)	.583
Behavioral health care	44 (93.6)	7,693 (87.8)	.224	99 (71.2)	8,962 (35.9)	<.001
Integrative medicine	8 (17.0)	1,648 (18.8)	.754	14 (10.1)	1,132 (4.5)	.002
End of life care	20 (42.6)	5,354 (61.1)	.009	22 (15.8)	7,071 (28.3)	.001
Buprenorphine treatment	44 (93.6)	886 (10.1)	<.001	26 (83.9)	285 (5.7)	<.001 [‡]
Pharmacologic management of HIV/AIDS	15 (31.9)	1,554 (17.8)	.012	1 (3.2)	197 (3.9)	1.000 [‡]
Pharmacologic management of hepatitis C	23 (48.9)	1,688 (19.3)	<.001	6 (19.4)	183 (3.6)	<.009 [‡]

ABFM, American Board of Family Medicine; ADM, addiction medicine; HIV/AIDS, human immunodeficiency virus/acquired immuno deficiency syndrome; ICU-CCU, intensive care unit-coronary care unit.

*Unless otherwise noted, χ^2 test was to calculate the *P* value.

†Data on prenatal care provision was not available for the National Graduate Survey.

‡Fisher exact test was used to calculate this *P* value.

CERQ respondents. These findings suggest that dually certified physicians may disproportionately work in underserved settings and in a teaching capacity.

Many dually certified physicians have maintained similar scope of practice as non-dually certified physicians, including hospital-based care and obstetric care, which suggests that dually certified physicians are not leaving primary care to practice addiction medicine solely.²³ Furthermore, we found that dually certified physicians are more likely to provide hepatitis C pharmacologic treatment and that early career dually certified physicians are more likely to provide HIV/AIDS treatment. This mechanism could provide increased access to evidence-based treatments for HIV/AIDS and hepatitis C, which are found at higher rates in individuals with SUD, both through direct patient care and mentoring/educational initiatives.

The small number of dual-certified physicians makes it clear that the burden of treating SUD cannot solely rest on these physicians. Rather, dually certified physicians could potentially increase access to SUD treatment services by educating and mentoring fellow primary care physicians about SUD management. Recent successful models of this care include Project Echo,¹⁹ the Provider Clinical Support System,²⁴ and the Hub-and-Spoke model in Vermont.²⁰ Future studies could examine how to use these models to best expand the workforce treating addiction by using the limited supply of dually certified physicians to provide more mentoring and technical assistance to community primary care physicians who might otherwise be hesitant to provide SUD care in their practices.

Several limitations exist. First, although our overall sample size is large, the total number of FPs who are ADM certified included in our sample is small and, thus, potentially limited generalizability. Second, examining ABMS board certification in ADM may underestimate FPs with expertise in ADM since they do not include those certified by mechanisms that existed before ABMS recognition of ADM as a specialty or those with expertise without any certification. Third, a response bias could potentially exist for the NGS. This limitation does not exist for the CERQ since it is mandatory.

Conclusion

While few in number, FPs with board certification in ADM are more likely to work in underserved areas, teach capacity, and provide HIV/AIDS and hepatitis C care while maintaining an otherwise

similar scope of practice to non-dually certified FPs. Further work is needed to examine whether dually certified physicians play an important role in educating and mentoring other primary care physicians about SUD care and how dually certified physicians help address treatment gaps in SUD.

To see this article online, please go to: <http://jabfm.org/content/34/4/814.full>.

References

- Centers for Disease Control and Prevention. Provisional Drug Overdose Death Counts. Available from: <https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm>. Published 2020. Accessed August 4, 2020.
- Substance Abuse and Mental Health Services Administration. Key substance use and mental health indicators in the United States: Results from the 2018 National Survey on Drug Use and Health (HHS Publication No. PEP19-5068, NSDUH Series H-54). 2019. Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Services Administration.
- Overdose Detection Mapping Application Program. The Consequences of COVID-19 on the Overdose Epidemic: Overdoses are Increasing. Available from: <http://www.odmap.org/Content/docs/news/2020/ODMAP-Report-May-2020.pdf>. Published 2020. Accessed August 4, 2020.
- Becker WC, Fiellin DA. When epidemics collide: coronavirus disease 2019 (COVID-19) and the opioid crisis. *Ann Intern Med* 2020;173(1):59–60.
- Slavova S, Rock P, Bush HM, Quesinberry D, Walsh SL. Signal of increased opioid overdose during COVID-19 from emergency medical services data. *Drug Alcohol Depend* 2020;214:108176.
- Edelman EJ, Oldfield BJ, Tetrault JM. Office-based addiction treatment in primary care: approaches that work. *Med Clin* 2018;102(4):635–652.
- Korthuis PT, McCarty D, Weimer M, et al. Primary care-based models for the treatment of opioid use disorder: A scoping review. *Ann Intern Med* 2017;166(4):268–278.
- Wakeman SE, Pham-Kanter G, Donelan K. Attitudes, practices, and preparedness to care for patients with substance use disorder: results from a survey of general internists. *Subst Abuse* 2016;37(4):635–641.
- Survey Research Laboratory. Missed Opportunity: National Survey of Primary Care Physicians and Patients on Substance Abuse. Chicago, IL: University of Illinois at Chicago; 2000.
- Tong S, Sabo R, Aycock R, et al. Assessment of addiction medicine training in family medicine residency programs: a CERA study. *Fam Med* 2017;49(7):537.
- Peterson LE, Morgan ZJ, Borders TF. Practice predictors of buprenorphine prescribing by family physicians. *J Am Board Fam Med* 2020;33(1):118–123.

12. Peterson LE, Morgan ZJ, Eden AR. Early-Career and graduating physicians more likely to prescribe buprenorphine. *J Am Board Fam Med* 2020;33(1):7–8.
13. Abraham R, Wilkinson E, Jabbarpour Y, Petterson S, Bazemore A. Characteristics of office-based buprenorphine prescribers for Medicare patients. *J Am Board Fam Med* 2020;33(1):9–16.
14. Tong ST, Hochheimer CJ, Peterson LE, Krist AH. Buprenorphine provision by early career family physicians. *Ann Fam Med* 2018;16(5):443–446.
15. Nishiaoki M, McNally D, Bey RM, Arias AJ. New ACGME-accredited addiction medicine fellowship programs and their impact on the field. *Curr Treat Options Psychiatry* 2020;7(1):1–8.
16. Kunz K, Wiegand T. Addiction medicine: current status of certification, maintenance of certification, training, and practice. *J Med Toxicol* 2016;12(1):76–78.
17. American Board of Preventive Medicine. Addiction Medicine Requirements. Available from: <https://www.theabpm.org/become-certified/subspecialties/addiction-medicine/>. Published 2020. Accessed August 4, 2020.
18. American Board of Preventive Medicine. The American Board of Preventive Medicine and the American Society of Addiction Medicine Announce a Four-Year Extension of the Practice Pathway for Addiction Medicine. Available from: <https://www.theabpm.org/2020/11/30/the-american-board-of-preventive-medicine-and-the-american-society-of-addiction-medicine-announce-a-four-year-extension-of-the-practice-pathway-for-addiction-medicine/>. Published 2020. Accessed December 9, 2020.
19. Komaromy M, Duhigg D, Metcalf A, et al. Project ECHO (Extension for Community Healthcare Outcomes): A new model for educating primary care providers about treatment of substance use disorders. *Subst Abus* 2016;37(1):20–24.
20. Brooklyn JR, Sigmon SC. Vermont hub-and-spoke model of care for opioid use disorder: development, implementation, and impact. *J Addict Med* 2017;11(4):286.
21. Peterson L, Fang B, Phillips Jr R, Avant R, Puffer J. A certification board's tracking of their specialty: the American Board of Family Medicine's data collection strategy. *J Am Board Fam Med* 2019;32(1):89–95.
22. Weidner AKH, Chen FM, Peterson LE. Developing the National Family Medicine Graduate Survey. *J Grad Med Educ* 2017;9(5):570–573.
23. Louis JS, Eden AR, Morgan ZJ, Barreto TW, Peterson LE, Phillips RL. Maternity care and buprenorphine prescribing in new family physicians. *Ann Fam Med* 2020;18(2):156–158.
24. Egan JE, Casadonte P, Gartenmann T, et al. The Physician Clinical Support System-Buprenorphine (PCSS-B): a novel project to expand/improve buprenorphine treatment. *J Gen Intern Med* 2010;25(9):936–941.