

ORIGINAL RESEARCH

Impact of the COVID-19 Pandemic on Exercise Habits Among US Primary Care Patients

Natalie Wood Byrne, PharmD, Daniel J. Parente, MD, PhD, and
Nicole T. Yedlinsky, MD, CAQSM, FAAFP

Background: To explore how the COVID-19 pandemic has affected exercise habits, we hypothesized that participants' physical activity would have increased by at least 30 min/wk after the onset of the pandemic.

Methods: We distributed an anonymous survey to ambulatory patients at the Family Medicine Clinic, University of Kansas Medical Center to analyze changes in exercise habits and weight.

Results: Of the 500 adult patients surveyed, 382 were included. Results were stratified by demographics, including employment status before and during COVID-19. The median change in weekly exercise duration was 0.0 minutes, but the mean change was -25.7 minutes; total exercise duration decreased after the pandemic's onset (paired Wilcoxon signed rank test $P < .001$). More individuals reported participation in virtual group classes (6.3% before the pandemic vs 13.1% during the pandemic; McNemar's $P < .001$). Individuals with home exercise equipment before the pandemic were more likely to acquire more than were those who had none before (Chi square test $P < .005$). Overall, there is a significant trend in the direction of weight gain (Wilcoxon signed rank test $P < .001$).

Conclusions: Most participants decreased physical activity during the unprecedented COVID-19 pandemic, expanding our understanding of how exercise habits change during stressful life events. (J Am Board Fam Med 2022;35:295–309.)

Keywords: Behavioral Sciences, COVID-19, Cross-Sectional Studies, Disease, Management, Employment, Family Medicine, Health Behavior, Lifestyle, Pandemics, Primary Health Care, Retrospective Studies, Surveys and Questionnaires

Introduction

The overall health benefits of physical activity are well documented. Regular physical activity can improve sleep, upregulate the immune system, and reduce all-cause mortality.^{2,11,21,26} Current guidelines recommend that adults weekly perform at least 150

minutes of moderate-intensity aerobic activity, or 75 minutes vigorous-intensity aerobic activity, or an equivalent combination of moderate and vigorous intensity aerobic activity.²¹ The guidelines also recommend muscle-strengthening activities that involve all muscle groups at least 2 days per week.²¹ Since the first guidelines were released in 2008, however, only 26% of men and 19% of women reported enough activity to meet the aerobic and muscle strengthening recommendations put forth by these guidelines.²¹

Although exercise has health benefits, there are many perceived barriers to increasing exercise time. One barrier commonly identified in adults across multiple studies is lack of time.^{4,8,17,20} A 2017 systematic review found that individuals who reported lack of time were less likely to participate in physical activity.⁸ Greater total work hours and overtime work hours were also both negatively correlated with participation in physical activity.⁸

Despite many adults using their waking time working, few studies have evaluated how work

This article was externally peer reviewed.
Submitted 21 July 2021; revised 7 October 2021; accepted 12 October 2021.

Drs Daniel J. Parente and Nicole T. Yedlinsky contributed equally.

From School of Medicine, University of Kansas Medical Center, Kansas City, KS (NWB); Department of Family Medicine and Community Health, University of Kansas Medical Center, Kansas City, KS (DJP, NTY).

Funding: This work was supported in part by the University of Kansas Medical Center, who permitted the use of their REDCap electronic management data system for our data analysis.

Competing and conflicting interests: None declared.

Corresponding author: Nicole T. Yedlinsky, MD, CAQSM, FAAFP, Department of Family Medicine and Community Health, University of Kansas Medical Center, 3901 Rainbow Boulevard, MS 4010, Kansas City, KS 66160 (E-mail: nyedlinsky@kumc.edu).

affects time spent exercising. Data indicate, however, that leisure-time physical activity levels were suboptimal among all major US worker groups.⁵ Moreover, workers who report feeling fatigued after work are less likely to engage in leisure time physical activity.³ In workers younger than 50 years old, this decrease could be as much as 50 min/wk among workers reporting that they were “very tired” after work.³ This is especially problematic because increased occupational physical activity has been found to be paradoxically harmful to health.¹⁵

In 2020, a novel coronavirus (COVID-19) outbreak traveled worldwide, resulting in over 540,000 fatalities in the US alone.⁷ This outbreak triggered unprecedented changes in all aspects of life. Lawmakers in most states imposed “Stay at Home” or “Safer at Home” orders to mitigate virus spread. These orders varied among states and often included actions such as closing indoor dining and gyms, restricting business hours, and mask mandates. Some states, such as Kansas (the study location), had extreme restrictions that led to the closure of all businesses and activities unless deemed “essential.”¹¹ This led to a shift in working behavior, causing many US Americans to begin working from home.

We assessed how the unprecedented change in the US American workforce during the COVID-19 crisis affected leisure time physical activity. With almost all fifty states issuing a “Stay at Home” or “Shelter in Place” order as well as national social distancing practices, many workers found themselves spending a significant amount of time at home. All sporting events, and other various social events were canceled in the US during the spring of 2020 due to the COVID-19 outbreak. We evaluated the impact of the COVID-19 pandemic on (1) duration of exercise in min/wk, (2) setting of physical activity, (3) acquisition of home exercise equipment, and (4) weight change.

Methods

Participants

We surveyed English-speaking adults living in the US who attended a primary care appointment in the University of Kansas Medical Center Department of Family Medicine between September 2020 through November 2020.

Exclusion criteria included persons who did not primarily reside in the US between February 1 and the time of the survey as well as incarcerated individuals.

Survey Implementation

A 1-time, written, anonymous survey was distributed by front desk personnel to 500 patients (pre-specified) attending ambulatory appointments and collected by a nurse when the patient was brought to an examination room. Surveys inquired about employment status, time spent exercising, type of exercise activities, location of exercise activities, and purchase of at home exercise equipment (exact formatting shown in the Appendix materials). Social and demographic information was collected, including race, gender, marital status, education attainment, and whether there were other adults or children in the home. The evaluation of Leisure Time Physical Activity each week was modeled from the International Physical Activity Questionnaire¹² to calculate exercise in time in min/wk and compare with goals set forth by current guidelines. Study data were collected and managed using REDCap electronic data capture tools hosted by the University of Kansas Medical Center.^{18,19} Written surveys were manually entered into the REDCap electronic data management system for further analysis.

Statistical Methods

Data were exported from REDCap and analyzed using R v. 3.6.1. Descriptive statistics were compiled using standard methods. Change in continuous and ordinal variables that were not normally distributed were analyzed with the Wilcoxon signed rank test (paired data) or Wilcoxon rank sum test (not paired). Proportions captured at a single point in time were compared using the Chi square test. Repeated proportions were analyzed with McNemar’s test. We present unadjusted p-values throughout this article. Adjustment for multiple comparisons using the Benjamini-Hochberg procedure (which controls the false discovery rate) results in no change to the interpretation of any result. Using the more conservative Holm-Sidak procedure (which controls the familywise error rate), results only in the change in proportion of persons who exercised by running (see the Results section for details) to become nonsignificant.

Human Subjects Protection

This project was approved by the Institutional Review Board at the University of Kansas Medical Center.

Results

Characteristics of the sample are shown in Table 1. Of the 500 adult patient surveys collected, 382 were included after complete case analysis. Surveys that were improperly filled out were excluded in the final analysis. Respondents were widely distributed in age (median = 36 years; range = 18–74 years), sex, race, ethnicity, and educational attainment. Respondents were most commonly female (64.9%), White (65.7%), non-Hispanic (88%), college-educated (38%), single (41.6%) or married (40.8%), and residing without children (63.9%). Before the beginning of the pandemic, most participants were employed and worked away from home (56.8%); this proportion declined during the pandemic to 34.6%. Over the same period, the proportion of the population who was employed and working from home rose from 10.7% to 27%. Similarly, persons with a hybrid work setting (both at and away from home) rose from 9.4% to 13.1%.

First, we evaluated the change in weekly total duration of exercise (Table 1). This distribution exhibited left skew (Appendix Figure 1): The median change in weekly exercise duration showed no change (0.0 min/wk), but the mean change was –25.7 min/wk. Across all participants, total duration of exercise was decreased after the onset of the pandemic (paired Wilcoxon signed rank test $P < .001$).

Next, we evaluated the proportion of individuals reporting exercising >150 min/wk, as recommended by the Physical Activity Guidelines for Americans. Before the pandemic, 157 of 382 (41.1%) persons met this guideline. During the pandemic, only 119 of 382 (31.1%) persons reported exercising >150 min/wk. Many individuals meeting the threshold before the pandemic fell short of meeting the threshold during the pandemic (63 of 157, 40.1%). Meanwhile, only 25 of the 225 (11.1%) of individuals not meeting the threshold started meeting the threshold during the pandemic. Significantly fewer individuals therefore met this guideline during the pandemic as compared with before the pandemic (McNemar's test $P < .001$).

We then evaluated the number and duration of weekly exercise sessions before and during the

pandemic (Table 1 and Appendix Figure 2). The median interquartile range (IQR) exercise session duration before the pandemic was 36.2 minutes (20.0 to 60.0) as compared with 30.0 minutes (10.0 to 45.0) during the pandemic (paired Wilcoxon signed rank test $P < .001$). Likewise, the median (IQR) number of days exercised before the pandemic was 3 (2 to 4) days per week, while during the pandemic it was 2.5 (1 to 4) days per week (paired Wilcoxon signed rank test $P < .001$).

We next evaluated acquisition of home exercise equipment. Before the pandemic, 170 of 382 (44.5%) had home exercise equipment. Interestingly, only 90 of 382 (23.6%) purchased home exercise equipment, including 52 of the 170 (30.6%) of those who already had home equipment and only 38 of the 212 (17.9%) who lacked such equipment before the pandemic. Individuals who already had home exercise equipment before the pandemic were therefore more likely to acquire more equipment than those who had none to begin with (Chi square test $P = .005$).

Significant differences were found in the setting of exercise activity (Table 2). Individuals reported decreased participation in sports, on-site group classes, weightlifting, running and swimming (McNemar's P values shown in Table 2). In contrast, a greater proportion of individuals reported participation in virtual group classes (6.3% before the pandemic vs 13.1% during the pandemic; McNemar's $P < .001$).

We then evaluated changes in weight. The plurality (162 of 382, 42.4%) of respondents reported no change in their weight (Table 3). Among those with a weight change more than 4 lb, more individuals reported weight gain (148 of 382, 38.7%) than did weight loss (72 of 382, 18.8%). Overall, this represents a significant trend in the direction of weight gain (Wilcoxon signed rank test $P < .001$).

Finally, we evaluated whether change in employment setting was associated with change in either (1) total weekly exercise duration (Table 4) or (2) weight (Table 5). Among persons who worked away from home before the pandemic ($n = 217$), we failed to detect a difference in change in weekly exercise (Wilcoxon rank sum $P = .141$) or change in weight (Wilcoxon rank sum $P = .974$) among persons who continued to work away from home after the pandemic ($n = 123$) versus those who did not ($n = 94$, including work at home setting, hybrid work away and at home setting and unemployed

Table 1. Sample Demographics, Exercise Session Days/Wk, Session Duration, and Total Exercise Min/Wk

Category	No. (%)	Session days/wk, median (IQR)			Session duration, minutes, median (IQR)			Total exercise min/wk, median (IQR)		
		Before	During	Change	Before	During	Change	Before	During	Change
All respondents	382 (100%)	3.0 (2.0)	2.5 (3.0)	0.0 (1.0)	36.2 (40.0)	30.0 (35.0)	0.0 (10.0)	120 (162.5)	85.0 (167.8)	0.0 (60.0)
Age, years										
18 to 24	46 (12.0)	3.0 (2.0)	2.8 (2.9)	0.0 (1.0)	40.0 (30.0)	30.0 (29.4)	0.0 (13.8)	120.0 (120.0)	75.0 (118.1)	0.0 (82.5)
25 to 34	123 (32.2)	3.0 (2.0)	2.5 (3.0)	0.0 (1.0)	40.0 (30.0)	30.0 (37.5)	0.0 (12.5)	120.0 (150.0)	90.0 (169.4)	0.0 (63.8)
35 to 44	91 (23.8)	3.0 (2.0)	2.5 (3.2)	0.0 (2.0)	45.0 (30.0)	30.0 (42.5)	0.0 (20.0)	120.0 (160.0)	90.0 (170.1)	0.0 (90.0)
45 to 54	65 (17.0)	3.0 (3.0)	2.0 (4.0)	0.0 (1.0)	30.0 (40.0)	30.0 (45.0)	0.0 (7.5)	75.0 (157.5)	60.0 (150.0)	0.0 (37.5)
55 to 64	39 (10.2)	3.0 (3.0)	3.0 (2.5)	0.0 (0.0)	30.0 (25.0)	30.0 (32.5)	0.0 (0.0)	135.0 (156.2)	100.0 (180.0)	0.0 (0.0)
65 to 74	18 (4.7)	3.5 (3.8)	2.0 (4.5)	0.0 (0.8)	30.0 (23.8)	30.0 (24.5)	0.0 (0.0)	120.0 (127.5)	75.0 (151.8)	0.0 (23.8)
Sex										
Female	248 (64.9)	3.0 (2.0)	2.5 (3.0)	0.0 (1.0)	30.0 (32.5)	30.0 (35.0)	0.0 (7.5)	90.0 (135.0)	72.5 (145.2)	0.0 (60.0)
Male	134 (35.1)	3.0 (3.0)	3.0 (3.0)	0.0 (1.5)	45.0 (30.0)	30.0 (54.9)	0.0 (15.0)	142.5 (170.6)	102.5 (206.2)	0.0 (90.0)
Gender										
Man	135 (35.3)	3.0 (3.0)	3.0 (3.0)	0.0 (1.5)	45.0 (30.0)	30.0 (53.2)	0.0 (15.0)	135.0 (168.8)	100.0 (203.4)	0.0 (90.0)
Woman	247 (64.7)	3.0 (2.0)	2.5 (3.0)	0.0 (1.0)	30.0 (32.5)	30.0 (35.0)	0.0 (7.5)	90.0 (135.0)	70.0 (145.2)	0.0 (60.0)
Race										
Asian	14 (3.7)	3.0 (0.9)	3.0 (2.5)	-0.5 (2.1)	60.0 (15.0)	47.5 (30.0)	0.0 (9.4)	176.2 (60.0)	120.0 (93.8)	-5.0 (95.6)
Black	86 (22.5)	3.0 (1.9)	2.0 (3.0)	0.0 (1.0)	30.0 (53.8)	20.0 (37.5)	0.0 (10.0)	90.0 (160.0)	38.8 (110.6)	0.0 (45.0)
None	5 (1.3)	5.0 (1.0)	4.0 (4.0)	1.0 (1.5)	30.0 (30.0)	5.0 (30.0)	10.0 (30.0)	120.0 (210.0)	30.0 (120.0)	0.0 (15.0)
Other/multiple	26 (6.8)	3.0 (1.9)	3.5 (3.4)	0.0 (1.9)	30.0 (25.0)	16.0 (30.0)	0.0 (18.8)	90.0 (107.5)	30.0 (101.2)	0.0 (71.2)
White	251 (65.7)	3.0 (3.0)	3.0 (3.0)	0.0 (1.0)	40.0 (30.0)	30.0 (25.0)	0.0 (7.5)	120.0 (154.4)	90.0 (150.0)	0.0 (60.0)
Ethnicity										
Hispanic	46 (12.0)	3.0 (2.0)	2.0 (4.0)	0.0 (1.5)	33.8 (37.5)	30.0 (45.0)	0.0 (15.0)	102.5 (131.2)	63.8 (159.4)	0.0 (60.0)
Not Hispanic	336 (88.0)	3.0 (2.0)	2.8 (3.0)	0.0 (1.0)	36.2 (40.0)	30.0 (33.1)	0.0 (8.1)	120.0 (165.0)	90.0 (165.0)	0.0 (60.0)
Educational attainment										
Doctorate/professional degree	25 (6.5)	3.0 (2.0)	4.0 (4.0)	0.0 (2.0)	45.0 (22.5)	40.0 (30.0)	0.0 (7.5)	180.0 (135.0)	135.0 (142.5)	0.0 (160.0)
Master's degree	42 (11.1)	3.0 (2.0)	2.0 (2.4)	0.0 (1.8)	40.0 (30.0)	38.8 (28.1)	0.0 (0.0)	150.0 (194.1)	167.5 (172.5)	0.0 (96.9)
College graduate	383 (51.1)	3.0 (2.0)	3.0 (3.5)	0.0 (1.0)	45.0 (30.0)	30.0 (30.0)	0.0 (0.0)	120.0 (158.8)	96.0 (145.2)	0.0 (0.0)
Some college	171 (30.6)	3.0 (2.0)	3.0 (3.0)	0.0 (1.5)	30.0 (40.0)	30.0 (45.0)	0.0 (11.0)	90.0 (131.0)	69.0 (157.5)	0.0 (0.0)
High school graduate	42 (11.1)	3.0 (2.4)	2.0 (2.9)	0.0 (0.4)	27.5 (39.4)	22.5 (26.1)	0.0 (2.2)	83.8 (125.8)	69.0 (104.9)	0.0 (27.8)
Some high school or less	11 (2.9)	3.0 (3.0)	3.0 (2.5)	0.0 (1.5)	15.0 (27.5)	0.0 (22.5)	0.0 (2.5)	45.0 (87.5)	0.0 (60.0)	0.0 (15.0)

Continued

Table 1. Continued

Category	No. (%)	Session days/wk, median (IQR)			Session duration, minutes, median (IQR)			Total exercise min/wk, median (IQR)		
		Before	During	Change	Before	During	Change	Before	During	Change
Before pandemic employment status										
Employed: work at home	41 (10.7)	3.0 (2.0)	3.0 (4.0)	0.0 (1.5)	37.5 (40.0)	30.0 (45.0)	0.0 (30.0)	135.0 (135.0)	60.0 (180.0)	0.0 (90.0)
Employed: work away from home	217 (56.8)	3.0 (2.0)	3.0 (3.0)	0.0 (1.0)	37.5 (30.0)	30.0 (30.0)	0.0 (10.0)	120.0 (150.0)	90.0 (165.0)	0.0 (60.0)
Employed: work at home and away from home	36 (9.4)	3.0 (2.1)	2.8 (3.1)	0.0 (0.6)	42.5 (30.0)	32.5 (31.9)	0.0 (1.2)	120.0 (137.8)	80.0 (161.2)	0.0 (37.5)
Retired	20 (5.2)	2.0 (3.5)	2.0 (2.6)	0.0 (0.0)	30.0 (18.8)	25.0 (28.9)	0.0 (1.2)	60.0 (112.5)	45.0 (108.0)	0.0 (3.8)
Student	26 (6.8)	3.2 (2.4)	3.0 (3.0)	0.0 (2.8)	45.0 (30.0)	30.0 (28.1)	0.0 (15.0)	142.5 (168.8)	90.0 (125.6)	0.0 (195.0)
Unemployed	42 (11.0)	3.0 (3.0)	2.2 (3.8)	0.0 (1.0)	30.0 (37.5)	27.5 (43.8)	0.0 (0.0)	90.0 (140.3)	60.0 (134.1)	0.0 (29.1)
During pandemic employment status										
Employed: work at home	103 (27.0)	3.0 (2.0)	3.0 (3.2)	0.0 (1.0)	45.0 (30.0)	30.0 (36.2)	0.0 (5.0)	120.0 (120.0)	90.0 (164.9)	0.0 (60.0)
Employed: work away from home	132 (34.6)	3.0 (2.0)	2.5 (2.6)	0.0 (1.0)	37.5 (30.0)	30.0 (41.9)	0.0 (20.6)	120.0 (166.2)	84.4 (169.6)	0.0 (80.0)
Employed: work at home and away from home	50 (13.1)	3.0 (2.5)	2.0 (2.9)	0.0 (1.4)	32.5 (28.2)	30.0 (22.5)	0.0 (0.2)	108.8 (135.0)	68.8 (150.0)	0.0 (56.2)
Retired	18 (4.7)	2.0 (4.5)	2.0 (2.4)	0.0 (0.0)	30.0 (18.8)	25.0 (24.5)	0.0 (0.0)	55.0 (137.5)	45.0 (103.0)	0.0 (0.0)
Student	23 (6.0)	3.0 (2.8)	3.0 (3.0)	0.0 (2.5)	45.0 (30.0)	30.0 (26.2)	0.0 (15.0)	135.0 (172.5)	90.0 (123.8)	0.0 (180.0)
Unemployed	56 (14.7)	3.0 (2.0)	2.0 (4.0)	0.0 (1.0)	30.0 (50.0)	27.5 (38.1)	0.0 (9.2)	90.0 (159.2)	60.0 (132.2)	0.0 (31.9)
Marital status										
Divorced and not remarried	74 (12.3)	3.0 (4.0)	2.0 (4.0)	0.0 (2.0)	30.0 (42.5)	30.0 (9.2)	0.0 (15.0)	105.0 (192.5)	60.0 (170.0)	0.0 (70.0)
Married	156 (40.8)	3.0 (2.5)	3.0 (3.1)	0.0 (1.0)	37.5 (30.0)	30.0 (27.0)	0.0 (7.5)	140.0 (138.8)	97.5 (191.2)	0.0 (60.0)
Married but separated	11 (2.9)	2.5 (1.5)	2.5 (2.2)	0.0 (0.2)	30.0 (47.5)	30.0 (17.5)	0.0 (8.8)	90.0 (132.5)	90.0 (132.5)	0.0 (37.5)
Single (never married)	159 (41.6)	3.0 (2.0)	2.0 (3.5)	0.0 (1.0)	30.0 (40.0)	30.0 (44.4)	0.0 (13.0)	90.0 (137.5)	60.0 (120.0)	0.0 (60.0)
Widowed and not remarried	9 (2.4)	3.0 (4.0)	3.0 (5.0)	0.0 (0.0)	20.0 (20.0)	30.0 (20.0)	0.0 (0.0)	60.0 (150.0)	90.0 (120.0)	0.0 (30.0)
Children in household 10 years or younger										
No	288 (5.4)	3.0 (2.5)	3.0 (3.0)	0.0 (1.0)	32.5 (40.0)	30.0 (5.0)	0.0 (7.5)	120.0 (191.2)	90.0 (191.0)	0.0 (60.0)
Yes	46 (4.2)	3.0 (2.0)	3.0 (2.4)	0.0 (1.0)	37.5 (33.8)	30.0 (43.4)	0.0 (15.0)	77.6 (135.0)	97.6 (147.4)	0.0 (60.0)
Children in household 11 years or older										
No	103 (8.8)	3.0 (2.0)	3.0 (3.0)	0.0 (1.0)	37.5 (33.8)	30.0 (30.0)	0.0 (0.0)	120.0 (120.0)	90.0 (165.0)	0.0 (0.0)
Yes	18 (2.1)	3.0 (2.0)	3.0 (3.0)	0.0 (1.0)	30.0 (40.0)	30.0 (30.0)	0.0 (0.0)	90.0 (157.5)	75.0 (165.0)	0.0 (45.0)

Abbreviation: IQR, interquartile range.

Table 2. Exercise Setting Before and During the Pandemic

Type of activity	No. (%) reporting activity		<i>P</i> ^a
	Before	During	
Sports	55 (14.4)	27 (7.1)	<.001
Group classes (virtual)	24 (6.3)	50 (13.1)	<.001
Group classes (on site)	58 (15.2)	9 (2.4)	<.001
Weightlifting	113 (29.6)	72 (18.8)	<.001
Walking	257 (67.3)	247 (64.7)	.229
Running	111 (29.1)	93 (24.3)	.010 ^b
Biking	57 (14.9)	47 (12.3)	.089
Swimming	26 (6.8)	13 (3.4)	.006

^aMcNemar's paired chi-square test.

^bControlling for multiple comparisons using the Holm-Sidak procedure, rather than the Benjamini-Hochberg, attenuates the statistical significance of this result (Holm-Sidak adjusted *P* = .073).

persons). Similarly, we failed to detect an association between change in total weekly exercise duration (Wilcoxon rank sum *P* = .946) or weight (Wilcoxon rank sum *P* = .085) among persons who were employed before the pandemic (*n* = 294), stratified by whether they remained employed (*n* = 276) or became unemployed (*n* = 18).

Discussion

The aim of this study was to investigate the changes in exercise habits in the US workforce during the "Stay at Home" order of the COVID-19 outbreak in the US. This study did not find any significant change in total weekly exercise duration when stratified by change in employment status. However, across all participants, total duration of exercise was decreased after the onset of the pandemic. This finding is consistent with a similar study on the change of exercise habits during COVID-19 confinement.¹ This large international study (*n* = 1047) found a negative impact on physical activity on all levels of intensity and a 33.5% decrease in overall weekly physical activity.¹ This same international study also found that food consumption and meal patterns (the type of food, eating out of control, snacks between meals, number of main meals) were unhealthier during confinement.¹ This could explain in part why 38.7% of our study participants reported weight gain during the pandemic. Our study is more applicable to the US population because the international study included only 31

participants who were included from continents other than Asia, Europe, and Africa.¹

Importantly, our study found that before the onset of the pandemic, 41.1% of participants reported exercising more than 150 min/wk as recommended by the Physical Guidelines for Americans. After the onset of the pandemic this had a significant fall to 31.1% meeting this threshold. This fall in exercise participation occurred despite 23.6% of participants purchasing new home exercise equipment during the pandemic.

These findings are important in light of emerging evidence that physical activity should be promoted as a nonpharmacologic intervention to protect against 2019 novel coronavirus.^{10,16} A large study (*n* = 48,440) found that patients with COVID-19 who were consistently inactive were at increased risk of hospitalization, ICU admission, and death.²² Exercise-induced immunomodulation has been recognized for over 3 decades, with around 5000 peer reviewed original and review articles available in both MEDLINE and PubMed databases.¹⁶ Researchers suggest physical exercise can act as an immunomodulator by increasing pro- and anti-inflammatory cytokines such as IL-10 and IFN- β .¹⁰ Exercise also increases lymphocyte circulation and cell recruitment.¹⁰ This can lower incidence, symptom intensity, and mortality in viral infections.¹⁰ Suppression of IFN- α and IFN- β has been theorized as a possible pathology to COVID-19.¹⁰ These cytokines are especially important because they act as antivirals by influencing activities of macrophages and lymphocytes.¹⁰ There has been evidence that physical activity leads to decreased acute respiratory illness incidence, duration, and intensity of symptoms, as well as mortality.¹⁰ Heffmerna et al have suggested that exercise training could also improve immune response to COVID-19 by augmenting the ACE2-Ang1-7-Mas receptor axis to reduce pulmonary fibrosis.¹³ Carter et al proposed that regular physical activity, when particularly done outside, can alleviate risk factors for COVID-19 by increasing vitamin D exposure as well as decreasing adiposity.⁶

Another manner in which physical activity may benefit US Americans during the pandemic is through the relief of psychological stress. Studies have consistently found a relationship exists between physical inactivity and an increased risk for depression and anxiety disorders.^{9,14} This is specifically important in the setting of a global

Table 3. Self-Reported Change in Weight Stratified by Demographic Variables

Category	No. (%)	Weight change >4 lbs, No. (%)		
		Increased	No change	Decreased
All respondents	382 (100.0)	148 (38.7)	162 (42.4)	72 (18.8)
Age, years				
18 to 24	46 (12.0)	17 (37.0)	25 (54.3)	4 (8.7)
25 to 34	123 (32.2)	46 (37.4)	50 (40.7)	27 (22.0)
35 to 44	91 (23.8)	36 (39.6)	38 (41.8)	17 (18.7)
45 to 54	65 (17.0)	32 (49.2)	22 (33.8)	11 (16.9)
55 to 64	39 (10.2)	13 (33.3)	17 (43.6)	9 (23.1)
65 to 74	18 (4.7)	4 (22.2)	10 (55.6)	4 (22.2)
Sex				
Male	134 (35.1)	50 (37.3)	62 (46.3)	22 (16.4)
Female	248 (64.9)	98 (39.5)	100 (40.3)	50 (20.2)
Gender				
Man	135 (35.3)	50 (37.0)	62 (45.9)	23 (17.0)
Woman	247 (64.7)	98 (39.7)	100 (40.5)	49 (19.8)
Race				
Asian	14 (3.7)	5 (35.7)	9 (64.3)	0 (0.0)
Black	86 (22.5)	44 (51.2)	26 (30.2)	16 (18.6)
None	5 (1.3)	3 (60.0)	2 (40.0)	0 (0.0)
Other/multiple	26 (6.8)	12 (46.2)	10 (38.5)	4 (15.4)
White	251 (65.7)	84 (33.5)	115 (45.8)	52 (20.7)
Ethnicity				
Hispanic	46 (12.0)	16 (34.8)	19 (41.3)	11 (23.9)
Non-Hispanic	336 (88.0)	132 (39.3)	143 (42.6)	61 (18.2)
Educational attainment				
Some high school or less	11 (2.9)	6 (54.5)	4 (36.4)	1 (9.1)
High school graduate	42 (11.0)	15 (35.7)	22 (52.4)	5 (11.9)
Some college	117 (30.6)	43 (36.8)	51 (43.6)	23 (19.7)
College graduate	145 (38.0)	59 (40.7)	59 (40.7)	27 (18.6)
Master's degree	42 (11.0)	14 (33.3)	20 (47.6)	8 (19.0)
Doctorate/Professional degree	25 (6.5)	11 (44.0)	6 (24.0)	8 (32.0)
Before pandemic employment status				
Unemployed	42 (11.0)	19 (45.2)	12 (28.6)	11 (26.2)
Employed: work away from home	217 (56.8)	81 (37.3)	94 (43.3)	42 (19.4)
Employed: work at home	41 (10.7)	19 (46.3)	21 (51.2)	1 (2.4)
Employed: work at home and away from home	36 (9.4)	12 (33.3)	13 (36.1)	11 (30.6)
Retired	20 (5.2)	8 (40.0)	9 (45.0)	3 (15.0)
Student	26 (6.8)	9 (34.6)	13 (50.0)	4 (15.4)
During pandemic employment status				
Unemployed	56 (14.7)	27 (48.2)	19 (33.9)	10 (17.9)
Employed: work away from home	132 (34.6)	51 (38.6)	53 (40.2)	28 (21.2)
Employed: work at home	103 (27.0)	36 (35.0)	51 (49.5)	16 (15.5)
Employed: work at home and away from home	50 (13.1)	21 (42.0)	17 (34.0)	12 (24.0)
Retired	18 (4.7)	7 (38.9)	9 (50.0)	2 (11.1)
Student	23 (6.0)	6 (26.1)	13 (56.5)	4 (17.4)
Marital status				
Single (never married)	159 (41.6)	61 (38.4)	70 (44.0)	28 (17.6)
Married	156 (40.8)	62 (39.7)	68 (43.6)	26 (16.7)
Married but separated	11 (2.9)	2 (18.2)	6 (54.5)	3 (27.3)

Continued

Table 3. Continued

Category	No. (%)	Weight change ≥4 lbs, No. (%)		
		Increased	No change	Decreased
Widowed and not remarried	9 (2.4)	1 (11.1)	6 (66.7)	2 (22.2)
Divorced and not remarried	47 (12.3)	22 (46.8)	12 (25.5)	13 (27.7)
Children in household 10 years or younger				
Yes	94 (24.6)	42 (44.7)	34 (36.2)	18 (19.1)
No	288 (75.4)	106 (36.8)	128 (44.4)	54 (18.8)
Children in household 11 years or greater				
Yes	81 (21.2)	36 (44.4)	30 (37.0)	15 (18.5)
No	301 (78.8)	112 (37.2)	132 (43.9)	57 (18.9)

pandemic. Several studies and articles have been published regarding the psychological effects of the COVID-19 pandemic. A study of 1210 respondents from 194 cities in China found that

54% of respondents reported the psychological impact of the COVID 19 outbreak as moderate to severe.²⁴ This same study also found that 29% of respondents reported moderate to severe

Table 4. Employment Status Before and During the Pandemic and Change in Total Weekly Exercise

Employment status during pandemic	No. (%)	Exercise min/wk, Median (IQR)		
		Before	During	Change
Before pandemic: unemployed				
Employed: work away from home	4 (1.0)	125.6 (81.6)	72.5 (93.8)	-53.1 (68.4)
Unemployed	36 (9.4)	88.8 (124.7)	60.0 (132.2)	0.0 (16.2)
Employed: work at home	2 (0.5)	153.8 (18.8)	105.0 (30.0)	-48.8 (48.8)
Before pandemic: employed, work away from home				
Employed: work away from home	123 (32.2)	120.0 (167.5)	90.0 (165.8)	0.0 (80.0)
Unemployed	15 (3.9)	120.0 (177.5)	30.0 (120.0)	0.0 (135.0)
Employed: work at home	58 (15.2)	112.5 (120.0)	112.5 (142.5)	0.0 (57.5)
Employed: work at home and away from home	21 (5.5)	135.0 (127.5)	90.0 (127.5)	0.0 (65.0)
Before pandemic: employed, work at home				
Employed: work away from home	1 (0.3)	90.0 (0.0)	0.0 (0.0)	-90.0 (0.0)
Unemployed	2 (0.5)	100.5 (79.5)	105.0 (105.0)	4.5 (184.5)
Employed: work at home	36 (9.4)	137.5 (138.8)	67.5 (195.0)	0.0 (63.8)
Employed: work at home and away from home	2 (0.5)	90.0 (90.0)	7.5 (7.5)	-82.5 (82.5)
Before pandemic: employed, work both at home and away from home				
Employed: work away from home	3 (0.8)	120.0 (120.0)	100.0 (112.5)	-15.0 (10.0)
Unemployed	1 (0.3)	120.0 (0.0)	120.0 (0.0)	0.0 (0.0)
Employed: work at home	5 (1.3)	180.0 (30.0)	120.0 (120.0)	0.0 (240.0)
Employed: work at home and away from home	27 (7.1)	100.0 (148.1)	67.5 (172.5)	0.0 (40.0)
Before pandemic: retired				
Employed: work away from home	1 (0.3)	80.0 (0.0)	0.0 (0.0)	-80.0 (0.0)
Employed: work at home	1 (0.3)	60.0 (0.0)	112.5 (0.0)	52.5 (0.0)
Retired	18 (4.7)	55.0 (137.5)	45.0 (103.0)	0.0 (0.0)
Before pandemic: student				
Unemployed	2 (0.5)	187.5 (52.5)	165.0 (135.0)	-22.5 (187.5)
Employed: work at home	1 (0.3)	240.0 (0.0)	120.0 (0.0)	-120.0 (0.0)
Student	23 (6.0)	135.0 (172.5)	90.0 (123.8)	0.0 (180.0)

Abbreviations: IQR, interquartile range.

Table 5. Self-Reported Change in Weight Stratified by Before and During Pandemic Employment Status

During pandemic employment status	No. (%)	Weight change more than 4 lbs, No. (%)		
		Increased	No change	Decreased
Before pandemic: unemployed				
Employed: work away from home	4 (1.0)	2 (50.0)	1 (25.0)	1 (25.0)
Unemployed	36 (9.4)	16 (44.4)	10 (27.8)	10 (27.8)
Employed: work at home	2 (0.5)	1 (50.0)	1 (50.0)	0 (0.0)
Before pandemic: employed, work away from home				
Employed: work away from home	123 (32.2)	47 (38.2)	51 (41.5)	25 (20.3)
Unemployed	15 (3.9)	8 (53.3)	7 (46.7)	0 (0.0)
Employed: work at home	58 (15.2)	17 (29.3)	28 (48.3)	13 (22.4)
Employed: work at home and away from home	21 (5.5)	9 (42.9)	8 (38.1)	4 (19.0)
Before pandemic: employed, work at home				
Employed: work away from home	1 (0.3)	1 (100.0)	0 (0.0)	0 (0.0)
Unemployed	2 (0.5)	1 (50.0)	1 (50.0)	0 (0.0)
Employed: work at home	36 (9.4)	16 (44.4)	19 (52.8)	1 (2.8)
Employed: work at home and away from home	2 (0.5)	1 (50.0)	1 (50.0)	0 (0.0)
Before pandemic: employed, work at home and away from home				
Employed: work away from home	3 (0.8)	0 (0.0)	1 (33.3)	2 (66.7)
Unemployed	1 (0.3)	0 (0.0)	1 (100.0)	0 (0.0)
Employed: work at home	5 (1.3)	1 (20.0)	3 (60.0)	1 (20.0)
Employed: work at home and away from home	27 (7.1)	11 (40.7)	8 (29.6)	8 (29.6)
Before pandemic: retired				
Employed: work away from home	1 (0.3)	1 (100.0)	0 (0.0)	0 (0.0)
Employed: work at home	1 (0.3)	0 (0.0)	0 (0.0)	1 (100.0)
Retired	18 (4.7)	7 (38.9)	9 (50.0)	2 (11.1)
Before pandemic: student				
Unemployed	2 (0.5)	2 (100.0)	0 (0.0)	0 (0.0)
Employed: work at home	1 (0.3)	1 (100.0)	0 (0.0)	0 (0.0)
Student	23 (6.0)	6 (26.1)	13 (56.5)	4 (17.4)

anxiety symptoms.²⁰ A meta-analysis of 6 studies suggests that exercise should be considered an evidence-based treatment for anxiety.²³ Zhang et al found that physical activity directly alleviated general negative emotions during the COVID-19 pandemic in a cohort of 66 college students in China.²⁵ From this study, it was found that about 108 minutes of light intensity, 80 minutes of moderate intensity, or 45 minutes of vigorous intensity exercise daily was a suitable amount of exercise to minimize negative emotions during the pandemic.²⁵ This is why it is important for physicians to promote exercise in their patients as a preventative measure against COVID-19 induced psychological issues.

This study has limitations. Self-reported physical activity data may be over or underestimated by participants. There is also the possibility of recall bias

because participants were asked to remember exercise habits from before February 2020, several months before the administration of the survey in the fall of 2020. An added bias may also be present by asking participants to evaluate their exercise habits before the pandemic and during the pandemic at the same point in time. There is possible selection bias of a healthy population by surveying patients willing to go to clinic during the COVID-19 pandemic when many delayed seeking routine medical care. Another limitation is that a nonvalidated form of the International Physical Activity Questionnaire was used to evaluate exercise habits. A shorter version was used to maintain brevity in the survey and increase participation. Other limitations include that it was conducted at a single center and was limited to English-speaking participants. Our results are likely generalizable to patients seeing primary

care at urban academic medical centers in the US; additional studies in community health centers and in rural locations are warranted.

Conclusion

In summary, the COVID-19 pandemic significantly altered exercise habits (decreasing total weekly duration of exercise, decreasing satisfaction of exercise guidelines, and changing the setting of exercise activity). These alterations occurred alongside a trend toward significant increases in weight. Although we observed large changes in employment setting—toward work-at-home and hybrid at-home/away settings—we did not find an association with post-pandemic work setting and either exercise duration or weight. Our findings illuminate the exercise experience of individuals in the United States seeking primary care; this group is important because it is the group primary care physicians can intervene on to promote health outcomes. Further research to define pandemic-related changes in exercise habits of individuals in community and rural settings would be helpful. Further studies are also needed to assess strategies to maintain healthy habits during and post pandemic.

We thank the nursing and administrative staff at the University of Kansas Medical Center family medicine clinic for their assistance with distributing and collecting the survey.

To see this article online, please go to: <http://jabfm.org/content/35/2/295.full>.


References

1. Ammar A, Brach M, Trabelsi K, et al. Effects of COVID-19 home confinement of eating behavior and physical activity: results of the ECLB-COVID19 international online survey. *Nutrients* 2020;12:1583.
2. Arem H, Moore SC, Patel A, et al. Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship. *JAMA Intern Med* 2015;175:959–67.
3. Bláfoss R, Micheletti JK, Sundstrup E, Jakobsen MD, Bay H, Andersen LL. Is fatigue after work a barrier for leisure-time physical activity? cross-sectional study among 10,000 adults from the general working population. *Scand J Public Health* 2019; 47:383–91.
4. Blake H, Stanulewicz N, McGill F. Predictors of physical activity and barriers to exercise in nursing and medical students. *J Adv Nurs* 2017;73: 917–29.
5. Caban-Martinez A, Lee D, Fleming LE, et al. Leisure-time physical activity levels of the US workforce. *Prev Med* 2007;44:432–6.
6. Carter SJ, Baranuskas MN, Fly AD. Considerations for obesity, vitamin D, and physical activity amid the COVID-19 pandemic. *Obesity* 2020;28:1176–7.
7. CDC COVID Data Tracker [Internet]. Centers for Disease Control and Prevention; 2020 [cited 2020 December 2]. Available at: https://covid.cdc.gov/covid-data-tracker/#cases_casesper100klast7days.
8. Choi J, Lee M, Lee J, Kang D, Choi J. Correlates associated with participation in physical activity among adults: a systematic review of reviews and update. *BMC Public Health* 2017;17:356.
9. Crowley S. Exercise is medicine: the role of exercise in the prevention and treatment of mood and anxiety disorders in children and adults. *Psychoneuroendocrinology* 2015;61:18.
10. da Silveira MP, da Silva Fagundes KK, Bizuti MR, et al. Physical exercise as a tool to help the immune system against COVID-19: an integrative review of the current literature. *Clin Exp Med* 2021;21:15–28.
11. Executive Order No. 20-16 [Internet]. State of Kansas, Governor Laura Kelly; 2020 [cited 2020 December 2]. Available at: <https://governor.kansas.gov/wp-content/uploads/2020/03/EO20-16.pdf>.
12. Hagströmer M, Oja P, Sjörström M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr* 2006;9:755–62.
13. Heffernan KS, Jae SY. Exercise as medicine for COVID-19: an ace in the hole? *Med Hypotheses* 2020;142:109835.
14. Hiles SA, Lamers F, Milaneschi Y, Penninx BWJH. Sit, step, sweat: longitudinal associations between physical activity patterns, anxiety and depression. *Psychol Med* 2017;47:1466–77.
15. Holtermann A, Krause N, Van der Beek AJ, Straker A. The physical activity paradox: six reasons why occupational activity (OPA) does not confer the cardiovascular health benefits that leisure time physical activity does. *Br J Sports Med* 2018;52:149–50.
16. Leandro CG, Ferreira e Silva WT, Lima-Silva AE. COVID-19 and exercise-induced immunomodulation. *Neuroimmunomodulation* 2020;27:75–8.
17. Martinez Y, Harmon B, Nigg C, Bantum E, Strayhorn S. Diet and physical activity intervention strategies for college students. *Health Behav Policy Rev* 2016;3:336–47.
18. Harris P, Taylor AR, Minor BL, REDCap Consortium, et al. The REDCap Consortium: building an international community of software partners. *J Biomed Inform* 2019;95:103208.
19. Harris P, Taylor A, Thielke R, Payne R, Gonzalez JN, Conde JG. Research Electronic Data Capture (REDCap)—a metadata-driven methodology and


- workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
20. Parra D, Wetherell JL, Van Zandt A, Brownson RC, Abhishek J, Lenze E. A qualitative study of older adults' perspectives on initiating exercise and mindfulness practice. *BMC Geriatr* 2019;19:354.
 21. Physical Activity Guidelines Advisory Committee. *Physical Activity Guidelines for Americans*. 2nd ed. US Dept of Health and Human Services; 2018.
 22. Sallis R, Young DR, Tartof SY, et al. Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48440 adult patients. *Br J Sports Med* 2021;55:1099–105. Published April 13. Access April 19, 2021. <https://pubmed.ncbi.nlm.nih.gov/33849909/>.
 23. Stubbs B, Vancampfort D, Rosenbaum S, et al. An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: a meta-analysis. *Psychiatry Res* 2017;249:102–8.
 24. Wang C, Pan R, Wan X, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *IJERPH* 2020;17:1729.
 25. Zhang Y, Zhang H, Xindong M, Qian D. Mental health problems during COVID-19 pandemics and mitigation effects of exercise: a longitudinal study of college students in China. *IJERPH* 2020;17:3722.
 26. Zhao M, Veeranki S, Li S, Steffen LM, Xi B. Beneficial associations of low and large doses of leisure time physical activity with all-cause cardiovascular disease and cancer mortality: a national cohort study of 88,140 US adults. *Br J Sports Med* 2019;53:1405–11.

Appendix.


Appendix Figure 1. Distribution of Change in Total Minutes of Weekly Exercise Among Respondents.



Appendix Figure 2. Distribution of Change in Total Weekly Exercise, Stratified by Demographic Variables.



Appendix Figure 3. Distribution of Change in Total Weekly Exercise, Stratified by Before the Pandemic Employment Status (Panels) and During the Pandemic Employment Status.



Research Survey About Exercise Habits Before and After the COVID-19 Pandemic.

This research survey asks about exercise habits before and after the COVID-19 pandemic. We are recruiting research participants for this study. Participation involves completing this survey. It will take about 10 minutes. No identifiable information will be collected about you. The survey is anonymous. There are no personal benefits or risks to participating in this study. Participation is voluntary, and you can stop taking the survey at any time. You will get the same medical care whether or not you complete the survey. If you are an employee or student at the University of Kansas Medical Center, your participation or answers will not affect employee or school evaluations. If you have any questions, please contact Nicole Yedlinsky, MD, at nyedlinsky@kumc.edu. For questions about the rights of research participants, you may contact the KUMC Institutional Review Board (IRB) at (913) 588-1240 or humansubjects@kumc.edu

1. What is your age?	Write in: _____ years
2. What was your biological sex at birth?	Male Female Other
3. What is your gender?	Man Woman Other
4. What is your married status?	Single (never married) Married Married but separated Widowed and not remarried Divorced and not remarried
5. What is your race? (Circle ALL that apply)	American Indian or Alaskan Native Asian Black or African American White or Caucasian Native Hawaiian or Other Pacific Islander Other
6. Are you of Hispanic, Latino or Spanish origin?	Yes No
7. What was the highest level of school you completed?	Some high school or less High school graduate Some college College graduate Master's degree Doctorate or professional degree
8. What state did you live in on February 1, 2020?	Write in: _____ (example: Kansas) Or circle: I did not live in the United States
9. What was your employment status on <u>February</u> 1, 2020? (Circle the one best answer)	Unemployed Employed: work away from home Employed: work at home Employed: work at home <u>and</u> away from home Retired Student
10. What was your employment status on <u>April</u> 15, 2020? (Circle the one best answer)	Unemployed Employed: work away from home Employed: work at home Employed: work at home <u>and</u> away from home Retired Student

(Continued)

11. Are there any other adults (18 years or older) in your home?	Yes	No
12. Are there any children 10 years old or younger in your home?	Yes	No
13. Are there any children 11 years old or older in your home?	Yes	No
14. Before the COVID-19 pandemic, how many days per week did you exercise in your free time?	Write in: _____ days per week	
15. Before the COVID-19 pandemic, how many minutes did you spend per exercise session?	Write in: _____ minutes per session	
16. Before the COVID-19 pandemic, what types of exercise did you participate in? (Circle <u>all</u> that apply)	Sports Group classes (virtual) Group classes (on-site) Weightlifting Walking Running Biking Swimming Other (specify): _____	
17. Before the COVID-19 pandemic, where did you normally exercise?	Outdoors Home	Indoor studio Gym Other
18. After the beginning of the COVID-19 pandemic (March 11, 2020), how many days per week did you exercise in your free time?	Write in: _____ days per week	
19. After the beginning of the COVID-19 pandemic (March 11, 2020), how many minutes did you spend per exercise session?	Write in: _____ minutes per session	
20. After the beginning of the COVID-19 pandemic (March 11, 2020), what types of exercise did you participate in? (Circle <u>all</u> that apply)	Sports Group classes (virtual) Group classes (on-site) Weightlifting Walking Running Biking Swimming Other (specify): _____	
21. After the beginning of the COVID-19 pandemic (March 11, 2020), where did you normally exercise?	Outdoors Home	Indoor studio Gym Other
22. Did you have exercise equipment at home on February 1, 2020?	Yes	No
23. After the beginning of the COVID-19 pandemic (March 11, 2020), did you purchase any home exercise equipment?	Yes	No
24. After the beginning of the COVID-19 pandemic (March 11, 2020), how did your weight change?	Increased by more than 4 pounds Stayed the same Decreased by more than 4 pounds	