

ORIGINAL RESEARCH

Primary Care Evaluation of a Hepatitis C Virus (HCV) Screening Education Intervention

Thomas Ludden, PhD, Lindsay Shade, PA-C, Jeremy Thomas, MSW, Mark W. Russo, MD, Michael Leonard, MD, Philippe J. Zamor, MD, Charity G. Patterson, PhD, and Hazel Tapp, PhD

Introduction: Increased screening efforts and the development of effective antiviral treatments have led to marked improvement in Hepatitis C (HCV) patient outcomes. However, many people in the US are believed to have undiagnosed HCV. Successful screening strategies and access to a coordinated system of care are critical for HCV affected adults. The objective of this study was to evaluate a primary care HCV screening education intervention that took place 2018 to 2019 to improve primary care training and management of patients after the implementation of the electronic medical record (EMR) screening alert.

Methods: Using 15 primary care practices located in vicinity of neighborhoods at-risk for higher rates of HCV infections, a stepped wedge randomized control study design was utilized to deliver an educational screening intervention. The education intervention was implemented sequentially with 5 practices being presented to every 3 months. Number of patients within the Baby Boomer cohort (birth years 1945–1965) were collected 3 months before the first practice receiving the intervention to 3 months after the last practice receiving the education intervention. The main outcome collected was the HCV screening. Generalized linear mixed models were used to test the hypothesis of improved screening rates after intervention implementation.

Results: There were a total of 85,697 patients within the Baby Boomer cohort seen at the 15 practices. Practices receiving the intervention had patients who were more likely to be screened for HCV ($\beta = 0.259$, $P < .001$; Odds Ratio [OR] [95%CI] 1.296 [1.098-1.529]). In terms of demographics, results showed that females are less likely to be screened than males ($\beta = -0.141$, $P < .001$; OR [95%CI] 0.868[0.813 to 0.927]), Baby Boomer patients aged less than 65 were more likely to be screened than Baby Boomer patients aged 65 and older ($\beta = 0.293$, $P < .001$; OR [95%CI] 1.340[1.251 to 1.436]).

Discussion: This study looked at screening rates before and after an educational intervention which happened subsequent to the activation of an EMR alert. Whereas HCV EMR alerts showed an increase in HCV screenings before the education intervention, the addition of the education showed a modest increase in HCV screening rates for Baby Boomer patients. (J Am Board Fam Med 2022;35:990–997.)

Keywords: Early Medical Intervention, Electronic Medical Record, Hepatitis C, Preventive Health Care, Primary Health Care, Screening Intervention

Introduction

Hepatitis C virus (HCV) infection is a disease that negatively impacts quality of life and results in

premature mortality if left untreated. An estimated 4.1 million persons living in the United States are HCV antibody-positive and 2.4 million persons are HCV positive with active viral replication (1% prevalence among adults). Of those people with chronic HCV, 75% are unaware that they are infected.^{1–3} Half of cases can be attributed to intravenous drug use. The rest of the cases result from sexual contact, birth mother to child, tattoos, intranasal drug use from shared paraphernalia, contaminated blood or transplants, and medical procedures.^{4–6} Persons born between 1945 and 1965 in the “Baby Boomer” generation were likely infected between 1960 and 1980 and account for the majority of all chronic HCV infections in adults. The Baby Boomer

This article was externally peer reviewed.
Submitted 16 December 2021; revised 20 May 2022; accepted 26 May 2022.

From Department of Family Medicine, Atrium Health, Charlotte, NC (TL, LS, JT, HT); Department of Hepatology, Atrium Health, Charlotte, NC (MWR, PJZ); Department of Infectious Diseases, Atrium Health, Charlotte, NC (ML); School of Health and Rehabilitation Sciences, University of Pittsburgh, Pittsburgh, PA (CGP).

Funding: This study, IN-US-337-4324, is supported by Gilead, Sciences, Inc.

Conflict of interest: None.

Corresponding author: Thomas Ludden, PhD, Atrium Health, 2001 Vail Avenue, Suite 400B, Charlotte, NC 28207 (E-mail: tom.ludden@atriumhealth.org).

generation is of interest because they represent more than 50% of newly identified cases in the past 10 years.⁷⁻⁹ HCV patients experienced a 5 to 20% increased risk for liver cirrhosis due to delayed treatment, putting patients at high risk for additional complications such as hepatocellular carcinoma (HCC).¹⁰ In 2019, more than 14,000 people had HCV as an underlying cause of death.¹¹ When this study was conducted, the Centers for Disease Control and Prevention (CDC) recommended all patients born between 1945 and 1965 be screened once for HCV. As of March 2020, the CDC began recommending a one-time HCV test for all adults age 18 to 79 to prevent HCV from spreading.¹²

The development of effective antiviral treatments along with increased screening efforts have led to improvements in HCV patient outcomes.¹⁰ Observational studies demonstrate a 40% reduction in HCC and liver failure in patients who achieve virologic cure along with reduction in mortality rates.¹³⁻¹⁶

Successful screening strategies and access to a coordinated system of care are critical for HCV affected adults.^{17,18} In May 2016, our large health care system implemented a one-time electronic medical record (EMR) alert for HCV screening for all patients born 1945-1965 who have no record of HCV antibody testing which resulted in a 17% absolute increase in screenings. A previous study also showing a 20% absolute increase in HCV screening when both HCV screening education and EMR alert activation took place at select practices took place during 2016.¹⁹

The objective of this study was to evaluate a primary care HCV screening education intervention using a stepped wedge randomized control trial design that took place from 2018 to 2019 to improve primary care training and management of patients by augmenting clinician and staff behavior after the implementation of the EMR screening alert. This study looked at screening rates before and after an education intervention which happened subsequent to the activation of the EMR alert. We hypothesized that HCV screenings would increase after the education intervention was rolled out at 15 primary care practices.²⁰ This stepped wedge design provides a more rigorous evaluation than a simple before and after intervention study which is scientifically flawed by confounding over time and does not include patients in a comparable control condition. The stepped wedge design

eventually included all practices so therefore prevents contamination and disappointment effects in practices not randomized to the intervention parallel randomized parallel designs.^{21,20}


Setting

The study was conducted at Atrium Health, a large, vertically integrated health care system, based in Charlotte, North Carolina with > 200 primary care offices and more than 12 million patient encounters per year.

Methods

The Institutional Review Board deemed this project as quality improvement because all patients born between 1945 and 1965 were receiving the screening alert and the project did not include research development, testing and evaluation designed to develop generalizable knowledge. Previous research identified 15 primary care practices located in vicinity of neighborhoods at-risk for higher rates of HCV infections.²² The research team, consisting of researchers, primary care providers, hepatologists, and physician assistants who were experienced in managing and treating HCV, worked together to develop an education intervention. An education presentation was developed using existing information from a previous study in a PowerPoint format with a prerecorded voice-over. Education content consisted of who to screen, what to do with positive results, and when and how to refer HCV patients. The education also included specific information around the needs for newly diagnosed HCV patients such as the referral process, treatment, side effects of treatment, potential interactions with current prescriptions, required labs, and guidance on how to prepare patients for treatment visits with specialists. In addition, algorithms providing decision support were developed to guide primary care providers with patients whose lab test returned with a positive HCV. Both the Infectious Disease department and the Center for Liver Disease and Transplantation within our health care system provided content for the intervention input and patient care for HCV positive patients. The information provided to participating departments followed the guidelines for the referral process within our health care system and treatment guidelines based on the medication options at the time of the study.

Figure 1. Study design for 15 practices into 3 wedges.



The intervention was rolled out as a 1 hour lunch time meeting at the practice. A PowerPoint presentation with the prerecorded voiceover was presented at the lunchtime meeting by a nonclinical research team member, and then questions about the presentation were answered and/or followed up by the appropriate clinical department. If a practice declined to participate with an in-person lunch time meeting, the PowerPoint was emailed to the practice manager to be distributed to the providers to view when convenient. Because this education was not required, there was no attestation for providers to view the presentation.

Study Design

The study utilized a stepped wedge randomized control trial design (Figure 1). Each practice was randomly assigned to 3 groups and the education

intervention was rolled out to 5 practices sequentially in 3-month intervals with the first 3 months used as a baseline. Practices were assigned randomly to wedges using a cluster randomization function in R. The evaluation consisted of data collection of patients within the Baby Boomer cohort across 4 time periods, (baseline and the 3 intervention rollout time periods) from July 1, 2018 to June 30, 2019. The education intervention was fully implemented by the end of the study, when all 15 practices receiving the education intervention, 5 in the first time period, 5 in the second time period, and 5 in the third time period.

The stepped wedge design was utilized to implement the intervention in a “phased, randomized roll-out,” where white cells indicate control time periods and gray cells indicate intervention time periods. Stepped wedge studies provide a more rigorous evaluation than a simple before/after study by: 1) allowing the research team and clinical teams to roll out the intervention in a timely, systematic manner, (2) increasing statistical power due to within and between cluster comparisons, (3) increasing clinical participation and leadership buy-in because all practices will eventually “flip” to the intervention, and (4) removing the confounding effects associated with time.²⁰

Data Collection

Patient data without a previous HCV screening was collected from our enterprise data warehouse aligning within each time period. Within each of the 15 practices, only 1 visit per person per time period without a previous HCV screening was used for the denominator. If the patient received an HCV screening during that time period, the patient was excluded from future time periods. Patients who were already screened in the past were excluded from data collection as the EMR alert would not be activated for those patients.

Table 1. Stepped Wedge Study Design and Number of Patients Eligible for Hepatitis C virus (HCV) Screening

	Time Period 1 Baseline (0–3 months)	Time Period 2 (3–6 months)	Time Period 3 (6–9 months)	Time Period 4 (9–12 months)
Practice 1 to 5 (Wedge 1)	6,206	<u>6,296</u>	<u>6,184</u>	<u>6,020</u>
Practice 6 to 10 (Wedge 2)	11,092	11,738	<u>11,460</u>	<u>10,727</u>
Practice 11 to 15 (Wedge 3)	4,244	3,962	4,087	<u>3,681</u>

Control periods before the HCV education intervention (non-underlined); intervention periods after the rollout of the HCV educational intervention (underlined).

Analysis

Generalized linear mixed models with a logit link were used to test the hypothesis of improved screening rates after intervention implementation controlling for random effects of practice location. The unit of analysis was any visit within the time frame where a patient was eligible for HCV screening (born between 1945 and 1965, no previous HCV diagnosis, no previous screening). The outcome of interest was HCV screening. The

primary variable of interest was the education intervention (yes/no). To prevent misclassification bias, we coded the patient data based on the date of the intervention provided to each practice. All patients before the presentation date were coded as control and patients after the presentation date as intervention. Patient level factors associated with the likelihood of screening were also explored such as gender, race, age at the date of screening, and ethnicity.

Table 2. Descriptive Statistics: Patient Population

	Control n = 41,329 ¹	Intervention n = 44,368 ¹	p-value
Wedge			<0.001
1	6,206 (15%)	18,500 (42%)	
2	22,830 (55%)	22,187 (50%)	
3	12,293 (30%)	3,681 (8.3%)	
Time Period			<0.001
1	21,542 (52%)	0 (0%)	
2	15,700 (38%)	6,296 (14%)	
3	4,087 (9.9%)	17,644 (40%)	
4	0 (0%)	20,428 (46%)	
White	29,526 (71%)	33,048 (74%)	<0.001
African American	9,507 (23%)	8,885 (20%)	<0.001
Asian	704 (1.7%)	807 (1.8%)	0.2
Multi-Race/Other	1,592 (3.9%)	1,628 (3.7%)	0.2
Age < 65	25,346 (61%)	26,540 (60%)	<0.001
Age 65 or older	15,983 (39%)	17,828 (40%)	<0.001
Baby Boomer	41,329 (100%)	44,368 (100%)	
Hispanic/Latino	1,234 (3.0%)	1,378 (3.1%)	0.3
Gender – Female	23,929 (58%)	25,364 (57%)	0.030
In Person	28,763 (70%)	30,588 (69%)	0.038
Site Name			<0.001
Practice 1	2,331 (5.6%)	2,115 (4.8%)	
Practice 2	1,796 (4.3%)	5,049 (11%)	
Practice 3	3,344 (8.1%)	3,260 (7.3%)	
Practice 4	3,912 (9.5%)	1,269 (2.9%)	
Practice 5	763 (1.8%)	2,304 (5.2%)	
Practice 6	1,411 (3.4%)	4,637 (10%)	
Practice 7	12,630 (31%)	12,292 (28%)	
Practice 8	2,106 (5.1%)	563 (1.3%)	
Practice 9	2,774 (6.7%)	791 (1.8%)	
Practice 10	1,623 (3.9%)	4,767 (11%)	
Practice 11	613 (1.5%)	1,743 (3.9%)	
Practice 12	1,555 (3.8%)	482 (1.1%)	
Practice 13	1,798 (4.4%)	1,901 (4.3%)	
Practice 14	1,946 (4.7%)	576 (1.3%)	
Practice 15	2,727 (6.6%)	2,619 (5.9%)	

¹n (%).


²Pearson's χ^2 test.

Preliminary sample size analysis predicted we would need at least 300 eligible patients per period per practice to detect a 10% absolute improvement in screening rates assuming the current screening rate is 20% in the Baby Boomer population.²³

Results

Table 1 shows the stepped wedge population by each wedge and time period. Rollout occurred at all 15 practices with 2 of the practices in time period 1, 4 in time period 2, and 2 in time period 3 that

Figure 2. Incidence rate of patients screened for Hepatitis C virus (HCV) for each time period without any previous record of HCV screening. Each wedge consists of 5 practices and each time period consist of 3 months.



received an in-person presentation with the presentation emailed to the remaining 7 practices.

Patient Characteristics

A total of 85 697 patient visits were included in our analyses (Table 2). This exceeded the overall number of patients originally proposed to be included in the model. Patient demographics were reflective of the population born between 1945 and 1965. Patient age demographic were divided into patients aged less than 65 and patients aged 65 and older. Both the preintervention and postintervention had higher proportions of female patients. The racial diversity of all the patients in the control condition vs the intervention condition were 71% to 74% for white, 23% to 21% for African American, 3% to 3% for Hispanic/Latino, and 3.8% to 3.6% for other race.

Screening Results

Figure 2 shows the screening rates at the 5 practices in each wedge and time period. Control periods were before the HCV education intervention and intervention periods are after the rollout of the HCV education intervention at the practice.

Results show patients born between 1945 and 1965 were more likely to be screened for HCV after their practice received the education intervention based on the interaction with an in-person presentation variable. (Table 3) ($\beta = 0.259$, $P < .001$; Odds Ratio [OR] [95%CI] 1.296 [1.098-1.529]).

In terms of demographics, results showed that females are less likely to be screened than males ($\beta = -0.141$, $P < .001$; OR [95%CI] 0.868[0.813 to 0.927]), Baby Boomer patients aged less than 65 were more likely to be screened than Baby Boomer patients aged 65 and older ($\beta = 0.293$, $P < .001$; OR [95%CI] 1.340[1.251 to 1.436]).

Discussion

The education intervention presented at the 15 practices, highlighting the screening need for HCV for the Baby Boomer generation, resulted in a significant increase in screening rates between the control time periods and the intervention time periods. Although the increase was statistically significant in aggregate with an absolute change of 0.3% (from 4.4% for the control time periods to 4.7% ($P < .001$) for the intervention time periods), the

Table 3. Generalized Linear Mixed Model Results

Variable	Beta (SE)	OR		
Constant	-3.327*** (0.395)	0.036	0.017	0.078
Intervention	-0.032 (0.090)	0.969	0.812	1.155
In Person	0.210 (0.528)	1.234	0.438	3.475
Gender - Female	-0.141*** (0.034)	0.868	0.813	0.927
Less than 65	0.293*** (0.035)	1.340	1.251	1.436
African American	-0.052 (0.041)	0.949	0.875	1.029
Asian	0.203* (0.113)	1.225	0.982	1.527
Multi-Race/Other	-0.092 (0.093)	0.912	0.760	1.094
Hispanic/Latino	0.039 (0.096)	1.039	0.861	1.254
Time Period 2	-0.202*** (0.053)	0.817	0.736	0.906
Time Period 3	-0.089 (0.068)	0.915	0.802	1.045
Time Period 4	-0.064 (0.079)	0.938	0.803	1.095
Intervention: In Person	0.259*** (0.085)	1.296	1.098	1.529
Observations	85,697			
Log Likelihood	-14,656.500			
Akaike Inf. Crit.	29,340.990			
Bayesian Inf. Crit.	29,472.010			

Abbreviations: SE, standard error; OR, odds ratio.

* $P < .1$; ** $P < .05$; *** $P < .01$.

absolute change observed did not meet the proposed absolute increase of 10% that would be clinically meaningful for the number of patients receiving the HCV screening. The number of screenings collected for this study far exceeded the requirement based on the power originally proposed. It is important to note that the practices evaluated here had previously experienced a 17% absolute increase in 2016 based on a health care system-wide EMR Baby Boomer HCV alert activation without an education intervention, that suggested that the 2016 alert activation successfully led to screening of a majority of eligible Baby Boomers before the start of this study. A previous study also showed a 20% absolute increase in HCV screening when both HCV screening education and EMR alert activation took place at select practices during 2016.¹⁹ Similarly, in support of the effectiveness of combined HCV alert activation and education, a study conducted in 2016 at the University of Kansas Health System achieved a 15% absolute increase in patients screened for HCV after HCV alert activation.²⁴ This absolute change increased to 25% when an education intervention was added shortly after.

Here, results indicated that younger Baby Boomers more likely to be screened than older Baby Boomers. There are a variety of reasons why this

might have happened and further implementation evaluation research would need to be conducted to determine why this difference occurred.

Similarly, the results indicated that females were less likely to be screened than males. There are a variety of reasons why this might have happened and further implementation evaluation research would need to be conducted to determine why this difference occurred.

The results indicate that practices with in-person presentations had higher screening rates than those who did not opt for the in-person presentation. Further research incorporating implementation evaluation would provide additional explanation around those differences. Barriers to screening, practice engagement, and other workflow issues could be evaluated to determine why these practices had difference screening rates.

The results indicate that Time Period 2 was different from the other time periods. There are a variety of reasons why this might have happened and further implementation evaluation research would need to be conducted to determine why this difference occurred.

These results and those from other studies suggest that a standalone EMR alert with additional education is effective at increasing screening rates for HCV. The education standalone intervention implemented in our health care system was likely less effective because of the previous impact of the Baby Boomer EMR alert activation.

Future Research

Because the United States Prevention Task Force is now recommending screening for all adults aged 18 to 79, future research should include the 18 to 79 age range as the variable in the logistic regression model.²⁵ To distinguish the difference in age groups, subgroups could be analyzed such as 18 to 34, 35 to 49, and 50 to 79 capturing the nuances of each age-group and their current and past behaviors.

Limitations

Because we did not track who viewed the education presentation, we do not know exactly how many physicians received the education intervention.

Because practices were not stratified based on size, clinically relevant demographic characteristics such as race, ethnicity and gender are not balanced in control and intervention conditions which may result in biased conclusions.

This study looked at screening rates before and after an education intervention which happened subsequent to the activation of the EMR alert. Significant patient populations in these practices were already screened before the education intervention which reduced the ability to measure the effect of the education intervention.

This study data were obtained from one health care system and therefore cannot be generalized to the entire population who do not seek services within our health care system.

Conclusions

This study evaluated screening rates before and after an education intervention which happened subsequent to the activation of the EMR alert. Whereas HCV EMR alerts showed an increase in HCV screenings before the education intervention, the addition of the education showed a modest additional increase in HCV screenings rates for Baby Boomer patients.

To see this article online, please go to: <http://jabfm.org/content/35/5/990.full>.

References

1. Edlin BR, Eckhardt BJ, Shu MA, et al. Toward a more accurate estimate of the prevalence of hepatitis C in the United States. *Hepatology* (Baltimore, Md.) 2015;62:1353–63.
2. Hofmeister MG, et al. Estimating prevalence of hepatitis C virus infection in the United States. *Hepatology* 2018; 2013–2016.
3. CDC. *Hepatitis C Questions and answers for health professionals*. 1/17/2019; Available from: <https://www.cdc.gov/hepatitis/hcv/hcvfaq.htm>.
4. Smith BD, et al. Recommendations for the identification of chronic hepatitis C virus infection among persons born during 1945–1965. *MMWR Recomm Rep* 2012;61:1–32.
5. Armstrong GL, Wasley A, Simard EP, et al. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med* 2006;144:705–14.
6. Rein DB, Wittenborn JS, Weinbaum CM, et al. Forecasting the morbidity and mortality associated with prevalent cases of pre-cirrhotic chronic hepatitis C in the United States. *Dig Liver Dis* 2011;43:66–72.
7. Wenger PJ, Rottnek F, Parker T, et al. Assessment of hepatitis C risk factors and infection prevalence in a jail population. *Am J Public Health* 2014;104:1722–7.

8. Ha S, Totten S, Pogany L, et al. Hepatitis C in Canada and the importance of risk-based screening. *Can Commun Dis Rep* 2016;42:57–62.
9. Akiyama MJ, Kaba F, Rosner Z, et al. Correlates of hepatitis C virus infection in the targeted testing program of the New York City jail system: epidemiologic patterns and priorities for action. *Public Health Rep* 2017;132:41–7.
10. Barth H. Hepatitis C virus: is it time to say goodbye yet? Perspectives and challenges for the next decade. *World J Hepatol* 2015;7:725–37.
11. CDC. *Hepatitis C: by the numbers*. 2020 8/17/2021]. Available from: <https://www.cdc.gov/nchhstp/newsroom/docs/factsheets/Hepatitis-c-by-the-numbers.pdf>.
12. Schillie S, W C, Osborne M, Wesolowski L, Ryerson AB. *CDC recommendations for hepatitis C screening among adults—United States, 2020*. *MMWR Recomm Rep* 2020;69:1–17.
13. El-Serag HB, Kanwal F, Richardson P, et al. Risk of hepatocellular carcinoma after sustained virological response in veterans with hepatitis C virus infection. *Hepatology* (Baltimore, Md.) 2016;64:p130–7.
14. Kanwal F, Kramer J, Asch SM, et al. Risk of hepatocellular cancer in HCV patients treated with direct-acting antiviral agents. *Gastroenterology* 2017;153:996–1005.e1.
15. Backus LI, Belperio PS, Shahoumian TA, et al. Impact of sustained virologic response with direct-acting antiviral treatment on mortality in patients with advanced liver disease. *Hepatology* 2019;69:487–97.
16. Tada T, Kumada T, Toyoda H, et al. Viral eradication reduces all-cause mortality, including non-liver-related disease, in patients with progressive hepatitis C virus-related fibrosis. *J Gastroenterol Hepatol* 2017;32:687–94.
17. Wallerstein N, Duran B. Community-based participatory research contributions to intervention research: the intersection of science and practice to improve health equity. *Am J Public Health* 2010;100 Suppl 1: S40–6.
18. Israel BA, Schulz AJ, Parker EA, Community-Campus Partnerships for Health, et al. Community-based participatory research: policy recommendations for promoting a partnership approach in health research. *Educ Health (Abingdon)* 2001; 14:182–97.
19. Tapp H, Ludden T, Shade L, et al. Electronic medical record alert activation increase hepatitis C and HIV screening rates in primary care practices within a large healthcare system. *Prev Med Rep* 2020;17:101036.
20. de Hoop E, van der Tweel I, van der Graaf R, et al. The need to balance merits and limitations from different disciplines when considering the stepped wedge cluster randomized trial design. *BMC Med Res Methodol* 2015;15:93.
21. Prost A, Binik A, Abubakar I, et al. Logistic, ethical, and political dimensions of stepped wedge trials: critical review and case studies. *Trials* 2015;16:351.
22. Ludden T, Shade L, Thomas J, et al. Novel models to identify census tracts for hepatitis C screening interventions. *J Am Board Fam Med Med* 2020;33:407–16.
23. Hooper R, Teerenstra S, de Hoop E, et al. Sample size calculation for stepped wedge and other longitudinal cluster randomised trials. *Stat Med* 2016; 35:4718–28.
24. Al-Hihi E, Shankweiler C, Stricklen D, et al. Electronic medical record alert improves HCV testing for baby boomers in primary care setting: adults born during 1945–1965. *BMJ Open Quality* 2017;6:e000084.
25. Hemker BG, Brousseau DC, Yan K, et al. When children with sickle-cell disease become adults: lack of outpatient care leads to increased use of the emergency department. *Am J Hematol* 2011; 86:863–5.