Associations Between Patient/Caregiver Trust in Clinicians and Experiences of Healthcare-Based Discrimination

Arshdeep Kaur, MSPH, Laura M. Gottlieb, MD, MPH, Stephanie Ettinger de Cuba, PhD, MPH, Elena Byhoff, MD, MSc, Eric W. Fleegler, MD, MPH, Alicia J. Cohen, MD, MSc, Nathaniel J. Glasser, MD, MPP, Mark J. Ommerborn, MPH, Cheryl R. Clark, MD, ScD, and Emilia H. De Marchis, MD, MAS

Background: Higher trust in healthcare providers has been linked to better health outcomes and satisfaction. Lower trust has been associated with healthcare-based discrimination.

Objective: Examine associations between experiences of healthcare discrimination and patients' and caregivers of pediatric patients' trust in providers, and identify factors associated with high trust, including prior experience of healthcare-based social screening.

Methods: Secondary analysis of cross-sectional study using logistic regression modeling. Sample consisted of adult patients and caregivers of pediatric patients from 11 US primary care/emergency department sites.

Results: Of 1,012 participants, low/medium trust was reported by 26% identifying as non-Hispanic Black, 23% Hispanic, 18% non-Hispanic multiple/other race, and 13% non-Hispanic White (P = .001). Experience of any healthcare-based discrimination was reported by 32% identifying as non-Hispanic Black, 23% Hispanic, 39% non-Hispanic multiple/other race, and 26% non-Hispanic White (P = .012). Participants reporting low/medium trust had a mean discrimination score of 1.65/7 versus 0.57/7 for participants reporting high trust (P < .001). In our adjusted model, higher discrimination scores were associated with lower trust in providers (aOR 0.74, 95%CI = 0.64, 0.85). A significant interaction indicated that prior healthcare-based social screening was associated with reduced impact of discrimination on trust: as discrimination score increased, odds of high trust were greater among participants who had been screened (aOR = 1.28, 95%CI = 1.03, 1.58).

Conclusions: Patients and caregivers reporting more healthcare-based discrimination were less likely to report high provider trust. Interventions to strengthen trust need structural antiracist components. Increased rapport with patients may be a potential by-product of social screening. Further research is needed on screening and trust. (J Am Board Fam Med 2024;37:607–636.)

Keywords: Caregivers, Cross-Sectional Studies, Doctor-Patient Relations, Emergency Departments, Healthcare Disparities, Health Services, Logistic Regression, Outcomes Assessment, Perceived Discrimination, Primary Healthcare, Screening, Social Determinants of Health, Trust

Background

The relationship between trust in healthcare providers and positive health behaviors is well established.^{1–18}

High levels of trust in providers has consistently been associated with adherence to treatment for a wide range of conditions, including hypertension,

(LMG); Department of Family and Community Medicine, University of California San Francisco, San Francisco, CA (LMG, EHDM); Children's HealthWatch, Boston, MA (SEDC); School of Public Health & Chobanian and Avedisian School of Medicine, Boston University, Boston, MA (SEDC); Department of Medicine, University of Massachusetts Chan Medical School, North Worcester, MA (EB);

This article was externally peer reviewed.

Submitted 12 May 2023; revised 4 March 2024; accepted 11 March 2024.

From the California University of Science and Medicine, Colton, CA (AK); Social Interventions Research and Evaluation Network, Center for Health and Community, University of California San Francisco, San Francisco, CA

diabetes, depression, and inflammatory bowel disease among adults, and asthma among children.^{1–7} High trust in providers has also been linked to increased use of preventive health services such as adolescent HPV vaccination and adult cancer screening, as well as greater patient satisfaction.^{8–18}

Critical to the healthcare system's efforts to advance health equity, previous studies have found levels of trust in providers to be lower among non-Hispanic Black and Hispanic patients, compared with non-Hispanic White patients.^{19–21} Poor selfrated health, which is similarly lower in non-Hispanic Black and Hispanic patients, has also been associated with lower trust.^{22–25} In part, these differences in trust by race and ethnicity may be attributed to experiences of healthcare discrimination; multiple studies have established links between patients' experiences of discrimination in healthcare settings and low trust in their providers.^{26–28}

Associations between trust and discrimination are particularly compelling areas of research amid

Funding: The parent study was supported by The Commonwealth Fund (CWF) (award 20181338), a national, private foundation based in New York City that supports independent research on healthcare issues and makes grants to improve healthcare practice and policy. The views presented here are those of the author and not necessarily those of CWF, its directors, officers, or staff. CWF had no role in study design; collection, analysis, or interpretation of data; writing the report; or the decision to submit the report for publication. EHD was supported by a grant from the National Research Service Award (NRSA) (T32 HP19025). EB was supported by a grant from the National Institute on Minority Health and Health Disparities (NIMHD) (K23 MD015267). NJG was supported by a grant from the Health Resources and Services Administration (HRSA) (T32 HP42019). AJC was supported by a grant from the US Department of Veterans Affairs (VA), Health Services Research and Development Service (CDA 20-037). The manuscript's contents are solely the responsibility of the authors and do not represent the official views of the CWF, NRSA, NIMHD, HRSA, or VA.

Conflict of interest: The authors have no conflicts of interest to report.

Corresponding author: Arshdeep Kaur, MSPH, California University of Science and Medicine, 1501 Violet Street, Colton, CA 92324 (E-mail: arshdeep.kaur@student.cusm. edu). increasing calls to integrate social screening (eg, screening for housing stability and food security) into care delivery, to identify and address social drivers of health and promote health equity.^{29,30} A growing body of literature explores how trust in providers may increase both patient and caregivers' perception of social screening acceptability.^{31–37} Similarly, studies have found that healthcare providers perceive that social screening may facilitate patient communication and trust.^{38–48} As healthcare-based social screening implementation expands,^{49,50} it is increasingly relevant to explore how social screening impacts trust across patient populations based on their experiences with discrimination and socioeconomic marginalization.

This study's primary aim was to examine associations between patients' and caregivers of pediatric patients' trust in healthcare providers and experiences of healthcare discrimination. A secondary aim was to identify additional factors associated with high patient/caregiver trust in providers, including self-rated health and prior experiences of healthcare-based social screening.

Methods

This study was a secondary analysis of a cross-sectional survey of adult patients and caregivers of pediatric patients from 2018 to 2019.31,51 Participants were recruited from 11 primary care and emergency department sites across 9 states, with each site recruiting convenience samples of 100 adult patients or caregivers of pediatric patients (1 site recruited 50 participants based on site capacity) (Appendix 1). Patients and caregivers were eligible for the study if they met the following criteria: 1) did not require immediate medical attention, 2) were 18 years of age or older, 3) were able to speak and read English or Spanish, 4) were able to provide informed consent, 5) were comfortable using a tablet, and 6) were themselves a patient or caregiver of a pediatric patient getting care at a study site. Recruitment and survey methods have been described in detail in previous publications.^{31,51} The original study was approved by the institutional review board of UCSF, as well as by 8 of the study sites.

The main outcome was participants' trust in their or their child's healthcare providers at the study site. Participants were asked, "How much do you trust your/your child's healthcare provider(s) at this clinic/emergency department?" Responses

Division of Emergency Medicine, Boston Children's Hospital, Boston, MA (EWF); Department of Pediatrics, Harvard Medical School, Boston, MA (EWF); Center of Innovation in Long Term Services and Supports, VA Providence Healthcare System, Providence, RI (AJC); Department of Family Medicine, Warren Alpert Medical School of Brown University (AJC); Department of Health Services, Policy, and Practice, Brown University School of Public Health (AJC); Section of General Internal Medicine, University of Chicago, Chicago, IL (NJG); Center for Community Health and Health Equity, Brigham and Women's Hospital, Boston, MA (MJO); Department of Medicine, Brigham and Women's Hospital, Boston, MA (CRC).

were measured on a scale of 1 to 10.⁵² Prior work has utilized a trichotomous cut point of 1 to 7, 8 to 9, and 10 for trust ratings; however, in this study, we aimed to explore the differences in groups with low/medium and high trust, rather than focusing on differentiating high versus complete trust.^{31,51} To that end, trust ratings were dichotomized into low/medium trust (1 to 7) and high trust (8 to 10). Caregivers of pediatric patients only responded regarding their trust in their child's healthcare provider(s).

Participants' experiences of healthcare discrimination were assessed using an adaptation of the Discrimination in Medical Settings Scale.^{53,54} Participants were asked whether they had ever experienced 7 distinct instances of discrimination in a healthcare setting due to their race, ethnicity, or socioeconomic status (eg, "felt like a doctor or nurse was not listening to what you were saying") (see Appendix 2 for the full survey). Healthcare discrimination score was defined as the number of events experienced out of the 7 queried types of discrimination. This score was treated as a continuous variable in regression analyses. In descriptive analyses, healthcare discrimination was treated as a binary variable; participants who reported 1 or more of the 7 types of discrimination were classified as having experienced any discrimination versus those who experienced none.

Exploratory secondary analyses focused on measures of self-rated health and prior experience with healthcare-based social screening/assistance. Participants rated their or their child's health status on a 5-point Likert scale; results were dichotomized into poor or fair versus good, very good, or excellent.⁵⁵ Self-perceived SES was assessed by asking participants to place themselves on a ladder relative to other people in the United States, with 1 being the bottom of the ladder and 10 being the top. This measure is designed to assess participants' subjective social status and has been validated among racially and ethnically diverse populations.^{56,57} SES ladder placement was treated as a continuous variable. To assess prior experiences with social screening/assistance, participants were asked, "In the past 12 months, not including today, have you been asked approximately any of the following in any healthcare setting?" and "In the past 12 months, not including today, have you received assistance from anyone in any healthcare setting related to..." Both questions were followed by a list of social

domains of which the participants could select multiple options (eg, housing, food).

The survey also collected participant demographics, including race, ethnicity, sex, age, years of education, income, and preferred language. Due to low sample sizes in the American Indian or Alaska Native, Asian, and other race groups, these categories were combined into a singular non-Hispanic multiple/other race category. Similarly, all participants identifying as Hispanic were collapsed into 1 category, regardless of origin (eg, Mexican American). The race and ethnicity variable in this study was used as a proxy for experiences of interpersonal and institutional racism.⁵⁸ In addition to participant-level characteristics, 2 site-level characteristics were included in the analysis: type of healthcare setting (primary care vs emergency department) and percentage of the patient population who were publicly insured/uninsured (dichotomized into less than 80% vs 80% or more).^{31,51} Aside from these 2 site-level measures, all measures were self-reported by participants.

Descriptive analyses using Chi Square tests were used to explore level of trust in providers and experiences of healthcare discrimination among the study sample. Univariable and multivariable logistic regressions were then used to examine associations between level of trust in providers and healthcare discrimination score, health status, having been screened for social risks in a healthcare setting in the past 12 months, and having been assisted with social needs in a healthcare setting in the past 12 months, while controlling for the following participant- and site-level characteristics: sex, race and ethnicity, age, education, income, SES ladder position, health status, preferred language, healthcare setting (primary care vs ED), percentage publicly insured/ uninsured at site, and type of participant (adult patient vs caregiver of a pediatric patient), as well as cluster analysis at the site level.^{12,24,26,27,31,51,59-61} Regression analyses utilized listwise deletion. We did not use multiple imputations because of the likelihood that data were not missing at random, but instead were missing due to the sensitive nature of the data (eg, self-reported income).⁶²⁻⁶⁵ Descriptive statistics for the disaggregated Hispanic group were calculated and provided in Appendix 3, as consistent with best practices in the reporting of race and ethnicity data despite small sample sizes.^{58,66} Unfortunately, this was not possible for the non-Hispanic multiple/other race group, due to very low sample sizes and concerns about participant confidentiality.

We used interaction terms to explore statistical interactions between healthcare discrimination score and other participant-level characteristics (demographics, health status, having been screened for social risks in a healthcare setting in the past 12 months, having been assisted with social needs in a healthcare setting in the past 12 months). These analyses aimed to explore potential differences in the relationship between healthcare discrimination and trust, given the known differences in levels of trust among different populations, the broad nature of the Everyday Discrimination Scale (with participants reporting discrimination based on race, ethnicity, or socioeconomic status), and gaps in knowledge regarding social screening among diverse populations.^{19–21,24,25,67} Results were stratified by race and ethnicity. Model goodness-of-fit was assessed using R-squared and Akaike information criterion values; models were also evaluated for multicollinearity and specification error. Sensitivity analyses were performed to examine the effect of using a tertile-based cut point (1 to 7, 8 to 9, and 10) for the provider trust outcome. Because of the overall high ratings of trust among the sample, additional sensitivity analyses using a dichotomous 1 to 9 and 10 cut point, as well as a National Quality Forum-recommended 1 to 8 and 9 to 10 cutpoint, were also performed.⁶⁸ In addition, we stratified by participant type (adult patient vs caregiver of pediatric patient) to evaluate for variations in participant report of trust in their own providers compared with trust in a child's provider. Similarly, we stratified by healthcare setting to evaluate for differences in trust in providers based on participants being recruited from primary care versus emergency department sites. Data collection took place between July 2018 and February 2019, and analyses were completed between June 2021 and November 2022. All analyses were conducted using Stata 15.1.69

Results

Of the 1,771 adult patients and adult caregivers of pediatric patients who were approached for recruitment, 1,090 provided consent and initiated the survey. The final sample (n = 1,012) consisted of those who completed \geq 50% of the survey questions, including the question about level of trust in providers

(Appendix 1). 77% (n = 781) of participants were adult patients (Table 1). Participants identifying as non-Hispanic White made up the largest racial and ethnic group in the sample (n = 357, 37%), followed by Hispanic (n = 310, 33%), non-Hispanic Black (n = 207, 22%), and non-Hispanic multiple/other race (n = 83, 9%) participants. 77% (n = 755) of participants rated themselves or their child as being in good, very good, or excellent health.

Twenty percent (n = 197) of participants were in the low/medium trust group. Participants had a median trust rating of 10/10 (interquartile range = 8 to 10). Those reporting low/medium trust in their or their child's providers had a mean healthcare discrimination score of 1.65/7, compared with 0.57/7 for participants reporting high trust (P < .001) (Table 1). Participants identifying as non-Hispanic Black (n = 54, 26%) reported the highest percentage of low/medium trust in providers (P < .001) (Table 2). The highest percentage of any reported healthcare discrimination was among participants identifying as non-Hispanic multiple/other race (n = 32, 39%) (P = .012). Significant differences were noted among racial and ethnic groups when comparing each of the 7 types of healthcare discrimination experienced (Table 2). Participants identifying as non-Hispanic Black and non-Hispanic multiple/ other race reported higher rates of experiencing each type of discrimination compared with non-Hispanic White and Hispanic participants. Across racial and ethnic groups, the most common type of discrimination reported was "felt like a doctor or nurse was not listening to what [they] were saying."

In the multivariable logistic regression model, participants with higher discrimination scores were less likely to report high trust in their providers (adjusted OR [aOR] 0.74, 95% CI 0.64, 0.85), whereas participants with good, very good, or excellent self-rated health were more likely to report high trust in providers (aOR 3.15, 95% CI 2.01, 4.92) (Table 3). The interaction term between healthcare discrimination score and having been screened for social risks in the past was statistically significant. Increased healthcare-based discrimination was associated with lower odds of high trust regardless of whether the participant had been screened for social risks. However, there was a difference in magnitude - the decrease in odds of high trust was smaller for those who had been previously

Þ	
Ę	
Во	
arc	
am	
2	
ed	
 fi	
ß	
р	
b	
she	
ð	
as	
10	
ω	
12	
2/3	
ðf	
3	
20	
23	
i,	
ő	
8	
R	
~	
Ĕ	
25	
Q)
융	
ğ	
õ	
- N 1	
24.	
24. D	
24. Dow	
24. Downlo	
24. Downloac	
24. Downloadec	
24. Downloaded fr	
24. Downloaded from	
24. Downloaded from h	
24. Downloaded from http:	
24. Downloaded from http://v	
24. Downloaded from http://ww	
24. Downloaded from http://www.j	
24. Downloaded from http://www.jab	
24. Downloaded from http://www.jabfm.	
24. Downloaded from http://www.jabfm.org	
24. Downloaded from http://www.jabfm.org/ o	
24. Downloaded from http://www.jabfm.org/ on	
24. Downloaded from http://www.jabfm.org/ on 6 l	
24. Downloaded from http://www.jabfm.org/ on 6 Ma	
24. Downloaded from http://www.jabfm.org/ on 6 May 2	
24. Downloaded from http://www.jabfm.org/ on 6 May 202	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 i	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by gu	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by gues	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. F	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Prc	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protec	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protecte	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protected t	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protected by o	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protected by cop	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protected by copyr	
24. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protected by copyrigh	

Table 1. S	Sample	Characteristics	by	Level	of Trust	in	Healthcare Provider	
------------	--------	-----------------	----	-------	----------	----	----------------------------	--

	n (%)			
		Low/Medium Trust (1–7)	High Trust (8–10)	
	Total	(n = 197)	(n = 815)	P-value
Biological sex ($n = 998$)				
Female	701 (70)	136 (19)	565 (81)	0.96
Male	297 (30)	58 (20)	239 (80)	
Race and ethnicity $(n = 957)$				
Non-Hispanic White	357 (37)	48 (13)	309 (87)	0.001
Non-Hispanic Black	207 (22)	54 (26)	153 (74)	
Hispanic	310 (32)	70 (23)	240 (77)	
Non-Hispanic multiple/other	83 (9)	15 (18)	68 (82)	
Age $(n = 1.003)$				
18 to 44	541 (54)	116 (21)	425 (79)	0.25
45 to 64	301 (30)	51 (17)	250 (83)	
65+	161 (16)	29 (18)	132 (82)	
Years of education $(n = 1.005)$				
Less than 12 years	170 (17)	37 (22)	133 (78)	0.44
12 or more years	835 (83)	160 (19)	675 (81)	
Income $(n = 857)$				
\$0-50k	595 (69)	136 (23)	459 (77)	< 0.001
\$50.001–75k	82 (10)	7 (9)	75 (92)	20.001
\$75.001+	180 (21)	23 (13)	157 (87)	
SFS ladder position* $(n = 922)$	5 66 (2 23)	5.06(2.11)	5 80 (2 23)	< 0.001
Healthcare setting $(n = 1.012)$	5.00 (2.25)	5.00 (2.11)	5.00 (2.25)	20.001
Primary care	620 (61)	118 (19)	502 (81)	0.66
Emergency department	392 (39)	79 (20)	313 (80)	0.000
Site type $(n = 1.012)$	572 (57)	() (20)	515 (00)	
Urban academic	720 (71)	146 (20)	574 (80)	0.56
Urban non-academic	97 (10)	18 (19)	79 (81)	
Non-urban non-academic	195 (19)	33 (17)	162 (83)	
Percentage publicly insured/uninsured at site $(n = 1.012)$			()	
Less than 80% publicly insured/uninsured	725 (72)	132 (18)	593 (82)	0.11
80% or more publicly insured/uninsured	287 (28)	65 (23)	222 (77)	
Participant type $(n = 1.012)$			(/ / /	
Adult patient	781 (77)	151 (19)	630 (81)	0.85
Caregiver of pediatric patient	231 (23)	46 (20)	185 (80)	0.05
Preferred language ($n = 1.012$)				
English	845 (83)	167 (20)	678 (80)	0.59
Spanish	167 (17)	30 (18)	137 (82)	0.077
Patient/caregiver-reported health ($n = 985$)		()		
Poor. Fair	230 (23)	69 (30)	161 (70)	< 0.001
Good. Very Good. Excellent	755 (77)	119 (16)	636 (84)	20.001
Experienced any of 7 categories perceived discrimination (n = 1,002)		/ (/)		
Yes	275 (27)	98 (36)	177 (64)	< 0.001
No	727 (73)	96 (13)	631 (87)	
Healthcare discrimination score $*(n = 927)$	0.78 (1.65)	1.65 (2.15)	0.57 (1.43)	< 0.001
Asked about needs in any domains in the past 12 months ($n = 993$)	()	/	()	
Yes	307 (31)	63 (21)	244 (80)	0.60
No	686 (69)	131 (19)	555 (81)	

Continued

		n (%)		
	Total	Low/Medium Trust (1–7) (n = 197)	High Trust (8–10) (n = 815)	<i>P</i> -value
Assisted with needs in any domains in the past 12 months $(n = 993)$				
Yes	174 (18)	48 (28)	126 (72)	0.003
No	819 (82)	145 (18)	674 (82)	

*mean (S.D.).

Abbreviation: SES, socioeconomic status.

screened (26% vs 6%, P=.026). In other words, with each 1-point increase in healthcare discrimination score (a higher level of reported discrimination), participants who had not been screened for social risks in a healthcare setting in the prior 12 months experienced a 26% lower odds of high trust (greater odds of low trust), whereas those who had been screened experienced 6% lower odds of high trust. Prior social screening by itself was not associated with trust.

In analyses stratified by race and ethnicity, the trends in associations between trust and discrimination, as well as between trust and self-rated health, were similar across groups (Table 4). In sensitivity analyses examining alternative cut points of the trust outcome, as well as stratification by type of participant (adult patient vs caregiver of pediatric patient) and healthcare setting (primary care vs emergency department), results showed consistent trends in direction and magnitude of associations (Appendix 3 Tables 1–8). Disaggregated descriptive statistics for Hispanic participants are displayed in Appendix 3 Tables 9 and 10.

Discussion

In this study, we found that as patients and caregivers reported more types of healthcare-based discrimination, they were less likely to report high trust in their or their child's providers. This finding is consistent with prior work on the relationship between healthcare discrimination and trust.^{26,27}

indicate and the first with high should be interested of first minimum of the ball of the	Table 2.	Trust in Healthcare	Provider and Exp	periences of Heal	thcare Discrimination	by Race and Ethnicit
---	----------	----------------------------	-------------------------	-------------------	-----------------------	----------------------

		n (%)					
	Total	Non-Hispanic White (n = 367)	Non-Hispanic Black (n = 212)	Hispanic (n = 329)	Non-Hispanic Multiple/ Other (n = 84)	<i>P</i> -value	
Low/medium trust (1 to 7)	187 (20)	48 (13)	54 (26)	70 (23)	15 (18)	0.001	
High trust (8 to 10)	770 (81)	309 (87)	153 (74)	240 (77)	68 (82)		
Experienced any of 7 categories perceived discrimination	268 (27)	94 (26)	67 (32)	75 (23)	32 (38)	0.01	
Item 1: Felt like a doctor or nurse was not listening to what you were saying	183 (19)	54 (15)	47 (23)	59 (18)	23 (27)	0.02	
Item 2: Treated you with less respect than other people	123 (13)	40 (11)	38 (18)	28 (9)	17 (20)	0.001	
Item 3: Received poorer services than other people	93 (10)	25 (7)	34 (17)	20 (6)	14 (17)	< 0.001	
Item 4: Treated with less courtesy than other people	107 (11)	34 (9)	28 (14)	29 (9)	16 (19)	0.03	
Item 5: Had a doctor or nurse act as if he or she was better than you	117 (12)	50 (14)	30 (15)	21 (7)	16 (20)	0.001	
Item 6: Had a doctor or nurse act as if he or she thinks you were not smart	122 (13)	52 (14)	33 (16)	24 (7)	13 (16)	0.006	
Item 7: Had a doctor or nurse act as if he or she was afraid of you	31 (3)	6 (2)	16 (8)	3 (1)	6 (7)	<0.001	

Table 3. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest (n = 690)

	Adjusted Odds Ratio (95% CI)
Healthcare discrimination score	0.74 (0.64, 0.85)
Race and ethnicity	
Non-Hispanic White (reference)	
Non-Hispanic Black	0.56 (0.27, 1.17)
Hispanic	0.67 (0.30, 1.52)
Non-Hispanic multiple/other	0.96 (0.39, 2.72)
Age	
18 to 44 (reference)	
45 to 64	1.30 (0.82, 2.06)
65+	1.05 (0.39, 2.81)
Education 12 or more years	0.62 (0.36, 1.05)
Income	
\$0–50k (reference)	
\$50,001–75k	2.19 (0.82, 5.84)
\$75,001+	1.41 (0.90, 2.22)
SES ladder	1.04 (0.92, 1.16)
Self-rated health good, very good, or excellent	3.15 (2.01, 4.92)
Interaction: Self-rated health * discrimination score	0.84 (0.74, 0.96)
80% or more publicly insured/uninsured	0.76 (0.52, 1.12)
Previously screened for social risks in prior 12 months	0.99 (0.63, 1.55)
Interaction: Previously screened * discrimination score	1.28 (1.03, 1.58)
Previously assisted with social risks in prior 12 months	0.86 (0.60, 1.23)

Abbreviations: SES, socioeconomic status; CI, confidence interval.

We additionally found evidence that the negative association between discrimination and trust was different among patients who had previously received social risk screening, based on a significant interaction term coefficient.

Overall, trust in our sample was high. Although it is difficult to make direct comparisons to the level of trust found in other similar samples given variability in measures of patient trust reported in the literature, qualitative assessment of these studies suggests that our findings are substantively similar to what is published. Level of trust in healthcare providers differed significantly between racial and ethnic groups, with participants identifying as non-Hispanic Black or Hispanic reporting higher rates of low trust in their own or their child's healthcare providers. Reporting healthcare discrimination was common in our sample; more than a quarter of participants reported at least 1 experience of discrimination. Previous estimates from studies with similarly diverse samples have ranged from 6% to 21%.^{70,71} We also found significant differences between racial and ethnic groups in terms of reporting experiences of healthcare discrimination, with participants identifying as non-Hispanic multiple/ other race or non-Hispanic Black reporting higher levels of discrimination compared to the non-Hispanic White participants.

Prior research has demonstrated that trust in providers plays an important role in treatment adherence, use of preventive healthcare, and patient satisfaction.^{1-6,8-16} The responsibility to build trust falls on individual providers, care teams, and healthcare systems, and there must be comprehensive system-wide efforts to improve trustworthiness.⁷² Given our finding that healthcare discrimination is associated with low trust, antiracist policies and programs should be evaluated as ways to improve trust alongside preventing and addressing healthcare discrimination. Many institutions have implemented individuallevel interventions such as trainings in antiracism and cultural competence, however, multiple systematic reviews have found little to no evidence to support the idea that these programs impact healthcare workers' behavior or patient outcomes.73-75 Studies show that racial and ethnic concordance between patients and providers is associated with better communication,^{76–78} as well as increased use of preventive care and decreased emergency department visits.^{79,80} Recent research, including qualitative studies of Black women's reproductive health, has demonstrated the role of patient-provider racial concordance in facilitating trust and combating the effects of racism in healthcare.^{81,82} Thus, the recruitment and retention of more diverse providers may be 1 strategy for improving minoritized patients' experiences and outcomes. This is the rationale for educational pipeline programs focused on diversity, for instance. Per Calliste and Dei, antiracism is "an action-oriented, educational and/or political strategy for systemic and political change that addresses issues of racism and interlocking systems of social oppression"; however, existing research is heavily weighted toward individual-level efforts.83 Evaluation of antiracism efforts should include both process and impact measures, at individual, care team, and organizational levels. In addition, evaluation should examine not only discrimination, but also associated factors, such as trust and health outcomes.

	Adjusted Odds Ratio (95% CI)					
	Non-Hispanic White (n = 275)	Non-Hispanic Black (n = 161)	Hispanic (n = 196)	Non-Hispanic Multiple/Other (n = 48)		
Healthcare discrimination score	0.80 (0.63, 1.00)	0.71 (0.53, 0.96)	0.52 (0.40, 0.68)	0.57 (0.24, 1.37)		
Age						
18 to 44 (reference)						
45 to 64	1.27 (0.72, 2.22)	1.21 (0.46, 3.18)	0.85 (0.23, 3.19)	7.63 (0.67, 87.08)		
65+	1.21 (0.65, 2.25)	1.18 (0.15, 9.26)	0.71 (0.27, 1.88)	0.42 (0.01, 28.58)		
Education 12 or more years	0.90 (0.12, 7.12)	1.72 (0.80, 3.68)	0.15 (0.05, 0.42)	0.70 (0.04, 12.82)		
Income						
\$0–50k(reference)						
\$50,001–75k	2.48 (0.45, 13.69)	2.30 (1.01, 5.27)	1.06 (0.30, 3.78)	-		
\$75,001+	1.24 (0.54, 2.81)	2.12 (0.56, 8.07)	0.55 (0.12, 2.50)	2.07 (0.10, 40.87)		
SES ladder	1.07 (0.87, 1.33)	1.01 (0.82, 1.26)	1.00 (0.82, 1.24)	0.63 (0.23, 1.67)		
Self-rated health good/very good/excellent	4.77 (2.30, 9.88)	1.59 (0.78, 3.26)	5.23 (1.99, 13.74)	12.75 (0.16, 1020.04)		
Interaction: Self-rated health *discrimination score	0.88 (0.65, 1.17)	0.97 (0.74, 1.26)	0.31 (0.14, 0.72)	0.79 (0.43, 1.42)		
80% or more publicly insured/uninsured	1.25 (0.21, 7.41)	1.24 (0.52, 2.95)	0.32 (0.11, 0.95)	0.84 (0.12, 5.80)		
Previously screened for social risks in prior 12 months	1.38 (0.53, 3.53)	1.28 (0.56, 2.92)	0.77 (0.27, 2.19)	0.21 (0.01, 4.54)		
Interaction: Previously screened* discrimination score	1.36 (0.86, 2.15)	1.03 (0.79, 1.35)	4.65 (1.92, 11.29)	1.49 (0.53, 4.24)		
Previously assisted with social risks in prior 12 months	1.54 (0.47, 5.02)	0.73 (0.35, 1.53)	1.03 (0.27, 3.99)	0.26 (0.01, 5.53)		

Table 4. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest, Stratified by Race and Ethnicity

Abbreviations: SES, socioeconomic status; CI, confidence interval.

Study findings also suggest that being screened for social risks in healthcare settings may help build rapport and trust with patients' healthcare teams, thus reducing the impact of prior discrimination. These findings align with the results of several recent studies of social resource interventions, in which screening and assistance with social needs was associated with more trust in, and better relationships with, providers.⁸⁴⁻⁸⁶ Previously published findings from the parent study from which our data were derived also reported that patients/caregivers perceived value in screening for the purpose of making healthcare providers aware of their social context and did not expect the healthcare team to resolve their social needs.³² Similarly, in a separate study of adult patients who were screened for social risks, 43% wanted their healthcare team to be aware of their screening results, but did not expect assistance.⁸⁷ This supports the idea that even in the absence of interventions, social screening may help to build trust, especially when delivered by personnel trained in empathic inquiry and trauma-informed approaches to screening.36 Universal social screening may provide

opportunities to strengthen patient-provider relationships, beyond the goal of reducing social needs.⁸⁸ It is notable that our dataset did not offer opportunities to examine more nuanced associations between trust and the context of social screening, however. It is possible that screening may exacerbate feelings of healthcare discrimination and/or worsen mistrust if not accompanied by adequate assistance, or if screening itself causes patients to feel stigmatized.⁸⁹ Perceived failures in the process of providing assistance could also affect patient/caregiver willingness to engage with future screening and/or assistance programs. Future research might explore best practices for social screening across different patient populations.⁶⁷ This research should prioritize examining both the intended and unintended impacts of screening, especially as it relates to trust and health equity. Factors such as patient-provider racial concordance, culture, nativity, and language may affect patients' receptivity to answering screening questions.^{77,90}

This study has several limitations. First, due to the cross-sectional study design, we were not able to assess the temporality of participants' experiences of healthcare discrimination, or if/how their trust in providers changed in relationship to experiences of discrimination. Participants' report of discrimination could have been influenced by their previous levels of trust. Although participants were asked about their trust in providers at the study site where they were completing the study survey, they were asked to report any prior discrimination across healthcare settings more generally. Regardless of setting, the survey did not assess care continuity. Care continuity should be assessed in future research as it may influence patients' trust in their healthcare team or organization, as well as moderate/mediate the relationship between trust and self-reported experiences of discrimination.^{91,92} Similarly, we cannot establish temporality between experience of discrimination and screening for social risks. The healthcare setting at which participants were surveyed and asked about their trust levels may not have been the same setting in which they were screened for social risks. Study findings cannot be used to determine causation. Second, the study is subject to selection bias, given the convenience sample design, as well as social desirability bias. Dropout due to missing data may also have biased our study, likely away from the null. In addition, although we included a diverse set of primary care and emergency department study sites, findings may not be generalizable across settings. The use of a single aggregated Hispanic category and a single non-Hispanic multiple/other race category due to small sample sizes also limits generalizability and may mask differing levels of trust and differing experiences of discrimination within these groups. Participants were also limited to those who could speak and read English and/or Spanish. Finally, our study survey did not capture all known factors anticipated to influence trust and discrimination, for example, racial concordance between patient and provider team and care continuity. Future research should consider incorporating a wider set of key variables that can be used to more carefully understand and improve experiences of social care. Despite these limitations, study findings deepen our understanding of factors that may influence patients' and caregivers' trust in providers, as well as provide directions for future research.

Conclusions

Experiences of healthcare-based discrimination were associated with low trust in healthcare providers. These findings underscore the need for systemic antiracist interventions to prevent and address healthcare discrimination and improve healthcare system trustworthiness. In addition, a potential by-product of social screening in healthcare settings may be building rapport with patients, possibly reducing the impact of prior discrimination on trust. More research is needed on the relationship between social screening and trust, particularly as social screening programs are implemented more widely.

We thank the study sites and participants for sharing their time and experiences.

To see this article online, please go to: http://jabfm.org/content/ 37/4/607.full.

References

- Abel W, Efird J. The association between trust in health care providers and medication adherence among Black women with hypertension. Front Public Health 2013;1:66. Accessed April 15, 2022. Available at: https://www.frontiersin.org/article/ 10.3389/fpubh.2013.00066.
- Schoenthaler A, Montague E, Manwell LB, Brown R, Schwartz MD, Linzer M. Patient-physician racial/ethnic concordance and blood pressure control: the role of trust and medication adherence. Ethn Health 2014;19:565–78.
- Reach G, Pellan M, Crine A, Touboul C, Ciocca A, Djoudi Y. Holistic psychosocial determinants of adherence to medication in people with type 2 diabetes. Diabetes Metab 2018;44:500–7.
- Bonds DE, Camacho F, Bell RA, Duren-Winfield VT, Anderson RT, Goff DC. The association of patient trust and self-care among patients with diabetes mellitus. BMC Fam Pract 2004;5:26.
- Bauer AM, Parker MM, Schillinger D, et al. Associations between antidepressant adherence and shared decision-making, patient-provider trust, and communication among adults with diabetes: Diabetes Study of Northern California (DISTANCE). J Gen Intern Med 2014;29:1139–47.
- Nguyen GC, LaVeist TA, Harris ML, Datta LW, Bayless TM, Brant SR. Patient trust-in-physician and race are predictors of adherence to medical management in inflammatory bowel disease. Inflamm Bowel Dis 2009;15:1233–9.
- Rotenberg KJ, Petrocchi S. A longitudinal investigation of trust beliefs in physicians by children with asthma and their mothers: relations with children's adherence to medical regimes and quality of life. Child Care Health Dev 2018;44:879–84.
- Gupta S, Brenner AT, Ratanawongsa N, Inadomi JM. Patient trust in physician influences colorectal cancer screening in low-income patients. Am J Prev Med 2014;47:417–23.

- Liu Y, Zupan NJ, Shiyanbola OO, et al. Factors influencing patient adherence with diabetic eye screening in rural communities: a qualitative study. PLOS ONE 2018;13:e0206742.
- Musa D, Schulz R, Harris R, Silverman M, Thomas SB. Trust in the health care system and the use of preventive health services by older Black and white adults. Am J Public Health 2009;99:1293–9.
- Hong HC, Ferrans CE, Park C, Lee H, Quinn L, Collins EG. Effects of perceived discrimination and trust on breast cancer screening among Korean American women. Womens Health Issues 2018;28:188–96.
- O'Malley AS, Sheppard VB, Schwartz M, Mandelblatt J. The role of trust in use of preventive services among low-income African-American women. Prev Med 2004;38:777–85.
- 13. Lee YY, Lin JL. How much does trust really matter? A study of the longitudinal effects of trust and decision-making preferences on diabetic patient outcomes. Patient Educ Couns 2011;85:406–12.
- Platonova EA, Kennedy KN, Shewchuk RM. Understanding patient satisfaction, trust, and loyalty to primary care physicians. Med Care Res Rev MCRR 2008;65:696–712.
- Thom DH, Kravitz RL, Bell RA, Krupat E, Azari R. Patient trust in the physician: relationship to patient requests. Fam Pract 2002;19:476–83.
- Baker R, Mainous AG, Gray DP, Love MM. Exploration of the relationship between continuity, trust in regular doctors and patient satisfaction with consultations with family doctors. Scand J Prim Health Care 2003;21:27–32.
- Fu LY, Zimet GD, Latkin CA, Joseph JG. Associations of trust and healthcare provider advice with HPV vaccine acceptance among African American parents. Vaccine 2017;35:802–7.
- Fuzzell LN, LaJoie AS, Smith KT, Philpott SE, Jones KM, Politi MC. Parents' adherence to pediatric health and safety guidelines: importance of patient-provider relationships. Patient Educ Couns 2018;101:1570–6.
- Sewell AA. Disaggregating ethnoracial disparities in physician trust. Soc Sci Res 2015;54:1–20.
- Halbert CH, Armstrong K, Gandy OH, Shaker L. Racial differences in trust in health care providers. Arch Intern Med 2006;166:896–901.
- Doescher MP, Saver BG, Franks P, Fiscella K. Racial and ethnic disparities in perceptions of physician style and trust. Arch Fam Med 2000;9:1156–63.
- Borrell LN, Dallo FJ. Self-rated health and race among Hispanic and non-Hispanic adults. J Immigr Minor Health 2008;10:229–38.
- Liang J, Quiñones AR, Bennett JM, et al. Evolving self-rated health in middle and old age: how does it differ across Black, Hispanic, and White Americans? J Aging Health 2010;22:3–26.

- 24. Mohseni M, Lindstrom M. Social capital, trust in the health-care system and self-rated health: the role of access to health care in a population-based study. Soc Sci Med 1982 2007;64:1373–83.
- Kim AM, Bae J, Kang S, Kim YY, Lee JS. Patient factors that affect trust in physicians: a cross-sectional study. BMC Fam Pract 2018;19:187.
- Bazargan M, Cobb S, Assari S. Discrimination and medical mistrust in a racially and ethnically diverse sample of California adults. Ann Fam Med 2021;19:4–15.
- Hausmann LRM, Kwoh CK, Hannon MJ, Ibrahim SA. Perceived racial discrimination in health care and race differences in physician trust. Race Soc Probl 2013;5:113–20.
- Adegbembo AO, Tomar SL, Logan HL. Perception of racism explains the difference between Blacks' and Whites' level of healthcare trust. Ethn Dis 2006; 16:792–8.
- 29. National Academies of Sciences, Engineering, and Medicine. Integrating social care into the delivery of health care: moving upstream to improve the nation's health. Washington (DC): National Academies Press (US); 2019.
- Gottlieb LM, Lindau ST, Peek ME. Why add "abolition" to the National Academies of Sciences, Engineering, and Medicine's Social Care Framework? AMA J Ethics 2022;24:E170–180.
- De Marchis EH, Hessler D, Fichtenberg C, et al. Part I: a quantitative study of social risk screening acceptability in patients and caregivers. Am J Prev Med 2019;57:S25–S37.
- Byhoff E, De Marchis EH, Hessler D, et al. Part II: a qualitative study of social risk screening acceptability in patients and caregivers. Am J Prev Med 2019;57:S38–S46.
- 33. Byhoff E, De Marchis EH, Gottlieb L, Halperin-Goldstein S, Nokes K, LeClair AM. Screening for immigration-related health concerns in a federally qualified health center serving a diverse Latinx community: a mixed methods study. J Immigr Minor Health 2020;22:988–95.
- Wallace AS, Luther BL, Sisler SM, Wong B, Guo JW. Integrating social determinants of health screening and referral during routine emergency department care: evaluation of reach and implementation challenges. Implement Sci Commun 2021;2:114.
- Drake C, Batchelder H, Lian T, et al. Implementation of social needs screening in primary care: a qualitative study using the health equity implementation framework. BMC Health Serv Res 2021;21:975.
- 36. Broaddus-Shea ET, Fife Duarte K, Jantz K, Reno J, Connelly L, Nederveld A. Implementing healthrelated social needs screening in western Colorado primary care practices: qualitative research to inform improved communication with patients. Health Soc

Care Community 2022;30:e3075–e3085. Published online February 16.

- Spain AK, Monahan EK, Alvarez K, Finno-Velasquez M. Latinx family perspectives on social needs screening and referral during well-child visits. MCN Am J Matern Nurs 2021;46.
- Eismann EA, Theuerling J, Maguire S, Hente EA, Shapiro RA. Integration of the Safe Environment for Every Kid (SEEK) model across primary care settings. Clin Pediatr (Phila) 2019;58:166–76.
- Freibott CE, Beaudin E, Frazier BJ, Dias A, Cooper MR. Toward successful and sustainable statewide screening for social determinants of health: testing the interest of hospitals. Popul Health Manag 2021;24:567–75.
- Hamity C, Jackson A, Peralta L, Bellows J. Perceptions and experience of patients, staff, and clinicians with social needs assessment. Perm J 2018;22:18–105.
- O'Toole TP, Roberts CB, Johnson EE. Screening for food insecurity in six Veterans Administration clinics for the homeless, June-December 2015. Prev Chronic Dis 2017;14:E04.
- 42. Onyekere C, Ross S, Namba A, Ross JC, Mann BD. Medical student volunteerism addresses patients' social needs: a novel approach to patient-centered care. Ochsner J 2016;16:45–9.
- Palakshappa D, Vasan A, Khan S, Seifu L, Feudtner C, Fiks AG. Clinicians' perceptions of screening for food insecurity in suburban pediatric practice. Pediatrics 2017;140:e20170319.
- Tong ST, Liaw WR, Kashiri PL, et al. Clinician experiences with screening for social needs in primary care. J Am Board Fam Med 2018;31:351–63.
- 45. Adams E, Hargunani D, Hoffmann L, Blaschke G, Helm J, Koehler A. Screening for food insecurity in pediatric primary care: a clinic's positive implementation experiences. J Health Care Poor Underserved 2017;28:24–9.
- Swamy P, Monterrey AC, Wood MS, Troisi CL, Greeley CS. Caregiver and pediatric health care provider views on social needs identification. J Prim Care Community Health 2020;11:2150132720923085.
- 47. Kostelanetz S, Pettapiece-Phillips M, Weems J, et al. Health care professionals' perspectives on universal screening of social determinants of health: a mixed-methods study. Popul Health Manag 2022; 25:367–74.
- Leary JC, Rijhwani L, Bettez NM, et al. Clinical stakeholder perspectives on pediatric inpatient screening for social needs. Acad Pediatr 2022;22:470–7.
- Centers for Medicare and Medicaid Services. 87 FR 48780. Published online August 10, 2022. Accessed November 24, 2022. Available at: https://www. federalregister.gov/documents/2022/08/10/2022-16472/medicare-program-hospital-inpatient-prospectivepayment-systems-for-acute-care-hospitals-and-the.

- National Committee for Quality Assurance. HEDIS MY 2023 measure descriptions. Accessed November 29, 2022. Available at: https://www.ncqa.org/wpcontent/uploads/2022/07/HEDIS-MY-2023-Measure-Description.pdf.
- 51. De Marchis EH, Hessler D, Fichtenberg C, et al. Assessment of social risk factors and interest in receiving health care–based social assistance among adult patients and adult caregivers of pediatric patients. JAMA Netw Open 2020;3:e2021201.
- 52. Agency for Healthcare Research and Quality. CAHPS clinician & group surveys supplemental items for the adult surveys 2.0. Published online 2012. Accessed August 3, 2022. Available at: https:// archive.ahrq.gov/cahps/surveys-guidance/item-sets/ cultural/2357a_Adult_Supp_Eng_20.pdf.
- Bird ST, Bogart LM. Perceived race-based and socioeconomic status(SES)-based discrimination in interactions with health care providers. Ethn Dis 2001;11:554–63.
- Peek ME, Nunez-Smith M, Drum M, Lewis TT. Adapting the everyday discrimination scale to medical settings: reliability and validity testing in a sample of African American patients. Ethn Dis 2011;21:502–9.
- DeSalvo KB, Fan VS, McDonell MB, Fihn SD. Predicting mortality and healthcare utilization with a single question. Health Serv Res 2005;40:1234–46.
- 56. Adler NE, Epel ES, Castellazzo G, Ickovics JR. Relationship of subjective and objective social status with psychological and physiological functioning: preliminary data in healthy white women. Health Psychol Off Psychol 2000;19:586–92.
- Operario D, Adler NE, Williams DR. Subjective social status: reliability and predictive utility for global health. Psychol Health 2004;19:237–46.
- Lett E, Asabor E, Beltrán S, Cannon AM, Arah OA. Conceptualizing, contextualizing, and operationalizing race in quantitative health sciences research. Ann Fam Med 2022;20:157–63.
- Fields A, Abraham M, Gaughan J, Haines C, Hoehn KS. Language matters: race, trust, and outcomes in the pediatric emergency department. Pediatr Emerg Care 2016;32:222–6.
- Armstrong K, Ravenell KL, McMurphy S, Putt M. Racial/ethnic differences in physician distrust in the United States. Am J Public Health 2007;97:1283–9.
- Wiltshire JC, Person SD, Allison J. Exploring differences in trust in doctors among African American men and women. J Natl Med Assoc 2011;103:845–51.
- 62. Giusti C, Little RJA. An analysis of nonignorable nonresponse to income in a survey with a rotating panel design. J Off Stat 2011;27:211–29.
- King G, Honaker J, Joseph A, Scheve K. Analyzing incomplete political science data: an alternative algorithm for multiple imputation. Am Polit Sci Rev 2001;95:49–69.

- 64. Adjaye-Gbewonyo D, Bednarczyk RA, Davis RL, Omer SB. Using the Bayesian Improved Surname Geocoding Method (BISG) to create a working classification of race and ethnicity in a diverse managed care population: a validation study. Health Serv Res 2014;49:268–83.
- 65. Grundmeier RW, Song L, Ramos MJ, et al. Imputing missing race/ethnicity in pediatric electronic health records: reducing bias with use of U.S. census location and surname data. Health Serv Res 2015;50:946–60.
- 66. Ross PT, Hart-Johnson T, Santen SA, Zaidi NLB. Considerations for using race and ethnicity as quantitative variables in medical education research. Perspect Med Educ 2020;9:318–23.
- De Marchis EH, Brown E, Aceves B, et al. State of the science of screening in healthcare settings. Published online 2022;140. Available at: https:// sirenetwork.ucsf.edu/sites/default/files/2022-06/ final%20SCREEN%20State-of-Science-Report% 5B55%5D.pdf.
- National Quality Forum. Healthcare disparities and cultural competency consensus standards technical report. Published 2012. Available at: https://www. qualityforum.org/Publications/2012/09/Healthcare_ Disparities_Technical_Report.aspx.
- 69. StataCorp. Stata. Published online 2017. Available at: https://www.stata.com.
- Alcalá HE, Cook DM. Racial discrimination in health care and utilization of health care: a crosssectional study of California adults. J Gen Intern Med 2018;33:1760–7.
- Nong P, Raj M, Creary M, Kardia SLR, Platt JE. Patient-reported experiences of discrimination in the US health care system. JAMA Netw Open 2020;3:e2029650.
- Platt J, Nong P. An ecosystem approach to earning and sustaining trust in health care—too big to care. JAMA Health Forum 2023;4:e224882.
- Ricks TN, Abbyad C, Polinard E. Undoing racism and mitigating bias among healthcare professionals: lessons learned during a systematic review. J Racial Ethn Health Disparities 2022;9:1990–2000.
- 74. Lie DA, Lee-Rey E, Gomez A, Bereknyei S, Braddock CH. Does cultural competency training of health professionals improve patient outcomes? A systematic review and proposed algorithm for future research. J Gen Intern Med 2011;26:317–25.
- 75. Renzaho AMN, Romios P, Crock C, Sønderlund AL. The effectiveness of cultural competence programs in ethnic minority patient-centered health care–a systematic review of the literature. Int J Qual Health Care J Int 2013;25:261–9.
- Cooper LA, Roter DL, Johnson RL, Ford DE, Steinwachs DM, Powe NR. Patient-centered communication, ratings of care, and concordance of patient and physician race. Ann Intern Med 2003; 139:907–15.

- 77. Shen MJ, Peterson EB, Costas-Muñiz R, et al. The effects of race and racial concordance on patient-physician communication: a systematic review of the literature. J Racial Ethn Health Disparities 2018;5:117–40.
- Blanchard J, Nayar S, Lurie N. Patient-provider and patient-staff racial concordance and perceptions of mistreatment in the health care setting. J Gen Intern Med 2007;22:1184–9.
- Ma A, Sanchez A, Ma M. The impact of patientprovider race/ethnicity concordance on provider visits: updated evidence from the medical expenditure panel survey. J Racial Ethn Health Disparities 2019;6:1011–20.
- Jetty A, Jabbarpour Y, Pollack J, Huerto R, Woo S, Petterson S. Patient-physician racial concordance associated with improved healthcare use and lower healthcare expenditures in minority populations. J Racial Ethn Health Disparities 2022;9:68–81.
- Thompson T Ann M, Young YY, Bass TM, et al. Racism runs through it: examining the sexual and reproductive health experience of black women in the South. Health Aff (Millwood) 2022;41:195–202.
- 82. Hunte R, Klawetter S, Paul S. "Black nurses in the home is working": advocacy, naming, and processing racism to improve Black maternal and infant health. Matern Child Health J 2022;26:933–40.
- Hassen N, Lofters A, Michael S, Mall A, Pinto AD, Rackal J. Implementing anti-racism interventions in healthcare settings: a scoping review. Int J Environ Res Public Health 2021;18:2993.
- Bryant A, Walsh-Felz A, Jacklitz J, Lindberg S. The impact of a community resource navigator program on patient trust. WMJ Off Publ State Med Soc Wis 2020;119:190–3.
- Hickey E, Phan M, Beck AF, Burkhardt MC, Klein MD. A mixed-methods evaluation of a novel food pantry in a pediatric primary care center. Clin Pediatr (Phila) 2020;59:278–84.
- Decker MR, Flessa S, Pillai RV, et al. Implementing trauma-informed partner violence assessment in family planning clinics. J Womens Health 2002 2017;26: 957–65.
- Davis J, Zinck L, Kelly S, et al. Screened social risk factors and screening acceptability among oncology patients in Philadelphia. JCO 2021;39: 12125–12125.
- Byhoff E, Gottlieb LM. When there is value in asking: an argument for social risk screening in clinical practice. Ann Intern Med 2022;175: 1181–2.
- 89. Garg A, Boynton-Jarrett R, Dworkin PH. Avoiding the unintended consequences of screening for social determinants of health. JAMA 2016;316:813–4.
- 90. Wrangle J, Fisher JW, Paranjape A. Ha sentido sola? Culturally competent screening for intimate

partner violence in Latina women. J Womens Health 2008;17:261–8.

91. Mainous AG, Baker R, Love MM, Gray DP, Gill JM. Continuity of care and trust in one's physician: evidence from primary care in the United States and the United Kingdom. Fam Med 2001; 33:22–7.

92. Pandhi N, Saultz JW. Patients' perceptions of interpersonal continuity of care. J Am Board Fam Med 2006;19:390–7.

Participant Recruitment and Survey Completion



Appendix 2

The SIREN Accountable Health Communities screening tool study

RA marks one of the following:

____ Participant is receiving care today

Participant is the adult caregiver of a child receiving care today

We appreciate your participation in this survey. Your responses will help us understand more about how to develop programs that can help respond to the needs of our patients.

AHC Screening instrument

Housing Stability and Conditions

1. What is your housing situation today?

- a) I have a steady place to live
- b) I have a place to live today, but I am worried about losing it in the future
- c) I do not have a steady place to live (I am temporarily staying with others, in a hotel, in a shelter, living outside on the street, on a beach, in a car, abandoned building, bus or train station, or in a park)
- d) I prefer not to answer

2. Think about the place you live. Do you have problems with any of the following? (Check all that apply)

- a) Pests such as bugs, ants, or mice
- b) Mold
- c) Lead paint or pipes
- d) Lack of heat
- e) Oven or stove not working
- f) Smoke detectors missing or not working
- g) Water leaks
- h) None of the above
- i) I prefer not to answer

Food Security

Some people have made the following statements about their food situation. Please answer whether the statements were OFTEN, SOMETIMES, or NEVER true for you and your household in the last 12 months.

- 3. Within the past 12 months, you worried that your food would run out before you got money to buy more.
 - a) Often true
 - b) Sometimes true
 - c) Never true
 - d) I prefer not to answer
- Within the past 12 months, the food you bought just didn't last and you didn't have money to get more.
 - a) Often true
 - b) Sometimes true
 - c) Never true
 - d) I prefer not to answer

Transportation Needs

- 5. In the past 12 months, has lack of reliable transportation kept you from medical appointments, meetings, work or from getting things needed for daily living?
 - a) Yes
 - b) No
 - c) I prefer not to answer

Utility Needs

6. In the past 12 months has the electric, gas, oil, or water company threatened to shut off services in your home?a) Yes

u) 10

b) No

c) Already shut off

d) I prefer not to answer

Interpersonal Safety

Because violence and abuse happens to a lot of people and affects their health we are asking the following questions.

7. How often does anyone, including family and friends, physically hurt you?

a) Never (1)

b) Rarely (2)

c) Sometimes (3)

d) Fairly often (4)

e) Frequently (5)

f) I prefer not to answer

8. How often does anyone, including family and friends, insult or talk down to you?

a) Never (1)

b) Rarely (2)

c) Sometimes (3)

d) Fairly often (4)

e) Frequently (5)

f) I prefer not to answer

9. How often does anyone, including family and friends, threaten you with harm?

a) Never (1)

b) Rarely (2)

c) Sometimes (3)

d) Fairly often (4)

e) Frequently (5)

f) I prefer not to answer

10. How often does anyone, including family and friends, scream or curse at you?

a) Never (1)

b) Rarely (2)

c) Sometimes (3)

d) Fairly often (4)

e) Frequently (5)

f) I prefer not to answer

A score of 11 or more when the numerical values for answers to questions 7-10 are added shows that the person might not be safe.

Additional housing questions

11. In the past 12 months, was there a time when you were not able to pay the mortgage or rent on time?

a) Yes

b) No

12. In the past 12 months, how many times have you moved where you were living? (drop down)

(Answer is # of moves. Positive screen if answer is 2 or more moves in the last 12 months)

13. At any time in the past 12 months, were you homeless or living in shelter (including now)?

a) Yes

b) No

Desire for help with needs

14. Would you like to receive assistance with any of the issues below: (check all that apply)

- a) Housing
- b) Food access
- c) Medical or non-medical transportation
- d) Electric, gas, oil, or water utility services
- e) Your safety, or violence in your household
- f) None of these

Patient Acceptability of Social Screening

15. In the last 12 months, not including today, have you been asked about any of the following in any health care setting: (check all that apply)

- a) Yes, housing
- b) Yes, food access
- c) Yes, medical or non-medical transportation
- d) Yes, electric, gas, oil, or water utility services
- e) Yes, your safety, or violence in your household
- f) No, none of these

16. In the last 12 months, not including today, have you received assistance from anyone in any health care setting related to: (check all that apply)

- a) Housing
- b) Food access
- c) Medical or non-medical transportation
- d) Electric, gas, oil, or water utility services
- e) Your safety, or violence in your household
- f) None of these

17a. Branch if primary care: How long have you or your family been receiving care at this clinic?

- a) Less than 1 year
- b) 1 to less than 3 years
- c) 3 to less than 5 years
- d) 5 years of more

17b. *Branch if adult ED patient*: Is there a doctor or place that you usually go if you are sick or need advice about your health?

- a) Yes
- b) No

17c. Branch if adult caregiver of pediatric ED patient: Is there a doctor or place that you usually go if your child is sick or you need advice about your child's health?

- c) Yes
 - d) No

18. Do you think it is appropriate to be asked these questions about your social and economic needs at [BRANCH: "this clinic" OR "this emergency department"]?

- a) Very appropriate
- b) Somewhat appropriate
- c) Neither appropriate nor inappropriate
- d) Somewhat inappropriate
- e) Very inappropriate

19. Please check if you felt uncomfortable today being asked any of the questions about:

(check all that apply)

- a) Housing
- b) Food access
- c) Medical or non-medical transportation
- d) Electric, gas, oil, or water utility services
- e) Your safety, or violence in your household
- f) None of these

20. In general, where do you think people should be asked questions about their social and economic needs? Check all that apply.

- a) Emergency Department
- Primary Care/Pediatrician's Office b)
- Other health care settings (e.g. specialty clinics, hospital) c)
- Non-health care settings (e.g. school, community center) d)
- e) None of these places

21. How frequently do you think it is appropriate to be asked these questions about social and economic needs?

(Branch based on location)

- i. In a primary care outpatient clinic visit:
 - a. every time I receive care
 - b. once every 6 months
 - c. once a year
 - d. every 2 years e. every 5 years
 - f. never

ii. In the emergency room/urgent care:

- a. every time I receive care
- b. once every 6 months
- c. once a year d.
- every 2 years e. every 5 years
- f. never

22. Would you be comfortable having these kinds of needs included in your health records (also known as your medical record or chart)?

- a) Completely comfortable
- b) Somewhat comfortable
- Neither comfortable nor uncomfortable c)
- Somewhat uncomfortable d)
- e) Completely uncomfortable

Self-rated health

23a. In general, would you say your health is...(select one)

- a) Excellent
- b) Very good
- c) Good
- d) Fair
- Poor e)
- f) Don't know/refused to answer

BRANCH: If patient is a child, survey will branch to include parent-reported child health question below

23b. In general, would you say your child's health is... (select one)

- a) Excellent
- b) Very good
- c) Good
- d) Fair
- e) Poor
- f) Don't know/refused to answer

Patient trust in their provider

24. *BRANCH if patient in clinic*: How much do you trust your health care provider(s) at this clinic?

BRANCH if caregiver of pediatric patient in clinic: How much do you trust your child's health care provider(s) at this clinic?

BRANCH if ED: How much do you trust the health care providers at this emergency department?

1	2	3	4	5	6	7	8	9	10
Not at all									Completely

Demographics

25. What is your age? a) 18 to 24 b) 25 to 34 c) 35 to 44

d) 45 to 54
e) 55 to 64
f) 65 to 74

g) 75 or older

26. What is the highest grade (or year) of school you completed? (select one)

Elementary School	High School	College	Graduate School
01	09	13	17
02	10	14	18
03	11	15	19
04	12	16	20+
05			
06			
07			

08

27. How do you describe your gender?

- a) Male
- b) Female
- c) Trans male/Trans mand) Trans female/Trans woman
- e) Gender-queer/Gender non-conforming
- f) Different identity
- g) Prefer not to answer

28. What is your race? (mark all that apply)

- a) White
- b) Black or African American
- c) American Indian or Alaska Native
- d) Asian Indian
- e) Chinese
- f) Filipino
- g) Japanese
- h) Korean
- i) Vietnamese
- j) Native Hawaiian
- k) Guamanian or Chamorro
- l) Samoan
- m) Other Pacific Islander (specify)
- n) Other Asian (specify)_____
- o) Some other race (specify)

29. Are you of Hispanic or Latino origin?

- a) No, not Hispanic, Latino or Spanish origin
- b) Yes, Mexican, Mexican American, Chicano
- c) Yes, Puerto Ricand) Yes, Cuban
- e) Yes, another Hispanic, Latino, or Spanish origin ______

30a. Which of the following categories best describes your total combined household income for the past 12 months?

- a) 0 \$5,000
- b) \$5,001 \$10,000
 c) \$10,001 \$15,000
- \$10,001 \$15,000
 \$15,001 \$20,000
- d) \$15,001 \$20,000e) \$20,001 \$25,000
- e) \$20,001 \$25,000
 f) \$25,001 \$30,000
- g) \$30,001 \$35,000
- h) \$35,001 \$40,000
- i) \$40,001 \$50,000
- j) \$50,001 \$75,000
- k) \$75,001 \$100,000
- 1) \$100,001 \$150,000
- m) \$150,000 +
- n) Don't know
- o) Would rather not say

30b. How many people (kids and adults) are currently dependent on this income and living in your household, including yourself? (*drop down*)

(Answer is # of people in household dependent on income)

30c. Of these people, how many are 0-17 years old? (drop down)

(Answer is # of people in household 0-17 years old)

31. When getting health care, have you ever had any of the following things happen to you because of your race, ethnicity, or socioeconomic status?

- a) Felt like a doctor or nurse was not listening to what you were saying Drop down Yes/No
- b) Treated you with less respect than other people Drop down Yes/No
- c) Received poorer services than other people Drop down Yes/No
- d) Treated with less courtesy than other people
- Drop down Yes/No e) Had a doctor or nurse act as if he or she was better than you Drop down Yes/No
- f) Had a doctor or nurse act as if he or she thinks you were not smart Drop down Yes/No
- g) Had a doctor or nurse act as if he or she was afraid of you Drop down Yes/No

32. Finally, we would like to understand where you see yourself in relation to others in the United States.

Think of the ladder below as representing where people stand in the United States.

At the top of the ladder (10) are the people who are the best off—those who have the most money, the most education, and the most respected jobs.

At the bottom (1) are the people who are the worst off—who have the least money, least education, and the least respected jobs or no job. The higher up you are on this ladder, the closer you are to the people at the very top.

Where would you put yourself on this ladder?

Please respond with the number (1-10) where you think you stand at this time in your life, relative to other people in the United States.

Answer 1-10



Appendix 3

	Adjusted Odds Ratio (95%
Healthcare discrimination score	0.74 (0.60, 0.93)
Race and ethnicity	
Non-Hispanic White (reference)	
Non-Hispanic Black	0.80 (0.50, 1.23)
Hispanic	0.71 (0.44, 1.07)
Non-Hispanic multiple/other	0.61 (0.30, 1.13)
Age	
18 to 44 (reference)	
45 to 64	1.34 (0.84, 2.17)
65+	1.27 (0.68, 2.42)
Education 12 or more years	0.40 (0.27, 0.63)
Income	
\$0–50k (reference)	
\$50,001–75k	1.08 (0.55, 2.28)
\$75,001+	0.81 (0.47, 1.49)
SES ladder	1.06 (1.01, 1.11)
Self-rated health good/very good/excellent	1.73 (1.16, 2.55)
Interaction: Self-rated health * discrimination score	0.95 (0.80, 1.12)
80% or more publicly insured/uninsured	0.86 (0.57, 1.27)
Previously screened for social risks in prior 12 months	0.93 (0.57, 1.52)
Interaction: Previously screened * discrimination score	1.33 (1.02, 1.74)
Previously assisted with social risks in prior 12 months	0.91 (0.64, 1.29)

Table 1. Adjusted Associations Between Complete Trust in Healthcare Provider and Variables of Interest (n = 690) (Sensitivity Analysis Using 1–9 vs 10 Cut Point)

Abbreviations: SES, socioeconomic status; CI, confidence interval.

CI)

Table 2. Adjusted Associations Between Complete Trust in Healthcare Provider and Variables of Interest,Stratified by Race and Ethnicity (Sensitivity Analysis Using 1–9 vs 10 Cut Point)

		Adjusted Odds	s Ratio (95% CI)	
	Non-Hispanic White (n = 275)	Non-Hispanic Black (n = 161)	Hispanic (n = 196)	Non-Hispanic Multiple/Other (n = 55)
Healthcare discrimination score	0.69 (0.53, 0.88)	0.70 (0.49, 1.01)	0.73 (0.57, 0.93)	0.49 (0.26, 0.93)
Age				
18 to 44 (reference)				
45 to 64	1.50 (0.71, 3.18)	1.77 (0.88, 3.58)	0.55 (0.21, 1.44)	6.95 (1.72, 28.06)
65+	1.43 (0.64, 3.18)	1.60 (0.49, 5.30)	1.00 (0.18, 5.57)	_
Education 12 or more years	0.35 (0.09, 1.41)	0.46 (0.21, 1.02)	0.35 (0.18, 0.68)	0.12 (0.02, 0.82)
Income				
\$0–50k (reference)				
\$50,001–75k	1.92 (0.58, 6.30)	0.75 (0.29, 1.94)	0.60 (0.18, 2.06)	0.36 (0.03, 4.21)
\$75,001+	0.95 (0.41, 2.23)	0.67 (0.15, 3.03)	0.30 (0.05, 1.93)	0.30 (0.05, 1.86)
SES ladder	1.05 (0.91, 1.22)	1.06 (0.96, 1.17)	1.04 (0.94, 1.14)	1.55 (0.89, 2.70)
Self-rated health good/very good/excellent	1.65 (1.02, 2.67)	1.04 (0.32, 3.39)	2.67 (1.44, 4.93)	1.03 (0.30, 3.49)
Interaction: Self-rated health * discrimination score	1.01 (0.84, 1.21)	1.17 (0.74, 1.85)	0.57 (0.40, 0.81)	0.87 (0.35, 2.12)
80% or more publicly insured/uninsured	0.91 (0.48, 1.73)	0.86 (0.61, 1.20)	0.86 (0.57, 1.57)	0.61 (0.17, 2.20)
Previously screened for social risks in prior 12 months	0.98 (0.63, 1.53)	0.92 (0.42, 2.03)	0.60 (0.89, 1.95)	3.21 (0.51, 20.41)
Interaction: Previously screened * discrimination score	1.65 (1.27, 2.13)	1.09 (0.70, 1.71)	1.92 (1.17, 3.15)	1.49 (0.75, 2.98)
Previously assisted with social risks in prior 12 months	0.96 (0.32, 2.88)	0.54 (0.36, 0.80)	1.91 (0.63, 1.37)	0.32 (0.05, 2.07)

	Adjusted Odds Ratio (95% CI)
Healthcare discrimination score	0.82 (0.69, 0.98)
Race and ethnicity	
Non-Hispanic White (reference)	
Non-Hispanic Black	0.80 (0.48, 1.34)
Hispanic	0.81 (0.49, 1.33)
Non-Hispanic multiple/other	1.02 (0.49, 2.09)
Age	
18 to 44 (reference)	
45 to 64	1.33 (0.81, 2.18)
65+	1.52 (0.62, 3.72)
Education 12 or more years	0.51 (0.37, 0.71)
Income	
\$0–50k (reference)	
\$50,001–75k	1.36 (0.81, 2.28)
\$75,001+	1.11 (0.55, 2.24)
SES ladder	1.02 (0.95, 1.10)
Self-rated health good/very good/excellent	1.98 (1.54, 2.55)
Interaction: Self-rated health * discrimination score	0.86 (0.74, 1.00)
80% or more publicly insured/uninsured	0.81 (0.50, 1.32)
Previously screened for social risks in prior 12 months	1.00 (0.57, 1.76)
Interaction: Previously screened * discrimination score	1.20 (0.93, 1.55)
Previously assisted with social risks in prior 12 months	1.09 (0.77, 1.55)

Table 3. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest (n = 690) (Sensitivity Analysis Using 1–8 vs 9–10 Cut Point)

Table 4.	Adjusted	Association	s Between Hi	gh Trust in	Healthcare	Provider and	l Variables	of Interest,	Stratified by
Race and	l Ethnicity	(Sensitivity	Analysis Usi	ng 1–8 vs 9	–10 Cut Poi	int)			

	Adjusted Odds Ratio (95% CI)			
	Non-Hispanic White $(n = 275)$	Non-Hispanic Black (n = 161)	Hispanic (n = 196)	Non-Hispanic Multiple/Other (n = 58)
Healthcare discrimination score	0.82 (0.62, 1.08)	0.71 (0.53, 0.96)	0.66 (0.48, 0.89)	0.73 (0.48, 1.12)
Age				
18 to 44 (reference)				
45 to 64	1.53 (0.78, 3.02)	1.28 (0.52, 3.14)	0.49 (0.13, 1.78)	7.43 (1.42, 38.94)
65+	1.62 (0.69, 3.82)	1.28 (0.23, 7.20)	1.93 (0.41, 9.01)	2.19 (0.14, 34.67)
Education 12 or more years	0.22 (0.03, 1.73)	0.94 (0.43, 2.06)	0.30 (0.18, 0.48)	0.23 (0.05, 1.02)
Income				
\$0–50k (reference)				
\$50,001–75k	1.45 (0.46, 4.57)	1.25 (0.32, 4.85)	1.81 (0.71, 4.64)	0.91 (0.05, 15.84)
\$75,001+	1.39 (1.52, 4.62)	0.93 (0.17, 4.94)	0.33 (0.02, 5.21)	0.11 (0.01, 0.89)
SES ladder	1.00 (0.89, 1.14)	1.03 (0.88, 1.21)	1.03 (0.94, 1.13)	1.08 (0.55, 2.10)
Self-rated health good/very good/excellent	2.65 (1.52, 4.62)	0.77 (0.50, 1.20)	2.72 (1.77, 4.19)	4.38 (0.98, 19.55)
Interaction: Self-rated health * discrimination score	0.88 (0.69, 1.13)	1.12 (0.88, 1.43)	0.52 (0.24, 1.12)	0.80 (0.47, 1.35)
80% or more publicly insured/uninsured	0.65 (0.37, 1.13)	0.83 (0.56, 1.23)	1.04 (0.44, 2.46)	0.23 (0.08, 0.65)
Previously screened for social risks in prior 12 months	0.98 (0.54, 1.78)	1.16 (0.43, 3.10)	0.88 (0.31, 2.51)	0.36 (0.03, 4.59)
Interaction: Previously screened * discrimination score	1.45 (1.02, 2.05)	1.04 (0.73, 1.48)	1.96 (0.88, 4.35)	1.25 (0.61, 2.54)
Previously assisted with social risks in prior 12 months	0.95 (0.34, 2.63)	0.84 (0.34, 2.09)	2.02 (0.43, 9.56)	0.64 (0.08, 5.33)

	Adjusted Odds Ratio (95% CI)
Healthcare discrimination score	0.73 (0.63, 0.85)
Race and ethnicity	
Non-Hispanic White (reference)	
Non-Hispanic Black	0.72 (0.45, 1.16)
Hispanic	0.67 (0.43, 1.06)
Non-Hispanic multiple/other	0.73 (0.40, 1.33)
Age	
18 to 44 (reference)	
45 to 64	1.30 (0.87, 1.92)
65+	1.16 (0.61, 2.21)
Education 12 or more years	0.41 (0.28, 0.61)
Income	
\$0–50k (reference)	
\$50,001–75k	1.29 (0.72, 2.30)
\$75,001+	0.93 (0.55, 1.56)
SES ladder	1.05 (0.98, 1.11)
Self-rated health good/very good/excellent	2.05 (1.40, 2.99)
Interaction: Self-rated health * discrimination score	0.92 (0.81, 1.05)
80% or more publicly insured/uninsured	0.83 (0.62, 1.10)
Previously screened for social risks in prior 12 months	0.93 (0.61, 1.42)
Interaction: Previously screened * discrimination score	1.33 (1.01, 1.74)
Previously assisted with social risks in prior 12 months	0.88 (0.63, 1.24)

Table 5. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest (n = 690) (Sensitivity Analysis Using 1–7 vs 8–9 vs 10 Cut Point)

	Adjusted Odds Ratio (95% CI)				
	Non-Hispanic White $(n = 275)$	Non-Hispanic Black (n = 161)	Hispanic (n = 196)	Non-Hispanic Multiple/Other (n = 58)	
Healthcare discrimination score	0.73 (0.59, 0.90)	0.68 (0.50, 0.92)	0.58 (0.45, 0.76)	0.71 (0.55, 0.91)	
Age					
18 to 44 (reference)					
45 to 64	1.38 (0.68, 2.77)	1.58 (0.80, 3.15)	0.62 (0.24, 1.63)	3.93 (1.03, 14.99)	
65+	1.35 (0.67, 2.74)	1.42 (0.31, 6.54)	0.90 (0.20, 3.94)	0.35 (0.09, 1.34)	
Education 12 or more years	0.37 (0.08, 1.73)	0.69 (0.30, 1.60)	0.30 (0.16, 0.56)	0.26 (0.02, 2.78)	
Income					
\$0–50k (reference)					
\$50,001–75k	2.01 (0.70, 5.81)	1.06 (0.52, 2.20)	0.86 (0.39, 1.91)	0.93 (0.17, 5.04)	
\$75,001+	1.01 (0.46, 2.20)	0.95 (0.27, 3.43)	0.42 (0.10, 1.67)	0.74 (0.18, 2.96)	
SES ladder	1.05 (0.92, 1.20)	1.04 (0.92, 1.18)	1.02 (0.92, 1.12)	1.20 (0.76, 1.88)	
Self-rated health good/very good/excellent	2.27 (1.34, 3.84)	1.16 (0.48, 2.80)	3.08 (1.55, 6.10)	1.46 (0.27, 7.98)	
Interaction: Self-rated health * discrimination score	1.01 (0.84, 1.21)	1.10 (0.78, 1.54)	0.53 (0.32, 0.86)	0.87 (0.46, 1.66)	
80% or more publicly insured/uninsured	0.95 (0.46, 1.97)	1.01 (0.67, 1.52)	0.65 (0.33, 1.28)	0.63 (0.19, 2.10)	
Previously screened for social risks in prior 12 months	1.03 (0.62, 1.69)	0.97 (0.53, 1.77)	0.57 (0.22, 1.53)	1.85 (0.36, 9.53)	
Interaction: Previously screened * discrimination score	1.56 (1.15, 2.10)	1.09 (0.72, 1.65)	2.61 (1.47, 4.62)	1.05 (0.60, 1.85)	

1.08 (0.37, 3.17)

0.61 (0.36, 1.04)

1.73 (0.56, 5.33)

0.37 (0.06, 2.30)

Table 6. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest, Stratified by Race and Ethnicity (Sensitivity Analysis Using 1–7 vs 8–9 vs 10 Cut Point)

Abbreviations: SES, socioeconomic status; CI, confidence interval.

Previously assisted with social risks in prior

12 months

	Adjusted Odds Ratio (95% CI)		
	Adult Patients ($n = 516$)	Pediatric Caregivers (n = 163)	
Healthcare discrimination score	0.69 (0.57, 0.83)	1.16 (0.80, 1.67)	
Race and ethnicity			
Non-Hispanic White (reference)			
Non-Hispanic Black	0.50 (0.22, 1.15)	0.40 (0.32, 0.50)	
Hispanic	0.81 (0.33, 1.99)	0.22 (0.53, 0.92)	
Non-Hispanic multiple/other	1.38 (0.47, 4.03)	0.06 (0.01, 0.32)	
Age			
18 to 44 (reference)			
45 to 64	1.31 (0.86, 2.01)	1.00 (-)	
65+	1.21 (0.46, 3.21)		
Education 12 or more years	0.68 (0.34, 1.36)	0.24 (0.13, 0.43)	
Income			
\$0–50k (reference)			
\$50,001–75k	2.44 (0.83, 7.17)	1.02 (0.11, 9.38)	
\$75,001+	1.42 (0.81, 2.50)	2.70 (0.85, 8.54)	
SES ladder	0.98 (0.87, 1.11)	1.32 (1.25, 1.40)	
Self-rated health good/very good/excellent	3.32 (2.13, 5.16)	3.18 (0.75, 13.44)	
Interaction: Self-rated health * discrimination score	0.83 (0.67, 1.04)	0.69 (0.50, 0.95)	
80% or more publicly insured/uninsured	0.75 (0.41, 1.36)	1.16 (0.52, 2.58)	
Previously screened for social risks in prior 12 months	0.98 (0.52, 1.84)	1.04 (0.51, 2.09)	
Interaction: Previously screened * discrimination score	1.40 (1.05, 1.88)	0.96 (0.79, 1.17)	
Previously assisted with social risks in prior 12 months	0.91 (0.57, 1.45)	0.89 (0.39, 1.99)	

Table 7. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest, Stratified by Participant Type (Original 1–7 vs 8–10 Cut Point)

	Adjusted Odds Ratio (95% CI)		
	Primary Care ($n = 410$)	Emergency Department (n = 280)	
Healthcare discrimination score	0.64 (0.46, 0.89)	0.82 (0.67, 0.99)	
Race and ethnicity			
Non-Hispanic White (reference)			
Non-Hispanic Black	0.48 (0.9, 1.27)	0.73 (0.19, 2.77)	
Hispanic	0.57 (0.23, 1.42)	0.69 (0.10, 4.86)	
Non-Hispanic multiple/other	1.31 (0.35, 4.98)	0.64 (0.09, 4.29)	
Age			
18 to 44 (reference)			
45 to 64	0.89 (0.62, 1.27)	3.67 (1.69, 7.98)	
65+	0.83 (0.22, 3.17)	1.91 (0.54, 6.86)	
Education 12 or more years	0.38 (0.29, 0.50)	0.93 (0.24, 3.55)	
Income			
\$0–50k (reference)			
\$50,001–75k	2.94 (0.62, 13.92)	1.45 (0.37, 5.61)	
\$75,001+	1.80 (0.94, 3.43)	1.06 (0.57, 1.97)	
SES ladder	0.94 (0.82, 1.08)	1.17 (1.05, 1.30)	
Self-rated health good/very good/excellent	3.21 (2.04, 5.06)	3.36 (1.19, 9.51)	
Interaction: Self-rated health * discrimination score	0.80 (0.59, 1.08)	0.96 (0.77, 1.21)	
80% or more publicly insured/uninsured	0.83 (0.50, 1.38)	0.75 (0.34, 1.65)	
Previously screened for social risks in prior 12 months	1.18 (0.56, 2.50)	0.71 (0.64, 0.79)	
Interaction: Previously screened * discrimination score	1.59 (1.12, 2.26)	1.03 (0.87, 1.21)	
Previously assisted with social risks in prior 12 months	0.95 (0.55, 1.66)	0.92 (0.64, 1.34)	

Table 8. Adjusted Associations Between High Trust in Healthcare Provider and Variables of Interest, Stratified by Healthcare Setting (Original 1–7 vs 8–10 Cut Point)

Table 9. Characteristics of Disaggregated Hispanic Participants

	n (%)				
	Total	Mexican, Mexican American, Chicano (n = 137)	Puerto Rican (n = 42)	Other Hispanic Latino, or Spanish Origin (n = 150)	
Biological sex					
Female	242 (74)	103 (75)	30 (71)	109 (73)	
Age					
18 to 44	220 (68)	90 (67)	28 (67)	102 (69)	
45 to 64	80 (25)	38 (28)	11 (26)	31 (21)	
65+	26 (8)	7 (5)	3 (7)	16 (11)	
Years of education					
12 or more years	204 (62)	82 (60)	34 (81)	87 (59)	
Income					
\$0-50k	228 (89)	109 (96)	27 (75)	92 (86)	
\$50,001–75k	15 (6)	1 (1)	7 (19)	7 (7)	
\$75,001+	14 (5)	4 (4)	2 (6)	8 (8)	
SES ladder position*	5.15 (2.28)	5.30 (2.12)	5.71 (2.05)	4.88 (2.45)	
Healthcare setting	. ,	× ,	. ,		
Primary care	209 (64)	111 (81)	10 (24)	88 (59)	
Clinic type					
Urban academic	196 (60)	53 (39)	40 (95)	103 (69)	
Urban non-academic	58 (18)	37 (27)	1 (2)	20 (13)	
Non-urban non-academic	75 (23)	47 (34)	1 (2)	27 (18)	
Percentage publicly insured/uninsured at site					
80% or more publicly insured/uninsured	159 (48)	68 (50)	13 (31)	78 (52)	
Participant type					
Adult patient	275 (84)	124 (91)	29 (69)	122 (81)	
Preferred language					
Spanish	169 (51)	76 (56)	7 (17)	86 (57)	
Patient/caregiver-reported health					
Good, Very good, excellent	231 (74)	90 (68)	27 (66)	114 (81)	
Experienced any of 7 categories perceived discrimination	75 (23)	27 (20)	13 (31)	35 (23)	
Healthcare discrimination score*	0.55 (1.31)	0.54 (1.36)	1 (1.80)	0.43 (1.05)	
Asked about needs in any domains in the past 12 months	81 (26)	39 (30)	10 (24)	32 (22)	
Assisted with needs in any domains in the past 12 months	58 (18)	20 (15)	7 (17)	31 (22)	

*mean (S.D.).

Abbreviation: SES, socioeconomic status.

J Am Board Fam Med: first published as 10.3122/jabfm.2023.230182R1 on 25 October 2024. Downloaded from http://www.jabfm.org/ on 6 May 2025 by guest. Protected by copyright.

Table 10. Trust in Healthcare Provider and Experiences of Healthcare Discrimination Among Disaggregated Hispanic Participants

	n (%)			
	Total	Mexican, Mexican American, Chicano (n = 137)	Puerto Rican (n = 42)	Other Hispanic, Latino, or Spanish Origin (n = 150)
Low/medium trust (1 to 7)	70 (23)	27 (21)	13 (32)	30 (21)
High trust (8 to 10)	240 (77)	104 (80)	28 (68)	108 (78)
Experienced any of 7 categories perceived discrimination	75 (23)	27 (20)	13 (31)	35 (23)
Item 1: Felt like a doctor or nurse was not listening to what you were saying	59 (18)	23 (17)	11 (26)	25 (17)
Item 2: Treated you with less respect than other people	28 (9)	12 (9)	7 (17)	9 (6)
Item 3: Received poorer services than other people	20 (6)	8 (6)	6 (14)	6 (4)
Item 4: Treated with less courtesy than other people	29 (9)	12 (9)	7 (17)	10 (7)
Item 5: Had a doctor or nurse act as if he or she was better than you	21 (7)	8 (6)	5 (12)	8 (5)
Item 6: Had a doctor or nurse act as if he or she thinks you were not smart	24 (7)	9 (7)	5 (12)	10 (7)
Item 7: Had a doctor or nurse act as if he or she was afraid of you	3 (1)	2 (2)	0 (0)	1 (1)