

## BRIEF REPORT

## Association of Social Needs with Diabetes Outcomes in an Older Population

David M. Mosen, PhD, MPH, Stephanie L. Fitzpatrick, PhD, Erin M. Keast, MPH, John F. Dickerson, PhD, Briar L. Ertz-Berger, MD, MPH, and Matthew P. Banegas, PhD, MPH, MS

**Background:** Identifying and addressing social needs is important to improve health for older adults with type 2 diabetes mellitus (T2DM). However, few studies have examined the association between social needs and T2DM-related outcomes among older populations within integrated health care systems. This study examined the association between social needs and DM-related outcomes among older adults with T2DM receiving care at Kaiser Permanente Northwest.

**Methods:** From a cohort of 1954 Medicare members ages 65 and older who completed a social needs questionnaire, we examined the association between 5 dichotomous (yes vs no) social needs measures (financial strain, food insecurity, housing instability, social isolation, and transportation needs) and 3 DM-specific outcomes in the 12 months after assessment: 1) good glycemic control (HbA1c < 8%); 2) any DM-specific emergency department (ED) utilization; and 3) any DM-specific hospital admissions.

**Results:** About 40% of the study population reported 1 or more social needs. Financial strain (OR = 0.56, 95% CI = 0.36-0.85), food insecurity (OR = 0.47, 95% CI = 0.28-0.81) and housing instability (OR = 0.50, 95% CI = 0.25-0.99) were associated with lower odds of good glycemic control. All 5 social needs were associated with higher ED utilization (financial strain: OR = 1.65, 95% CI = 1.17-2.33; food insecurity: OR = 1.62, 95% CI = 1.02-2.57; housing instability: OR = 2.14, 95% CI = 1.23-3.75; social isolation: OR = 1.36, 95% CI = 1.06-1.74; transportation needs: OR = 1.83, 95% CI = 1.23-2.71). Financial strain was also associated with higher hospital admissions (OR = 1.77, 95% CI = 1.17-2.68).

**Discussion:** Associations between social needs and DM-specific outcomes demonstrate the need to develop programs to address social needs in the clinical setting and test whether such interventions improve DM-related outcomes. (J Am Board Fam Med 2025;38:125-132.)

**Keywords:** Financial Stress, Food Insecurity, Housing Instability, Integrated Health Care Systems, Medicare, Social Determinants of Health, Social Isolation, Surveys and Questionnaires, Type 2 Diabetes Mellitus

## Background

Examining social needs among older adults with type 2 diabetes mellitus (T2DM) in the United

States is an important research area, as nearly 30% of adults ages 65 and older have T2DM,<sup>1</sup> resulting in over \$400 billion in health care costs annually<sup>2</sup> and older adults experience high levels of social

This article was externally peer reviewed.

Submitted 3 April 2024; revised 10 April 2024; accepted 16 September 2024.

From the Kaiser Permanente Center for Health Research, Portland, OR (DMM, SLF, EMK, JFD, MPB); Radiation Medicine and Applied Sciences, University of California San Diego, La Jolla, CA (MPB); Northwell, New Hyde Park, NY, Institute of Health System Science, Feinstein Institutes of Medical Research, Manhasset, NY and Department of Medicine, Manhasset, NY (SLF); Northwest Permanente, Continuum of Care Department, Portland, OR (BLE).

**Funding:** This work was supported by the National Institutes of Health (NIH) National Cancer Institute (NCI) Grant No. R01CA253028, (C. Kroenke, PI); Kaiser Permanente's Office of Community Health through the Social Needs Network for Evaluation and Translation (SONNET). SONNET is a national Kaiser Permanente (KP) program that convenes

evaluators and researchers from its eight regional care delivery systems; the KP Bernard J. Tyson School of Medicine; KP's Office of Community Health; the Social Interventions Research and Evaluation Network (SIREN), a KP- and Robert Wood Johnson Foundation-supported network housed at the University of California, San Francisco. SONNET seeks to improve the health of KP members and the communities they live in by developing and implementing new, scientifically driven strategies to shape social health practice and policy; the Northwell, New Hyde Park, NY, Institute of Health System Science, Feinstein Institutes of Medical Research, Manhasset, NY, and Department of Medicine, Manhasset, NY.

**Conflict of interest:** The authors have no conflicts of interest to report.

needs. About 25% report social isolation and loneliness,<sup>3</sup> nearly 1 in 10 experience food insecurity,<sup>4</sup> and 5 to 7% experience financial strain.<sup>5</sup>

Further demonstrating the importance of studying social needs within the T2DM population, previous research found that social needs are associated with negative health and health care utilization outcomes among adults with T2DM. Specifically, food insecurity, financial strain, housing, and transportation needs are associated with suboptimal DM-related outcomes, including lower medication adherence<sup>6</sup> and poor glycemic control.<sup>7–9</sup> Moreover, the presence of social needs may lead to delayed or forgone needed medical care, as well as increased emergency department and inpatient visits.<sup>9–14</sup>

Although some studies have examined the association between social needs and T2DM-related outcomes for older adults,<sup>13–15</sup> few have been conducted in large, integrated health care systems. Studying these associations in large, integrated health care systems among older populations can both: 1) provide insights from systems with comprehensive clinical, demographic, social need, and health care utilization data and 2) help to target quality improvement interventions to those with specific social needs. With this background in mind, the primary objective of this study is to examine the association between social needs and DM-related outcomes among a population of Medicare patients receiving care in a large integrated health care system.

## Methods

### Study Design

This was an observational cohort study of 1954 Medicare Advantage enrollees (ages 65 years or older) who were members of Kaiser Permanente Northwest (KPNW), an integrated health system that provides care for patients who reside in Oregon and southwest Washington.

Patients were part of a larger study<sup>16</sup> that included the following enrollment criteria: 1) completion of a Medicare Total Health Assessment (MTHA) survey during usual care between 8/18/2020 and 1/31/2022 (index date is defined as date of survey completion);

2) completion of all 5 social needs assessments on the MTHA survey: financial strain, food insecurity, housing instability, social isolation, and transportation needs; 3) health plan enrollment for 12 months before and after their index date; 4) no missing covariate data; 5) not on the KPNW research exclusion list; 6) on the KPNW T2DM registry for 12 months before their index date and 12 months after their index date; and 7) completion of an HbA1c test within 12 months before and 12 months after their index date. A process flow of this enrollment criteria is described in Figure 1. The MTHA<sup>16–18</sup> and KPNW T2DM Registry have been described previously.<sup>19</sup> A recent study analyzed similar social needs and all-cause utilization outcomes for the full older adult population using the MTHA survey.<sup>16</sup>

Survey data from the MTHA, as well as clinical and health care utilization outcome measures from the electronic health record (EHR), are housed in the Research Data Warehouse (RDW) at the Kaiser Permanente Center for Health Research (CHR), which has been described previously.<sup>20</sup> This study was approved by the KPNW Institutional Review Board.

## Measures

### Outcome Measures

Three outcome measures were assessed in the 12 months after the index date: 1) good glycemic control, defined as HbA1c < 8%; 2) DM-specific emergency department (ED) visits and 3) DM-specific hospital admissions. Although previous research has used a HbA1c cutoff < 7%,<sup>19</sup> we selected less restrictive of HbA1c < 8%, given guidelines suggest a less rigorous HbA1c cutoff is more appropriate in older adults.<sup>21</sup> If multiple HbA1c tests were conducted in the 12 months after the index date, the first HbA1c test was selected. DM-specific utilization outcomes were based on E10\*\*–E11\*\* ICD-10 codes as encounter diagnoses for ED visits and hospital admissions.


### Independent Variables

The primary independent variables were the presence of 5 social needs (financial strain, food insecurity, housing instability, social isolation, and transportation needs) measured as binary (yes/no) measures, assessed during the index MTHA

---

Corresponding author: David M. Mosen, PhD, 3800 North Interstate Avenue, Portland, OR 97227 (E-mail: david.m.mosen@kpchr.org).

**Figure 1. Population process flow. Abbreviation: MTHA, Medicare Total Health Assessment.**



survey. A description of these social needs is provided in Table 1.

#### **Covariate Measures**

Covariate measures for the analysis were identified from either the MTHA survey or the EHR. Measures identified from the MTHA survey included: highest education level completed, self-reported overall health, and self-reported mental

health. Measures identified from the EHR included: neighborhood deprivation index (NDI)<sup>22</sup>; age, sex, race, ethnicity, Charlson Comorbidity Index (CCI),<sup>23,24</sup> HbA1c status in the 12 months before index date, and DM-specific ED utilization and hospital admissions in the 12 months before index date. If multiple HbA1c tests were conducted in the 12 months before the index date, the first HbA1c test was selected.

**Table 1. Description of Social Needs Measures.**

Individual Social Needs	Responses below Indicate “Yes” Response for Social Need
Financial strain (1 item)	Ability to pay for basics such as food, housing, medical care, and heating rated as “very hard”, “hard” or “somewhat hard.”
Food insecurity (2 items)	Rated the following as “sometimes true” or “often true”: In the past 12 months, 1) worried food would run out before more could be bought or 2) worried food bought would not last and did not have money to get more.
Housing instability (3 items)	In the past 12 months (including now), any of the following three conditions occurred: 1) unable to pay the mortgage or rent on time, 2) lived in 3 or more places or 3) did not have a steady place to sleep or slept in a shelter.
Social isolation (1 item)	Respondent felt lonely or isolated from people around them “sometimes,” “always,” or “often.”
Transportation needs (2 items)	In the past 12 months, lack of transportation: 1) kept respondent from medical appointments or from getting medications or 2) kept respondent from meetings, work, or from getting things needed for daily living.

### Analysis

First, descriptive analyses were conducted for the study population. Next, collinearity diagnostics were used to assess collinearity among each of the social needs variables and the covariate measures. Because no collinearity was found, all covariate measures were included in final logistic regression models. Five logistic regression models were constructed for each dichotomous outcome measure, with each social need variable analyzed as a dichotomous measure (yes vs no [ref. group]).

### Results

**Sample Characteristics (Table 2).** About 60% of the study population were between ages 65 and 74, 52% were male, and nearly 90% were White and non-Hispanic. Nearly 35% reported having a college degree or higher level of education, and on average, patients lived in neighborhoods that had low levels of neighborhood deprivation (mean NDI score =  $-0.2 \pm 0.6$ ). About 30% of the population reported their overall health as “fair” or “poor”; over 60% had a CCI score of 2 or higher. About 20% and 10%, respectively, had any DM-specific ED utilization and DM-specific hospital admissions in the 12 months before the MTHA survey.

**Type and Number of Social Needs (Table 3).** Forty percent of respondents identified at least 1 social need on the MTHA; 3.5% reported 3 or more needs. The most reported social need was social isolation (31.0%), followed by financial strain (9.6%), transportation needs (6.7%), food insecurity (4.9%) and housing instability (3.1%).

### Logistic Regression Results (Table 4)

#### *Association Between Social Needs and HbA1c Control*

Presence of financial strain (OR = 0.56, 95% CI = 0.36–0.85), food insecurity (OR = 0.47, 95% CI = 0.28–0.81), and housing instability (OR = 0.50, 95% CI = 0.25–0.99) were associated with lower odds of having good glycemic control. No other social needs measures were associated with glycemic control as measured by HbA1c values.

#### *Association Between Social Needs and Diabetes-Specific ED Utilization and Hospital Admissions*

The presence of each of the 5 social needs was associated with higher ED utilization in the 12 months following the index date: financial strain (OR = 1.65, 95% CI = 1.17–2.33), food insecurity (OR = 1.62, 95% CI = 1.02–2.57), housing instability (OR = 2.14, 95% CI = 1.23–3.75), social isolation (OR = 1.36, 95% CI = 1.06–1.74) and transportation needs (OR = 1.83, 95% CI = 1.23–2.71). The presence of financial strain was associated with higher hospital admissions in the 12 months following the index date (OR = 1.77, 95% CI = 1.17–2.68).

### Discussion

We found nearly 4 in 10 adults ages 65 and older with T2DM reported at least 1 unmet social need, with over 10% reporting 2 or more needs. Financial strain, food insecurity, and housing instability were associated with poor glycemic control, and all 5 social needs were associated with DM-related ED utilization in the year following assessment. Finally, financial strain was

**Table 2. Population Characteristics**

Population Measures	Total Population (n = 1,954)
Demographic measures	
Age, N (%)	
65 to 74	1,203 (61.6%)
75 to 84	641 (32.8%)
85+	110 (5.6%)
Mean $\pm$ SD	73.7 $\pm$ 5.7
Sex, N (%)	
Female	938 (48.0%)
Male	1,016 (52.0%)
Race/Ethnicity, N (%)	
Asian/Asian-American	53 (2.7%)
Black/African-American	30 (1.5%)
Hawaiian or Pacific Islander	3 (0.2%)
Hispanic/Latinx	64 (3.3%)
Native American or Alaska Native	4 (0.2%)
White, non-Hispanic	1,734 (88.7%)
More than one race/ethnicity or other race/ethnicity	66 (3.4%)
Socioeconomic and education status	
Neighborhood Deprivation Index (NDI, mean $\pm$ SD; min = -1.7788, max = 2.9605)	-0.2 $\pm$ 0.6
NDI categories, N (%)	
Least deprivation (-1.7788 through -0.6141)	550 (28.2%)
Moderate deprivation (-0.6136 through -0.1673)	651 (33.3%)
Highest deprivation (-0.1667 through 2.9605)	753 (38.5%)
Highest education level, N (%)	
High school or less	480 (24.6%)
Some college or 2-year degree	776 (39.7%)
College graduate or higher	698 (35.7%)
Clinical characteristics	
HbA1c; 1 <sup>st</sup> value in the 12 months before index date (N, %)	
HbA1c < 8%	1,663 (85.1%)
HbA1c $\geq$ 8%	291 (14.9%)
Years on T2DM Registry before index date (mean $\pm$ SD; min = 1.0 max = 29.2)	11.4 $\pm$ 7.1
Self-reported general health, N, (%)	
Fair/poor	588 (30.1%)
Good, very good, excellent	1,366 (69.9%)
Self-reported mental health, N (%)	
Fair/poor	363 (18.6%)
Good, very good, excellent	1,591 (81.4%)
Comorbidities and prior health care utilization	
Charlson Co-morbidity Index, N (%)	
0	393 (20.1%)
1	363 (18.6%)
2+	1,198 (61.3%)
Prior DM-specific ED utilization N (%)	
1+ DM-related ED visits in 12 months before index date	381 (19.5%)
0 DM-related ED visits in 12 months before index date	1,573 (80.5%)
Prior DM-specific hospital utilization, N (%)	
1+ DM-related hospital admissions in 12 months prior	181 (9.3%)
0 DM-related hospital admissions in 12 months prior	1,773 (90.7%)

*Abbreviations:* ED, Emergency Department; DM, Diabetes Mellitus; NDI, Neighborhood Deprivation Index; T2DM, Type 2 Diabetes Mellitus.

**Table 3. Description of Social Needs and Outcome Measures in Population**

Need Characteristics	N (%)
Individual needs identified on MTHA	
Financial strain	187 (9.6%)
Food insecurity	95 (4.9%)
Housing instability	61 (3.1%)
Social isolation	605 (31.0%)
Transportation needs	131 (6.7%)
Number of needs identified on MTHA	
0	1,174 (60.1%)
1	580 (29.7%)
2	132 (6.8%)
3+	68 (3.5%)
Total needs (mean $\pm$ SD)	0.6 $\pm$ 0.8

Outcome Measures (1-year post Index Date)	N (%)
HbA1c < 8%	1,578 (80.8%)
Any DM-specific ED utilization	456 (23.3%)
Any DM-specific Hospital admissions	224 (11.5%)

*Abbreviations:* MTHA, Medicare Total Health Assessment; ED, emergency department; DM, Diabetes Mellitus.

associated with DM-related hospital admissions in the year following assessment.

Our study results complement previous research examining social needs and health outcomes among older people with diabetes. As in our study, Ryan et al. (2023)<sup>13</sup> found that loneliness and transportation needs were associated with higher ED utilization. In addition consistent with our findings, Massey et al. (2023)<sup>25</sup> found that food insecurity was associated with poorer glycemic control, while

Walker et al. (2021)<sup>26</sup> found that increased financial strain was associated with worse glycemic control. While previous research found that lack of housing was associated with poorer glycemic control in a veteran population<sup>27</sup> our study seems to be one of the first to find an association between housing instability and worse glycemic control in a Medicare population.

Our finding that food insecurity was associated with increased ED utilization is similar to other research findings in the published literature. Although not DM-specific,<sup>28</sup> Estrella and colleagues (2021) found that those with food insecurity were more likely to be frequent users of the ED. While the exact causal mechanisms are unclear, food insecurity among those with T2DM may lead to poor disease management, leading to complications resulting in increased ED utilization.<sup>29</sup>

The results of our study make a unique contribution to the literature on social determinants of health and T2DM outcomes. First, this study examined the association between social needs and DM-specific outcomes on a population level among Medicare patients seeking wellness visits in a large, integrated health care system. Second, this study examined a comprehensive set of social needs, and followed outcomes prospectively in the year after social needs assessment.

While we cannot determine if the relationship between social needs and study outcomes is causal due to the observational study design used in the analysis, our results have implications for clinical care and health care delivery. Based on our findings, further multi-level, multi-sector interventions are needed to potentially improve outcomes for populations with T2DM and social needs. These

**Table 4. Logistic Regression Results: Association Between Individual Social Needs and DM-Specific Outcomes**

Logistic Regression Models	Hb A1c < 8%		DM-Specific ED Utilization		DM-Specific Hospital Admissions	
	Or	95% CI	Or	95% CI	Or	95% CI
Individual social needs (yes vs no)						
Model 1: Financial Strain	<b>0.56</b>	<b>0.36-0.85</b>	<b>1.65</b>	<b>1.17-2.33</b>	<b>1.77</b>	<b>1.17-2.68</b>
Model 2: Food Insecurity	<b>0.47</b>	<b>0.28-0.81</b>	<b>1.62</b>	<b>1.02-2.57</b>	1.20	0.67-2.17
Model 3: Housing Instability	<b>0.50</b>	<b>0.25-0.99</b>	<b>2.14</b>	<b>1.23-3.75</b>	0.80	0.35-1.81
Model 4: Social Isolation	0.91	0.67-1.24	<b>1.36</b>	<b>1.06-1.74</b>	1.05	0.76-1.46
Model 5: Transportation Needs	1.06	0.63-1.80	<b>1.83</b>	<b>1.23-2.71</b>	1.34	0.81-2.21

<sup>1</sup>Models adjusted for age, sex, race/ethnicity, NDI, education level, prior HbA1c status ( $\leq 8$ ), years on T2DM Registry, self-reported general (overall) health, self-reported mental health, CCI, prior DM-specific ED visits and prior DM-specific hospital admissions.

*Abbreviations:* ED, Emergency Department; DM, Diabetes Mellitus; CCI, Charlson Comorbidity Index; NDI, Neighborhood Deprivation Index.

interventions should be built on the CMS Accountable Health Communities (AHC) Model implemented in 2016.<sup>30</sup> Core components of the AHC model include social needs screening, patient navigation and connection to community-based services, and engagement with community stakeholders to ensure sufficient services are available to address unmet social needs. Further research is warranted on how such interventions impact health outcomes for older adults with T2DM.

The current study has several limitations. First, study results may not be generalizable to settings with different Medicare insurance models, such as noncapitated health care environments (eg, fee-for-service). Second, data on social needs were collected via self-report and may have social desirability biases. Third, the dichotomization of social needs measures limit the ability to assess potential “dose effects” of social needs on study outcomes. Fourth, the population studied was mostly white and non-Hispanic, and was not representative of the US population as a whole. Fifth, results may not be generalizable to noninsured patients or those residing in neighborhoods that have high levels of neighborhood deprivation. In addition, although the relationship between social needs and study outcomes likely did not change during the COVID-19 pandemic period, there may have been more older adults with social needs during this period compared with before the COVID-19 pandemic period. Last, while the population studied is representative of the KPNW Medicare population seeking wellness visits, it may not be generalizable to the KPNW Medicare population overall.

## Conclusions and Learnings for Clinical Practice

Our study found that social needs were associated with worse DM-specific outcomes. These results have clear implications for clinical practice. Our study findings suggest the importance of developing further multi-level, multi-sector interventions to address social needs for patients with T2DM. Further research is warranted on how such interventions impact health outcomes for older adults with T2DM.

We acknowledge Jill Pope and Neon Brooks for providing editing services to the project and Robin Daily and Mary Early for

providing administrative support for the submission of this manuscript.

To see this article online, please go to: <http://jabfm.org/content/38/1/125.full>.

## References

- Centers for Disease Control and Prevention. National Diabetes Statistics Report 2020. Available at: <https://www.cdc.gov/diabetes/pdfs/data/statistics/national-diabetes-statistics-report.pdf>. Accessed March 1, 2024.
- Parker ED, Lin J, Mahoney T, et al. Economic costs of diabetes in the U.S. in 2022. *Diabetes Care* 2024;47:26–43.
- National Academies of Sciences Engineering and Medicine. *Social Isolation and Loneliness In Older Adults: Opportunities for The Health Care System*. National Academies Press; 2020.
- Feeding America. The state of senior hunger in 2021. 2023; Available at: <https://www.feedingamerica.org/research/state-senior-hunger>. Accessed May 26, 2023.
- Marshall GL, Bayaz-Ozturk G, Kahana E, Gallo WT, Seghal A. Dynamics of financial hardship in the United States: health and retirement study 2006–2016. *J Gerontol Soc Work* 2022;65:241–51.
- Drake C, Alfaro JM, Blalock DV, et al. Association of unmet social needs with metformin use among patients with type 2 diabetes. *Diabetes Care* 2023;46:2044–9.
- Chambers EC, McAuliff KE, Heller CG, Fiori K, Hollingsworth N. Toward understanding social needs among primary care patients with uncontrolled diabetes. *J Prim Care Community Health* 2021;12:2150132720985044.
- Gold R, Kaufmann J, Gottlieb LM, et al. Cross-sectional associations: social risks and diabetes care quality, outcomes. *Am J Prev Med* 2022;63:392–402.
- Fitzpatrick SL, Banegas MP, Kimes TM, Papajorgji-Taylor D, Fuoco MJ. Prevalence of unmet basic needs and association with diabetes control and care utilization among insured persons with diabetes. *Popul Health Manag* 2021;24:463–9.
- Walker RJ, Smalls BL, Campbell JA, Strom Williams JL, Egede LE. Impact of social determinants of health on outcomes for type 2 diabetes: a systematic review. *Endocrine* 2014;47:29–48.
- Berkowitz SA, Meigs JB, DeWalt D, et al. Material need insecurities, control of diabetes mellitus, and use of health care resources: results of the Measuring Economic Insecurity in Diabetes Study. *JAMA Intern Med* 2015;175:257–65.
- Jiang DH, O'Connor PJ, Huguet N, Golden SH, McCoy RG. Modernizing diabetes care quality measures. *Health Aff (Millwood)* 2022;41:955–62.
- Ryan JL, Franklin SM, Canterberry M, et al. Association of health-related social needs with

- quality and utilization outcomes in a medicare advantage population with diabetes. *JAMA Netw Open* 2023;6:e239316.
14. Hill-Briggs F, Adler NE, Berkowitz SA, et al. Social determinants of health and diabetes: a scientific review. *Diabetes Care* 2020;44:258–79.
15. Canterberry M, Figueroa JF, Long CL, et al. Association between self-reported health-related social needs and acute care utilization among older adults enrolled in Medicare Advantage. *JAMA Health Forum* 2022;3:e221874.
16. Mosen DM, Banegas MP, Keast EM, Dickerson JF. Examining the association of social needs with future health care utilization in an older adult population: which needs are most important? *Popul Health Manag* 2023;26:413–9.
17. Mosen DM, Banegas MP, Tucker-Seeley RD, et al. Social isolation associated with future health care utilization. *Popul Health Manag* 2021;24:333–7.
18. Mosen DM, Banegas MP, Friedman N, Shuster E, Brooks N. Food insecurity associated with self-reported falls among Medicare Advantage members. *Popul Health Manag* 2019;22:536–9.
19. Mosen DM, Pihlstrom DJ, Snyder JJ, Shuster E. Assessing the association between receipt of dental care, diabetes control measures and health care utilization. *J Am Dent Assoc* 2012;143:20–30.
20. Ross TR, Ng D, Brown JS, et al. The HMO Research Network Virtual Data Warehouse: a public data model to support collaboration. *EGEMS (Wash DC)* 2014;2:1049.
21. Strain WD, Down S, Brown P, Puttanna A, Sinclair A. Diabetes and frailty: an expert consensus statement on the management of older adults with type 2 diabetes. *Diabetes Ther* 2021;12:1227–47.
22. Messer LC, Laraia BA, Kaufman JS, et al. The development of a standardized neighborhood deprivation index. *J Urban Health* 2006;83:1041–62.
23. Kieszak SM, Flanders WD, Kosinski AS, Shipp CC, Karp H. A comparison of the Charlson comorbidity index derived from medical record data and administrative billing data. *J Clin Epidemiol* 1999;52:137–42.
24. Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol* 1992;45:613–9.
25. Massey M, Stewart MP, LaManna JB, Park C, Ng BP. Food insecurity and glycemic goals among Medicare beneficiaries with type 2 diabetes. *Chronic Illn* 2023;17423953231217346.
26. Walker RJ, Garacci E, Campbell JA, Harris M, Mosley-Johnson E, Egede LE. Relationship between multiple measures of financial hardship and glycemic control in older adults with diabetes. *J Appl Gerontol* 2021;40:162–9.
27. Axon RN, Gebregziabher M, Dismuke CE, et al. Differential impact of homelessness on glycemic control in veterans with type 2 diabetes mellitus. *J Gen Intern Med* 2016;31:1331–7.
28. Estrella A, Scheidell J, Khan M, et al. Cross-sectional analysis of food insecurity and frequent emergency department use. *West J Emerg Med* 2021;22:911–8.
29. Levi R, Bleich SN, Seligman HK. Food insecurity and diabetes: overview of intersections and potential dual solutions. *Diabetes Care* 2023;46:1599–608.
30. Alley DE, Asomugha CN, Conway PH, Sanghavi DM. Accountable health communities—addressing social needs through Medicare and Medicaid. *N Engl J Med* 2016;374:8–11.