


240^{TOTAL}

<35^{HDL}

Low HDL with elevated LDL and triglycerides: A common denominator of many heart attack victims

Mixed hyperlipidemias—elevated cholesterol *and* triglycerides—are common among heart attack victims, and nearly two thirds of people who developed myocardial infarction in the PROCAM trial had a low (< 35 mg/dL) baseline level of HDL cholesterol. TOPIID (gemfibrozil) is not indicated for the treatment of patients with low HDL cholesterol as their only lipid abnormality.

HEART ATTACK PATIENTS (PROCAM TRIAL)*



A powerful case for

LOPID[®]



BID

(gemfibrozil) **600-mg
Tablets**

LOPID is indicated for reducing the risk of coronary heart disease in type IIb patients with low HDL, in addition to elevated LDL and triglycerides, and who have had an inadequate response to weight loss, diet, exercise, and other pharmacologic agents such as bile acid sequestrants and nicotinic acid.

Raised low HDL 25%

—in patients whose baseline HDL was < 35 mg/dL and median baseline LDL was 186 mg/dL in the landmark Helsinki Heart Study (HHS).³

Reduced heart attack incidence up to 62%*

—in these HHS patients.³ Incidence of serious coronary events was similar for LOPID and placebo subgroups with baseline HDL above the median (46.4 mg/dL).³

RAISES HDL, LOWERS LDL AND TRIGLYCERIDES DRAMATICALLY REDUCES HEART ATTACK

Contraindicated in patients with hepatic or severe renal dysfunction, including primary biliary cirrhosis, preexisting gallbladder disease, or hypersensitivity to gemfibrozil. LOPID may increase cholesterol secretion into the bile, leading to cholelithiasis. Caution should be exercised when anticoagulants are given in conjunction with LOPID.

*Defined as a combination of definite coronary death and/or definite myocardial infarction.
P = .013; 95% CI 13.3 to 111.5.

References 1. Goldstein JL, Hazzard WR, Schrott HG, Bierman EL, Motulsky AG. Hyperlipidemia in coronary heart disease. I. Lipid levels in 500 survivors of myocardial infarction. *J Clin Invest.* 1973;52:1533-1543.
2. Assmann G, Schulte H. *PROCAM-Trial: Prospective Cardiovascular Münster Trial.* Zürich: Panscientia Verlag; 1986:8-9. 3. Data on file, Medical Affairs Dept, Parke-Davis.

Please see last page of this advertisement for warnings, contraindications, and brief summary of prescribing information.

Lopid® (Gemfibrozil Capsules and Tablets)

Before prescribing, please see full prescribing information. A Brief Summary follows.

CONTRAINDICATIONS. 1. Hepatic or severe renal dysfunction, including primary biliary cirrhosis.

2. Preexisting gallbladder disease (See WARNINGS).
3. Hypersensitivity to gemfibrozil.

WARNINGS. 1. Because of chemical, pharmacological, and clinical similarities between gemfibrozil and clofibrate, the adverse findings with clofibrate in two large clinical studies may also apply to gemfibrozil. In the first of those studies, the Coronary Drug Project, 1000 subjects with previous myocardial infarction were treated for five years with clofibrate. There was no difference in mortality between the clofibrate-treated subjects and 3000 placebo-treated subjects, but twice as many clofibrate-treated subjects developed cholelithiasis and cholecystitis requiring surgery. In the other study, conducted by the World Health Organization (WHO), 5000 subjects without known coronary heart disease were treated with clofibrate for five years and followed one year beyond. There was a statistically significant, 29%, higher total mortality in the clofibrate-treated than in a comparable placebo-treated control group. The excess mortality was due to a 33% increase in noncardiovascular causes, including malignancy, post-cholecystectomy complications, and pancreatitis. The higher risk of clofibrate-treated subjects for gallbladder disease was confirmed.

During the Helsinki Heart Study and in the 1½ year follow-up period since the trial was completed, mortality from any cause was 59 (2.9%) in the Lopid group and 55 (2.7%) in the placebo group. Mortality from any cause during the double-blind portion of the study was 44 deaths in the Lopid group and 43 in the placebo group. Because of the more limited size of the Helsinki Heart Study, this result is not statistically significantly different from the 29% excess mortality seen in the clofibrate group in the separate WHO study. Noncoronary heart disease related mortality showed a 58% greater trend in the Lopid group (43 vs 27 patients in the placebo group, $p=0.056$).

In the Helsinki Heart Study, the incidence of total malignancies discovered during the trial and in the 1½ years since the trial was completed was 39 in the Lopid group and 29 in the placebo group (difference not statistically significant). This includes 5 basal cell carcinomas in the Lopid group and none in the placebo group ($p=0.06$; historical data predicted an expected 4.7 cases in the placebo group). GI malignancies and deaths from malignancies were not statistically different between Lopid and placebo subgroups. Follow-up of the Helsinki Heart Study participants will provide further information on cause-specific mortality and cancer morbidity.

2. A gallstone prevalence substudy of 450 Helsinki Heart Study participants showed a trend toward a greater prevalence of gallstones during the study within the Lopid treatment group (75% vs 49% for the placebo group, a 55% excess for the gemfibrozil group). A trend toward a greater incidence of gallbladder surgery was observed for the Lopid group (17 vs 11 subjects, a 54% excess). This result did not differ statistically

from the increased incidence of cholecystectomy observed in the WHO study in the group treated with clofibrate. Both clofibrate and gemfibrozil may increase cholesterol excretion into the bile leading to cholelithiasis. If cholelithiasis is suspected, gallbladder studies are indicated. Lopid therapy should be discontinued if gallstones are found.

3. Since a reduction of mortality from coronary artery disease has not been demonstrated and because liver and interstitial cell testicular tumors were increased in rats, Lopid should be administered only to those patients described in the INDICATIONS AND USAGE section. If a significant serum lipid response is not obtained, Lopid should be discontinued.

4. Concomitant Anticoagulants—Caution should be exercised when anticoagulants are given in conjunction with Lopid. The dosage of the anticoagulant should be reduced to maintain the prothrombin time at the desired level to prevent bleeding complications. Frequent prothrombin determinations are advisable until it has been definitely determined that the prothrombin level has stabilized.

5. Concomitant therapy with Lopid and Mevacor® (lovastatin) has been associated with rhabdomyolysis, markedly elevated creatine kinase (CK) levels and myoglobinuria, leading in a high proportion of cases to acute renal failure. In most subjects who have had an unsatisfactory lipid response to either drug alone, the possible benefit of combined therapy with lovastatin and gemfibrozil does not outweigh the risks of severe myopathy, rhabdomyolysis, and acute renal failure (See Drug Interactions). The use of fibrates alone, including Lopid, may occasionally be associated with myositis. Patients receiving Lopid and complaining of muscle pain, tenderness, or weakness should have prompt medical evaluation for myositis, including serum creatine kinase level determination. If myositis is suspected or diagnosed, Lopid therapy should be withdrawn.

6. Cataracts—Subcapsular bilateral cataracts occurred in 10%, and unilateral in 6.3% of male rats treated with gemfibrozil at 10 times the human dose.

PRECAUTIONS. 1. **Initial Therapy**—Laboratory studies should be done to ascertain that the lipid levels are consistently abnormal. Before instituting Lopid therapy, every attempt should be made to control serum lipids with appropriate diet, exercise, weight loss in obese patients, and control of any medical problems such as diabetes mellitus and hypothyroidism that are contributing to the lipid abnormalities.

2. **Continued Therapy**—Periodic determination of serum lipids should be obtained, and the drug withdrawn if lipid response is inadequate after 3 months of therapy.

3. **Drug Interactions**—(A) **Lovastatin:** Rhabdomyolysis has occurred with combined gemfibrozil and lovastatin therapy. It may be seen as early as 3 weeks after initiation of combined therapy or after several months. In most subjects who have had an unsatisfactory lipid response to either drug alone, the possible benefit of combined therapy with lovastatin and gemfibrozil does not outweigh the risks of severe myopathy, rhabdomyolysis, and acute renal failure. There is no assurance that periodic monitoring of creatine kinase will prevent the occurrence of severe myopathy and kidney damage.

(B) **Anticoagulants:** CAUTION SHOULD BE EXERCISED WHEN ANTICOAGULANTS ARE GIVEN IN CONJUNCTION WITH LOPID. THE DOSAGE OF THE ANTICOAGULANT SHOULD BE REDUCED TO MAINTAIN THE PROTHROMBIN TIME AT THE DESIRED LEVEL TO PREVENT BLEEDING COMPLICATIONS. FREQUENT PROTHROMBIN DETERMINATIONS ARE ADVISABLE UNTIL IT HAS BEEN DEFINITELY DETERMINED THAT THE PROTHROMBIN LEVEL HAS STABILIZED.

4. **Carcinogenesis, Mutagenesis, Impairment of Fertility**—Long-term studies have been conducted in rats and mice at one and ten times the human dose. The incidence of benign liver nodules and liver carcinomas was significantly increased in high dose male rats. The incidence of liver carcinomas increased also in low dose males, but this increase was not statistically significant ($p=0.1$). In high dose female rats, there was a significant increase in the combined incidence of benign, and malignant liver neoplasms. In male and female mice, there were no statistically significant differences

Lopid® (Gemfibrozil Capsules and Tablets)

from controls in the incidence of liver tumors, but the doses tested were lower than those shown to be carcinogenic with other fibrates.

Male rats had a dose-related and statistically significant increase of benign Leydig cell tumors at 1 and 10 times the human dose.

Electron microscopy studies have demonstrated a florid hepatic peroxisome proliferation following Lopid administration to the male rat. An adequate study to test for peroxisome proliferation has not been done in humans but changes in peroxisome morphology have been observed. Peroxisome proliferation has been shown to occur in humans with either of two other drugs of the fibrate class when liver biopsies were compared before and after treatment in the same individual.

Administration of approximately three or ten times the human dose to male rats for 10 weeks resulted in a dose-related decrease of fertility. Subsequent studies demonstrated that this effect was reversed after a drug-free period of about eight weeks, and it was not transmitted to the offspring.

5. **Pregnancy Category B**—Reproduction studies have been performed in the rat at doses 3 and 9 times the human dose, and in the rabbit at 2 and 6.7 times the human dose. These studies have revealed no evidence of impaired fertility in females or harm to the fetus due to Lopid. Minor fetotoxicity was manifested by reduced birth rates observed at the high dose levels. No significant malformations were found among almost 400 offspring from 36 litters of rats and 100 fetuses from 22 litters of rabbits.

There are no studies in pregnant women. In view of the fact that Lopid is tumorigenic in male and female rats, the use of Lopid in pregnancy should be reserved for those patients where the benefit clearly outweighs the possible risk to the patient or fetus.

6. **Nursing Mothers**—Because of the potential for tumorigenicity shown for gemfibrozil in rats, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

7. **Hematologic Changes**—Mild hemoglobin, hematocrit and white blood cell decreases have been observed in occasional patients following initiation of Lopid therapy. However, these levels stabilize during long-term administration. Rarely, severe anemia, leukopenia, thrombocytopenia, and bone marrow hypoplasia have been reported. Therefore, periodic blood counts are recommended during the first 12 months of Lopid administration.

8. **Liver Function**—Abnormal liver function tests have been observed occasionally during Lopid administration, including elevations of AST (SGOT), ALT (SGPT), LDH, bilirubin, and alkaline phosphatase. These are usually reversible when Lopid is discontinued. Therefore periodic liver function studies are recommended and Lopid therapy should be terminated if abnormalities persist.

9. **Use in Children**—Safety and efficacy in children have not been established.

ADVERSE REACTIONS. In the double-blind controlled phase of the Helsinki Heart Study, 2046 patients received Lopid for up to 5 years. In that study, the following adverse reactions were statistically more frequent in subjects in the Lopid group (placebo incidence in parentheses): gastrointestinal reactions, 34.2%

(23.8%); dyspepsia, 19.6% (11.9%); abdominal pain, 9.8% (5.6%); acute appendicitis (histologically confirmed in most cases where data are available), 1.2% (0.6%); atrial fibrillation, 0.7% (0.1%).

Adverse events reported by more than 1% of subjects, but without a significant difference between groups (placebo incidence in parentheses) were: diarrhea, 7.2% (6.5%); fatigue, 3.8% (3.5%); nausea/vomiting, 2.5% (2.1%); eczema, 1.9% (1.2%); rash, 1.7% (1.3%); vertigo, 1.5% (1.3%); constipation, 1.4% (1.3%); headache, 1.2% (1.1%).

Gallbladder surgery was performed in 0.9% of Lopid and 0.5% of placebo subjects, a 64% excess, which is not statistically different from the excess of gallbladder surgery observed in the clofibrate compared to the placebo group of the WHO study.

Nervous system and special senses adverse reactions were more common in the Lopid group. These included hypesthesia, paresthesias, and taste perversion. Other adverse reactions that were more common among Lopid treatment group subjects but where a causal relationship was not established include cataracts, peripheral vascular disease, and intracerebral hemorrhage.

From other studies it seems probable that Lopid is causally related to the occurrence of **musculoskeletal symptoms** (See WARNINGS), and to **abnormal liver function tests and hematologic changes** (See PRECAUTIONS).

Reports of viral and bacterial infections (common cold, cough, urinary tract infections) were more common in gemfibrozil-treated patients in other controlled clinical trials of 805 patients.

Additional adverse reactions that have been reported for gemfibrozil are listed below by system. These are categorized according to whether a causal relationship to treatment with Lopid is probable or not established:

CAUSAL RELATIONSHIP PROBABLE: *Gastrointestinal:* cholestatic jaundice; *Central Nervous System:* dizziness, somnolence, paresthesia, peripheral neuritis, decreased libido, depression, headache; *Eye:* blurred vision; *Genitourinary:* impotence;

Musculoskeletal: myopathy, myasthenia, myalgia, painful extremities, arthralgia, synovitis, rhabdomyolysis (see WARNINGS and Drug Interactions under PRECAUTIONS); *Clinical Laboratory:* increased creatine phosphokinase, increased bilirubin, increased liver transaminases (AST [SGOT], ALT [SGPT]), increased alkaline phosphatase; *Hematopoietic:* anemia, leukopenia, bone marrow hypoplasia, eosinophilia; *Immunologic:* angioedema, laryngeal edema, urticaria; *Integumentary:* exfoliative dermatitis, rash, dermatitis, pruritus.

CAUSAL RELATIONSHIP NOT ESTABLISHED: *General:* weight loss; *Cardiac:* extrasystoles; *Gastrointestinal:* pancreatitis, hepatoma, colitis; *Central Nervous System:* confusion, convulsions, syncope; *Eye:* retinal edema; *Genitourinary:* decreased male fertility; *Clinical Laboratory:* positive antinuclear antibody; *Hematopoietic:* thrombocytopenia; *Immunologic:* anaphylaxis, Lupus-like syndrome, vasculitis; *Integumentary:* alopecia.

DOSAGE AND ADMINISTRATION. The recommended dose for adults is 1200 mg administered in two divided doses 30 minutes before the morning and evening meal.

MANAGEMENT OF OVERDOSE. While there has been no reported case of overdose, symptomatic supportive measures should be taken should it occur.

References: 1. Frick MH, Elo O, Haapa K, et al: Helsinki Heart Study: Primary prevention trial with gemfibrozil in middle-aged men with dyslipidemia. *N Engl J Med* 1987;317:1237-1245. 2. Manninen V, Elo O, Frick MH, et al: Lipid alterations and decline in the incidence of coronary heart disease in the Helsinki Heart Study. *JAMA* 1988; 260:641-651. 3. Nikkila EA: Familial lipoprotein lipase deficiency and related disorders of chylomicron metabolism. In Stanbury J. B. et al. (eds): *The Metabolic Basis of Inherited Disease*, 5th ed., McGraw-Hill, 1983, Chap. 30, pp. 622-642.

Caution—Federal law prohibits dispensing without prescription.

PARKE-DAVIS
Div of Warner-Lambert Co
Morris Plains, NJ 07950 USA

0737G013

PD-56-JA-676-B1(031)

LOPID® *Lopid*
BID
(gemfibrozil) 600-mg Tablets

**RAISES HDL, LOWERS LDL AND TRIGLYCERIDES
DRAMATICALLY REDUCES HEART ATTACK**

▼
**You're
 looking at
 the answer
 to your
 recruitment
 problems!**

JABFP
Volume 18 Number 4

The Journal of the American Board of Family Practice

Outpatient Antimicrobial Susceptibility	225
Postural Hypotension in the Elderly	234
EMR Use for Patient Health	246
Macronutrients	252
Nonsteroidal Anti-Inflammatory Drugs	257
Macronutrient Intake	279



It's the *Journal of the American Board of Family Practice*. Where you reach the best job candidates in Family Practice. More of them. More often. To find out more, just contact Classified Advertising at (617) 893-3800.



The Journal of the American Board of Family Practice



MICRO-K10
 (Potassium Chloride)
 Extended-Release Capsules, USP 750 mg
 (70 mEq K)



The appearance of the capsule is a registered trademark of A.H. Robins.

© 1991, Wyeth-Ayerst Laboratories

JUST PUBLISHED!

Many physicians are finding it difficult to keep current on the effectiveness and safety of prescribed products.


Drug Therapy, Volume 8 can help you review changes in drug use, the implications of using drugs in combination, and the comparative efficacy of different drugs.

Drug Therapy, Volume 8 includes 24 review articles originally published in the *New England Journal of Medicine* from 1985 to 1990, plus related Correspondence and Addenda by the original authors, providing updates from new research and clinical experience.

Drug Therapy, Volume 8

© 1991. □ Edited by John A.

Oates, M.D. and Alastair J.J. Wood, M.D., Department of Medicine, Vanderbilt University School of Medicine □ 300 pages, including index and over 75 charts & illustrations.



CONTENTS:

ANTIARRHYTHMICS

(Encaïnide, Flecainide, Tocainide, Amiodarone, Mexiletine)

VASODILATORS AND ANTITHROMBOTICS

(Combined Beta-Adrenergic and Calcium-Entry Blockade in Angina Pectoris, Nitrate Therapy in Stable Angina Pectoris, Converting-Enzyme Inhibitors, Tissue Plasminogen Activator, Dipyridamole)

ANTIBIOTICS

(Beta-Lactam Antibiotics, Bactericidal Activity as a Monitor of Antibiotic Therapy, Antimicrobial Prophylaxis)

ANTICANCER DRUGS

(Etoposide, Ketoconazole as an Inhibitor of Steroid Production)

BRONCHODILATORS

Ipratropium Bromide, New Approach to the Treatment of Asthma)

OTHERS

(Drug Therapy in the Elderly, Cyclosporine, Oral Hypoglycemic Agents, AZT (Zidovudine) and Related Dideoxynucleosides, Physiologic Replacement of Insulin, HMG-CoA Reductase Inhibitors, Cigarette Smoking and Nicotine Addiction)

THE LATEST REPRINT COLLECTION FROM THE NEW ENGLAND JOURNAL OF MEDICINE . . .

Yes, please send me
Drug Therapy, Volume 8:

Softcover, ISBN 0-910133-23-9,
@ \$34.00 + \$3.50 postage &
handling per copy. _____copies.

Hardcover, ISBN 0-910133-33-6,
@ \$49.00 + \$3.95 postage &
handling per copy. _____copies.

TOLL FREE (800) THE-NEJM
Inside MA: (617) 893-3800 x1199
FAX: (617) 893-0413

Enclosed is a check for \$ _____*

Please charge my Visa MasterCard AmEx in the amount of \$ _____.

Card# Exp. Date

Signature _____

Name/Specialty _____
(please print)

Organization _____

Address _____

City _____ State _____ ZIP _____

*Prepaid orders only. Please make payable to the *New England Journal of Medicine*. MA residents please add 5% sales tax. Allow 4-6 weeks for delivery. Prices subject to change without notice.

30-day return privileges guarantee your satisfaction.



The New England Journal of Medicine
P.O. Box 9130, Waltham, MA 02254-9130

SBD08

INFORMATION FOR READERS

THE JOURNAL OF THE AMERICAN BOARD OF FAMILY PRACTICE
2228 Young Drive
Lexington, KY 40505

*Official Publication of
The American Board of Family Practice*

John P. Geyman, M.D., *Editor*
Paul R. Young, M.D., *Executive Editor*
Alfred O. Berg, M.D., *Associate Editor*
Paul Brucker, M.D., *Associate Editor*
G. Gayle Stephens, M.D., *Associate Editor*
Claire Z. Fenwick, *Assistant Editor*
Ann Stockham, *Copy Editor and Assistant Executive Editor*
Virginia M. Gessner, *Senior Editorial Assistant*
Debbie Wilson, *Production Assistant*

PUBLISHING SERVICES

Publishing Division,
Massachusetts Medical Society
Robert D. Bovenshulte, *Vice President for Publishing*

Customer Service
M. Dolores Fletcher, *Director*

Circulation and Product Marketing
Laurie Priano, *Director*

Electronic Production
William H. Paige, *Director*
Ruth Goodman, *Associate Director*
Martha Soule, *Composition Coordinator*

Manufacturing and Distribution
William H. Paige, *Director*
James T. Clifton, *Assistant Director, Agency and International Services*

Mary Kaye Howe, *Assistant Director, Advertising Production*

Mark Davidson, *Assistant Director, Distribution and Postal Affairs*

International Services
Peter R. Cole, *Director*

Market Research and Analysis
Janet E. Halpern, *Director*

Management Information Services
Larry Altrich, *Director*
Michael McDonald, *Systems Analyst*

Finance
Richard Simoes, *Controller*

Office of Vice President for Publishing
Chris Lynch, *Product Manager*

Alberta L. Fitzpatrick, *Associate Director, Rights and Permissions*

Advertising Sales
Arthur Wilschek, *Director*

Account Managers
Bill Healy, *Midwest*

Lew Wetzels, *Eastern*

Wayne Wickman, *Eastern*

COPYRIGHT

Material appearing in the *Journal of the American Board of Family Practice* is covered by copyright. Copying beyond the quantities permitted under "fair use" as defined by U.S. copyright law is allowed provided the stated fee of \$.20 per page is paid through the Copyright Clearance Center, 21 Congress St., Salem, MA 01970. This consent does not extend to other copying, such as copying for advertising or promotional purposes. Single copies for personal or internal use are allowed at no charge. Nonprofit institutions may make copies provided they obtain prior consent from the *Journal of the American Board of Family Practice*, Rights and Permissions Department, 1440 Main Street, Waltham, MA 02154-1649, (617) 893-3800, ext. 1413.

SUBSCRIPTION INFORMATION AND SERVICES

The *Journal of the American Board of Family Practice* is supplied free of charge to 38,000 Diplomates of the American Board of Family Practice. For information please contact:

American Board of Family Practice
2228 Young Drive
Lexington, KY 40505
Tel: (606) 269-5626
FAX: (606) 266-9699

For all other subscribers please contact:

The Journal of the American Board of Family Practice
Subscription Department
1440 Main Street
Waltham, MA 02154-1649
(617) 893-3800, ext. 1199
Telex: 5106017779 NEJM BOS
FAX: (617) 893-0413

For international subscription information please contact:

The Journal of the American Board of Family Practice
Saxon Way, Melbourn, ROYSTON
Herts, SG8 6NJ, U.K.
Telephone: 07-6326-2368
Telex: 94020513 NEJM G
FAX: 07-6326-2401

SUBSCRIPTION RATES

	Domestic	International*
Institutions	\$58.00	\$60.00
Physicians	\$35.00	\$45.00
Residents/Students	\$20.00	\$45.00

*Pounds Sterling drawn on U.K. banks accepted and converted at current rate of exchange. U.S. dollars drawn on U.S. banks.

OTHER SUBSCRIPTION INFORMATION

Diplomates should make address changes on the form accompanying this issue and forward to the Diplomate address listed above. All other subscribers should forward changes to the Waltham, Mass., address listed on this page. Changes must be received at least six weeks in advance of intended move. Please send new address, old address, and expected date of change.

ISSUES NOT RECEIVED

Missing issues will be replaced for up to three months from the issue date without charge. Diplomates and other subscribers who fail to notify the Lexington, Ky., or the Waltham, Mass., office of address changes will not be eligible for free replacement issues. Claims beyond the three-month limit must be prepaid at the backcopy rates. Claims should be sent to either the Diplomate or regular subscriber address listed on this page.

BACK COPIES

If you wish to purchase back copies (issues published prior to your effective start date) of the *Journal of the American Board of Family Practice*, there is a charge of \$12.50 per issue. Contact the Waltham, Mass., address listed above for information.

REPRINTS

Individual copies of articles are available from the Waltham, Mass., office. If you wish to order bulk reprints (minimum order of 100) please contact the Reprint Department (617) 893-3800, ext. 1279, at the Waltham, Mass., office.

INDEXING AND MICROFORM

The *Journal of the American Board of Family Practice* is indexed in *Index Medicus* and is available in microform from University Microfilms International.

INFORMATION FOR AUTHORS

The *Journal of the American Board of Family Practice* welcomes for editorial review manuscripts that contribute to family practice as a clinical scientific discipline. High priority is given to reports of clinically relevant studies that have practical implications for improved patient care. Manuscripts are considered in relation to the extent to which they represent original work, their significance to the advancement of family medicine, and their interest to the practicing family physician. Some papers that are accepted by the *Journal* will be selected for an accompanying guest editorial or concurrent commentary by other invited authors addressing issues raised by the papers. The *Journal* publishes the following features:

Original Articles. Reports of original research, usually dealing with a clinical, health services, or other clinically relevant study.

Medical Practice. Scholarly articles that relate directly to clinical topics useful in everyday family practice, whether dealing with diagnostic or therapeutic roles of the family physician or reporting studies of what family physicians do in practice.

Clinical Review. In-depth reviews of specific clinical problems, disease entities, or treatment modalities; comprehensive and critical analysis of the literature is required (usual maximum length 5000 words).

Clinical Guidelines and Primary Care. Summaries of major clinical guidelines proposed by various specialty, governmental, or health care organizations, with critical commentary from a primary care perspective.

Special Articles. Articles in other areas that may relate to the role of the family physician, education for family practice, or other subjects important to family practice as a clinical specialty.

Brief Reports. Short reports of pilot studies or case reports with a teaching point of clinical relevance (usual length 1000–1500 words).

Editorial. Focused opinion or commentary that bears on an issue relevant to the field. May or may not accompany an original article in the same issue (usual length 1000–1500 words).

Letters to the Editor. Observations, opinion, or comment on topics under

discussion in the *Journal*, usually not to exceed 500 words.

Book Reviews. Books for review and book reviews should be sent to Dr. John P. Geyman, Editor, the *Journal of the American Board of Family Practice*, Department of Family Medicine (HQ-30), School of Medicine, University of Washington, Seattle, WA 98195.

The following guidelines are in accordance with the "Uniform Requirements for Manuscripts Submitted to Biomedical Journals." The current (fourth) edition was published in the February 7, 1991, issue of the *New England Journal of Medicine*.

MANUSCRIPT SUBMISSION

Manuscripts containing original material are accepted for consideration with the understanding that neither the article nor any part of its essential substance, tables, or figures has been or will be published or submitted for publication elsewhere before appearing in the *Journal*. This restriction does not apply to abstracts or press reports published in connection with scientific meetings. Copies of any possibly duplicative manuscripts should be submitted to the Editor along with the manuscript that is to be considered by the *Journal*. The *Journal* strongly discourages the submission of more than one article dealing with related aspects of the same study. In almost all cases, a single study is best reported in a single paper.

Submit an original and 3 copies of the complete manuscript, including text pages, legends, tables, references, and glossy prints of figures. Only typed copy, on standard-sized typewriter paper and double-spaced throughout, with margins of at least 2.5 cm, is acceptable. Address all submissions to John P. Geyman, M.D., Editor, the *Journal of the American Board of Family Practice*, Department of Family Medicine (HQ-30), School of Medicine, University of Washington, Seattle, WA 98195. A covering letter should identify the person (with the address and telephone number) responsible for negotiations concerning the manuscript; the letter should make it clear that the final manuscript has been seen and approved by all authors.

The *Journal* expects authors to disclose any commercial associations that might pose a conflict of interest in connection with the submitted article. Consultancies, stock ownership or other equity interests, patent-licensing

arrangements, and other kinds of associations that might involve conflict of interest should be disclosed to the Editor in a covering letter at the time of submission. Such information will be held in confidence while the paper is under review and will not influence the editorial decision. If the manuscript is accepted, the Editor will discuss with the authors how best to disclose the relevant information. Questions about this policy should be directed to the Editor.

MANUSCRIPTS

Titles and Authors' Names

With the manuscript, provide a page giving the title of the paper; a running foot of fewer than 40 letter spaces; the name(s) of the author(s), including first name(s) and academic degree(s); the name of the department and institution in which the work was done; and the name and address of the author to whom reprint requests should be addressed. All funding sources supporting the work should be routinely acknowledged on the title page, as should all institutional or corporate affiliations of the authors. Two to four key words should be submitted with the manuscripts to be used for purposes of classification by subject. Use terms from the Medical Subject Headings from *Index Medicus* when possible.

Abstracts

Use another page to provide an abstract of not more than 200 words. This abstract should be factual, not descriptive, with its content appropriate to the type of paper. For original articles reporting results of studies, a four-paragraph format should be used labeled Background, Methods, Results, and Conclusions. These should briefly describe, respectively, the object of the study, the methods used, the major results, and the author(s) conclusions. Abstracts are not necessary for Brief Reports.

Units of Measure

The *Journal* will print measurements in Système International (SI) and conventional units. Authors should use SI units as their principal system and indicate conventional units in parentheses.

Abbreviations

Except for units of measurement, abbreviations are discouraged. Consult the *Council of Biology Editors Style Manual* (Fifth edition. Bethesda, MD: Council of Biology Editors, 1983) for

lists of standard abbreviations. The first time an abbreviation appears, it should be preceded by the words for which it stands.

Drug Names

Generic names should, in general, be used. If an author so desires, brand names may be inserted in parentheses.

Inclusive Language

Sex bias should be avoided and gender-inclusive language used whenever possible.

References

References must be typed in double spacing and numbered consecutively as they are cited. References first cited in tables or figure legends must be numbered so that they will be in sequence with references cited in the text. The style of references is that of the *Index Medicus*. List all authors when there are 6 or fewer; when there are 7 or more, list the first 6, then "et al." Sample references are as follows:

Standard Journal Article

(List all authors, but if the number exceeds 6, give 6 followed by et al. Note that month and issue number are omitted when a journal has continuous pagination throughout a volume.)

Morrow JD, Margolies GR, Rowland J, Roberts LJ 2nd. Evidence that histamine is the causative toxin of scombroid-fish poisoning. *N Engl J Med* 1991; 324:716-20.

Organization as Author

Clinical Experience Network (CEN). A large-scale, office-based study evaluates the use of a new class of non-sedating antihistamines. A report from CEN. *J Am Board Fam Pract* 1990; 3:241-58.

Book

Rakel RE. Textbook of family practice. 4th ed. Philadelphia: WB Saunders, 1990.

Chapter in Book

Haynes RC Jr. Agents affecting calcification: calcium, parathyroid hor-

mone, calcitonin, vitamin D, and other compounds. In: Gilman AG, Rall TW, Nies AS, Taylor P, editors. Goodman and Gilman's the pharmacological basis of therapeutics. 8th ed. New York: Pergamon Press, 1990.

Government Agency

Schwartz JL. Review and evaluation of smoking cessation methods: the United States and Canada, 1978-1985. Bethesda, MD: Department of Health and Human Services, 1987. (NIH publication no. 87-2940.)

Personal Communications

Numbered references to personal communications, unpublished data, and manuscripts either "in preparation" or "submitted for publication" are unacceptable (see "Permissions"). If essential, such material may be incorporated in the appropriate place in the text.

Tables

Type tables in double spacing on separate sheets, and provide a title for each. For footnotes, use the following symbols, in this sequence: *, †, ‡, §, ||, ¶, **, ††, etc. Excessive tabular data are discouraged. If an article is accepted, the *Journal* will arrange to deposit extensive tables of important data with the National Auxiliary Publications Service (NAPS); we will pay for the deposit and add an appropriate footnote to the text. This service makes microfiche or photocopies of tables available at moderate charges to those who request them.

Illustrations

Figures should be professionally designed. Glossy, black-and-white photographs are requested. Symbols, lettering, and numbering should be clear, and these elements should be large enough to remain legible after the figure has been reduced to fit the width of a single column.

The back of each figure should include the sequence number, the name of the author, and the proper orientation (e.g., "top"). Do not mount the figure on cardboard. Photomicro-

graphs should be cropped to a width of 8 cm, and electron photomicrographs should have internal scale markers.

If photographs of patients are used, either the subjects should not be identifiable or their pictures must be accompanied by written permission to use the figure. Permissions forms are available from the Editor.

Legends for illustrations should be type-written (double-spaced) on a separate sheet and should not appear on the illustrations.

Color illustrations are used from time to time. Send both transparencies and prints for this purpose.

Permissions

Materials taken from other sources must be accompanied by a written statement from both author and publisher giving permission to the *Journal* for reproduction.

Obtain permission in writing from at least one author of papers still in press, of unpublished data, and of personal communications.

REVIEW AND ACTION

Manuscripts are examined by the editorial staff and are usually sent to outside reviewers. Authors will remain anonymous to outside reviewers and vice versa. External statistical review will be accomplished where appropriate. Every effort will be made to complete the review process as expeditiously as possible.

Copyright Transfer Forms


Transfer of copyright to the *Journal* is requested upon acceptance of the material for publication. Copyright transfer is required of all materials to be published in the *Journal*, including Letters to the Editor and Book Reviews.

Reprints


Authors will receive reprint information and rates when they are sent their galley proofs. Reprints ordered at that time will be shipped about 3 weeks after the publication date.

Your Best Lines of Defense Against Disease are

MEASLES (rubella) — By year, United States, 1950-1986



ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — Cases and known deaths, by 8-month periods of report to CDC, United States, 1981-1986



Disease	Total	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-39	40-49	50-59	60+
AIDS	12,922	66	72	77	81	87	91	106	1,028	2,846	990	28
Cholera	1	1	1	1	1	1	1	1	1	1	1	1
Diphtheria	800,881	1	2,224	1	1	1	1	1	1	1	1	1
Scarlet fever	21,222	1	1,168	1	1	1	1	1	1	1	1	1
Measles	29,157	116	156	200	224	235	236	232	210	180	150	100
Measles, unreported	3,965	15	122	134	151	157	157	157	157	157	157	157
Measles, total	4,281	131	178	219	235	236	236	236	236	236	236	236
Measles, reported	2,346	101	133	165	176	179	179	179	179	179	179	179
Measles, unreported	1,935	30	45	54	59	57	57	57	57	57	57	57
Polio (paralytic)	2,181	609	843	183	127	253	10	76	175	801	561	100
Polio (nonparalytic)	4,190	142	368	788	826	1,130	177	84	145	441	281	77
Polio, total	6,371	751	1,211	911	953	1,227	253	221	320	1,242	842	177
Polio, reported	3	3	3	3	3	3	3	3	3	3	3	3
Polio, unreported	6,368	748	1,208	908	950	1,224	250	218	317	1,239	839	174
Rubella (congenital)	151	50	78	48	21	44	86	12	53	251	41	10
Rubella (noncongenital)	11,138	108	1,029	1,881	2,160	2,160	2,160	2,160	2,160	2,160	2,160	2,160
Rubella, total	11,289	158	1,107	1,929	2,181	2,204	2,246	2,272	2,213	2,411	2,171	2,170
Syphilis	22,866	1	1	1	1	1	1	1	1	1	1	1
Syphilis, primary & secondary	22,866	1	1	1	1	1	1	1	1	1	1	1
Typhoid	28	1	1	1	1	1	1	1	1	1	1	1
Typhoid, reported	28	1	1	1	1	1	1	1	1	1	1	1
Typhoid, unreported	0	0	0	0	0	0	0	0	0	0	0	0
Typhoid, total	28	1	1	1	1	1	1	1	1	1	1	1



the Facts

Morbidity & Mortality Weekly Report gives you the facts.

Every week the experts at the Centers for Disease Control report the facts on disease trends, epidemiological reports and health recommendations. Better than mere summaries, these reports (complete with charts, maps and tables) *pinpoint disease trends by region*. You'll know exactly what's happening in your part of the country as it occurs. No other source can offer you this fast breaking information — *as it happens*.
 Subscribe today, and get the facts.

YES! I want the facts on the latest medical trends as reported by the C.D.C. Please enter my one-year subscription (51 weekly issues) to **MMWR**.
SPECIAL BONUS: Subscribe today and receive **FREE** surveillance summaries & supplements as released by the C.D.C.

Rush mine, 1st class delivery — \$ 58.00
 Send mine, 3rd class delivery — \$ 42.00

My payment for _____ is enclosed.*
 Please charge to my credit card:
 MasterCard Visa AmEx

 Expire Date _____
 Signature _____

Name _____
 Address _____
 City _____ State _____ Zip _____
 Medical Specialty _____

Please allow 4-6 weeks for delivery of first issue. Rates subject to change without notice.
 * Make checks payable to **MMWR**.

RISK-FREE GUARANTEE: You may cancel your subscription at any time and receive a full refund on all unmailed issues.

Massachusetts Medical Society, P.O. Box 9120, Waltham, MA 02254-9120
 Credit Card Orders: Call 1 (800) 843-6356