

Health Status Of Illiterate Adults: Relation Between Literacy And Health Status Among Persons With Low Literacy Skills

Barry D. Weiss, M.D., Gregory Hart, M.A., Daniel L. McGee, Ph.D., and Sandra D'Estelle

Abstract: *Background:* In nonindustrialized nations, illiteracy is independently associated with poor health. The objective of this research was to determine whether such a relation exists in the United States.

Methods: One hundred ninety-three persons were randomly selected from a group of adult students enrolled in a publicly funded literacy training program. Subjects' health status was measured with the Sickness Impact Profile (SIP), a behaviorally based measure of sickness-related dysfunction. Subjects' literacy skills were also measured. Multivariate statistical techniques were then used to evaluate the relation between health status and literacy level and to adjust for confounding sociodemographic factors.

Results: The physical health (measured by the SIP) of subjects with extremely low reading levels was poor compared with that of subjects with higher reading levels. The relation between reading level and physical health was statistically significant ($P < 0.002$), even after adjusting for confounding sociodemographic variables. Psychosocial health (measured by the SIP) was poor across all levels of reading skills and was comparable with the psychosocial health of populations with severe psychosocial disability. The relation between reading level and psychosocial health was statistically significant ($P < 0.02$) after adjusting for confounding variables.

Conclusions: In the United States, illiteracy and poor health status are independently associated. (J Am Board Fam Pract 1992; 5:257-64.)

Illiteracy is common in many nonindustrialized nations.¹ In those countries, health status indicators, such as life expectancy and infant-maternal survival rates, all improve as the population's literacy level rises.²⁻¹¹

Illiteracy is also common in the United States. Up to 15 million persons in the United States (10 percent of the adult population) lack basic reading skills, and as many as 27 million more have only rudimentary reading skills that are not sufficient to permit full participation in society's economic and social activities.¹²⁻²¹

It is logical, therefore, to hypothesize that a relation between literacy and health status can

exist also in the United States. No published studies in the United States, however, have evaluated such a relation. Thus, the goal of our research was to determine whether a relation exists between literacy and health status among a group of US adults with poor literacy skills. The effect of potentially confounding sociodemographic variables was evaluated with multivariate statistical techniques.

Methods

Subjects

Subjects for this research were adult students enrolled in the Pima County Adult Education Program (PCAE) in Tucson, AZ. PCAE is a publicly funded program that offers adult basic education, including literacy instruction, at more than 40 sites throughout Pima County. PCAE students are not a random sample of all illiterate adults in Pima County, but such a population is impossible to identify because no governmental or private agencies maintain registries of illiterate individuals. During the study there were 5536 enrollees in PCAE. Sociodemographic characteristics of the

Submitted, revised, 14 January 1992.

From the Department of Family and Community Medicine and the Arizona Cancer Center, University of Arizona College of Medicine, Tucson, and the Adult Education Program of Pima County, Arizona. Address reprint requests to Barry D. Weiss, M.D., Department of Family and Community Medicine, Arizona Health Sciences Center, Tucson, Arizona 85724.

This paper was supported by a grant from the University of Arizona Foundation. Results of this study were presented to the US Agency for Health Care Policy and Primary Care Research Conference, San Diego, California, 14 January 1991.

PCAE students studied in this research are shown in Table 1.

To be eligible for this research, potential subjects had to have reading skills at a grade-equivalent level between 0.0 (total inability to read) and grade 12.9. PCAE staff determined reading levels of all subjects using the Tests of Adult Basic Education.²² In some cases, staff additionally used the Mott Basic Language Skills Program²³ as a supplemental test to clarify reading level.

There were also several language eligibility requirements. Subjects had to speak and understand English well enough to participate in the study. They also had to respond affirmatively when asked whether English had been spoken in their home when they were young children. Thus, individuals were eligible for the study even if other languages had been spoken in their childhood home, as long as English was one of those languages.

Table 1. Demographic Characteristics of Study Population, n = 193.

Demographic Characteristics	Mean (SD)
Age (years)	28.53 ($\pm$ 10.6)
Annual income (\$ thousand)	7.61 ($\pm$ 7.02)
School grade completed	9.86 ($\pm$ 1.96)
	Percent
Sex	
Women	61.1
Marital status	
Married	29.0
Single	50.3
Divorced	20.2
Widowed	0.5
Ethnicity	
Hispanic	53.4
White	29.5
Black	9.8
Native American	6.7
Other	0.6
Occupation	
Blue collar	54.4
Unemployed	45.6
Country of birth	
United States	91.2
Mexico	6.7
Other	2.1
Languages spoken in childhood home	
English only	71.0
English and Spanish	26.9
Native American and English	1.6
Other and English	0.5

We excluded persons in whose homes English had not been spoken from early childhood regardless of their ability to read or write in another language. Such individuals could have been fully literate in Spanish, for example, despite poor reading skills in English. We excluded these persons to assure that literacy (and not English language literacy) was truly the variable being analyzed.

All PCAE students are at least 16 years old. We excluded the small number of PCAE students who are mentally retarded and those with known learning disabilities.

We selected subjects using stratified random sampling. Stratification was done by classroom (each class contained students of similar reading level) to assure that the final distribution of subjects would be representative of the overall distribution of reading levels among PCAE students.

Subjects were selected from within each class using a table of random numbers. If the chosen student met eligibility criteria, that student was asked to participate. If the individual was ineligible or unwilling to participate, the numbers table was used to select an alternate subject.

Subjects received a small payment for participation. The Human Subjects Committee at the University of Arizona College of Medicine reviewed and approved this research.

Health Status Measure

We used the Sickness Impact Profile (SIP) to measure health status.²⁴ The SIP has been used to quantify health status for a wide variety of populations and medical conditions, including healthy patients enrolled in prepaid health plans²⁵ and patients with chronic medical conditions, such as rheumatoid arthritis,²⁶ coronary artery disease,²⁷⁻²⁹ cancer,^{30,31} chronic obstructive pulmonary disease,³²⁻³⁴ back pain,^{35,36} and thyroid dysfunction.³⁷

The SIP is a behaviorally based measure of sickness-related dysfunction. SIP includes 136 items covering 12 categories of daily activity: ambulation, mobility, body care, social interaction, communication, alertness, emotional behavior, sleep, eating, work, home management, and recreational pastimes. A person's SIP responses are scored to yield quantitative ratings for a "physical dimension" and a "psychosocial dimension" of health. A total composite (overall) score can also

be computed. SIP scores have a high interrater reliability (0.92), test-retest reliability (0.88 to 0.92), and internal consistency (0.96).³⁸

There is no normal range for the SIP, but lower scores indicate better health. Persons from healthy populations typically have scores below 3.0, and patients with profound health-related disabilities often have scores as high as 20.0 to 30.0. It is unlikely, however, that a clinician would note an obvious difference between a patient with an SIP score of 2.0 and a patient whose score, for example, is 5.0. Both patients would appear healthy, even though the individual with the higher score can have some minor sickness-related limitation of activity. Reported scores on the SIP for persons in a variety of clinical conditions are provided in Table 2.

The SIP can be administered either orally or as a written questionnaire. For this research, we administered the SIP orally in one-on-one sessions with interviewers and subjects. Thus, no literacy skills were needed. Also, because the SIP is available in English only, non-English-speaking persons were excluded (as previously noted).

PCAE staff members served as the interviewers who administered the SIP. Each was trained in SIP administration techniques.³⁹ Interviewers did not necessarily test students known to them but rather tested students from various classes who had been identified by the study's sampling technique. After testing, SIP results were scored independently by other members of the project staff.

Other Variables

Interviewers also collected the following sociodemographic information from the subjects:

age, marital status, ethnicity, place of birth, language(s) spoken in childhood home, educational attainment, income, health insurance status, and occupation. Occupations were categorized as either blue collar or white collar. We defined blue-collar occupations as those that involved manual labor, unskilled nonmanual labor (e.g., baby sitter, cashier), and nonsupervisory positions requiring technical skills (e.g., typing, driving).

Interviewers asked each subject for a self-assessment of health graded on a 4-point scale: excellent (1), good (2), fair (3), poor (4). In addition, interviewers collected information about the number of hospitalizations and physician office visits the subject had experienced during the past year and about the number of prescription medications taken on a daily basis.

Statistical Methods

The independent variable for this research was the grade-equivalent reading level. The dependent variables were the physical, psychosocial, and total SIP scores.

We examined the relation between reading level and SIP scores in two ways. First, we used standard linear models with reading level entered as a continuous characteristic. This analysis used both grade-equivalent reading level and log₁₀ (grade-equivalent reading level + 0.5).

For the second analysis, we dichotomized subjects into two groups: (1) those with reading levels at or below grade level 4, and (2) those with reading levels above grade level 4. We chose the 4th grade as the dividing point because this education level is used by the US Census Bureau to define literacy.¹³ We used a general linear model

Table 2. Range of Mean (SD) Sickness Impact Profile (SIP) Scores Reported in the Medical Literature.*

Setting or Illness	Number of Subjects	SIP Score		
		Physical Mean (SD)	Psychosocial Mean (SD)	Overall Mean (SD)
Healthy enrollees in prepaid health plan ²⁵	144			2.6 (4.5)
Myocardial infarct survivors at 6 mo ²⁷	308	6.9 (11.2)	8.8 (1.2)	10.3 (10.8)
Patients undergoing cancer radiotherapy ³¹	12			11.5
Rheumatoid arthritis ARA Class 2 ²⁶	64	13.1 (9.1)	11.7 (9.8)	15.1 (8.8)
Chronic back pain ³⁵	80	18.3 (15.0)	18.8 (17.0)	18.7 (14.3)
Chronic lung disease, oxygen dependent ³⁴	66	19.8		20.5
Cardiac arrest survivors with cognitive defects ²⁹	10			18.0 (17.0)

*Blank cells indicate that no data were reported.

to test for differences in SIP scores between the groups and adjusted for confounding socio-demographic covariables (using dummy variables for categorical variables).

Power analysis, utilizing reported distributions of SIP scores and the known distribution of reading levels among PCAE students, indicated that at least 175 subjects were needed for this research. The Statistical Analysis System (SAS) and the Statistical Package for the Social Sciences (SPSS) were used for data analysis.

Results

Of the first 197 randomly selected individuals who met eligibility requirements, 193 agreed to participate.

Indicators of Health Services Utilization and Self-Evaluation of Physical Health

One hundred fifty-five (80.3 percent) of the subjects reported visiting a physician's office for health care during the past year. The average number of times these 155 subjects had been to a physician's office was 5.53 ($\pm$ SD 11.92), range was 1 to 100 visits per year. The median number of visits was 3.0.

Thirty-two (16.6 percent) of the subjects had been hospitalized during the past year. The number of hospitalizations per subject among those 32 individuals ranged from 1 to 30; median was 1.0 and mean was 3.25 ($\pm$ SD 6.81).

Fifty-seven of the 193 subjects (29.5 percent) reported taking prescription medication on a regular basis. Among these 57 subjects, the average number of medications taken per day was 1.98 ($\pm$ SD 1.79).

Only 49 (25.4 percent) of the 193 subjects self-rated their health as excellent, whereas 92 (47.7 percent) rated their health as good, 45 (23.3 percent) rated it as fair, and 7 (3.6 percent) indicated poor health.

Reading Levels

The subjects' mean reading level was grade 7.17 ($\pm$ 2.77). The assessment of reading levels had been made either at entry into PCAE or at the beginning of the current academic session; we accepted the most recent of the two assessments. The mean time interval between the subject's reading level assessment and SIP testing




Figure 1. Frequency distribution of grade-equivalent reading levels among the 193 adults who participated in the study, with reading levels rounded to nearest integer grade level. All subjects were enrolled in a publicly funded literacy instruction program.

for this study was 2.2 ($\pm$ 3.6) months. Figure 1 shows the distribution of the subjects' reading levels.

SIP Scores

Table 3 presents unadjusted and adjusted mean SIP scores according to reading level (divided at the 4th grade level). Physical, psychosocial, and total SIP scores were all related to reading level. The strongest relation, however, was between reading level and SIP physical score.

Table 3. Mean Sickness Impact Profile (SIP) Scores According to Reading Level.

SIP Scores	Subjects Who Read at or below 4th Grade Level (n = 37)	Subjects Who Read above 4th Grade Level (n = 156)	P Value
Mean physical score			
Unadjusted*	6.5	2.5	0.0008
Adjusted†	6.2	2.3	0.002
Mean psychosocial score			
Unadjusted*	17.8	12.8	0.10
Adjusted†	15.4	8.0	0.02
Mean total (overall) score			
Unadjusted*	11.7	8.3	0.04
Adjusted†	10.4	6.0	0.02

*No adjustment for covariables.

†Adjusted for age, sex, ethnicity, marital status, insurance status, occupation, and income.




Figure 2. Frequency distribution of the Sickness Impact Profile (SIP) physical health scores among the 193 study participants. Higher scores represent a greater degree of sickness-related dysfunction.

SIP Physical Score

Figure 2 displays the distribution of SIP physical scores. The mean score was $3.25 (\pm 6.67)$. Subjects with very poor reading skills had higher mean physical SIP scores (greater degree of sickness-related dysfunction) than did subjects with more advanced literacy skills (Figure 3).

The mean physical SIP score for subjects with reading levels at or below grade level 4 was higher than the mean score of subjects who read at more advanced reading levels (6.54 versus 2.48, $P < 0.0008$). The relation remained significant ($P < 0.002$, Table 3) after adjusting for potentially confounding covariables. This statistical difference was primarily due to the effect of a small




Figure 3. Sickness Impact Profile (SIP) physical health scores of subjects at each grade-equivalent reading level. Higher SIP scores indicate a greater degree of sickness-related dysfunction. Reading levels were rounded to nearest integer grade level. Both mean SIP score and range of SIP scores given for each grade level.

number of subjects with very poor reading levels (grades 0.0 to 1.9).

SIP Psychosocial Scores

Psychosocial SIP scores ranged from 0.0 to 68.0; the mean was $13.75 (\pm 16.41)$. This mean score is comparable with scores found among patients with significant medically related psychosocial dysfunction (Table 2). The relation between psychosocial SIP score and reading level was significant after adjustment for confounding variables ($P < 0.02$, Table 3).

Discussion

We initially considered two possible study designs for this research. One design involved comparing the health status of illiterate adults with the health status of adults with "normal" literacy. Because illiteracy occurs frequently among those of low socioeconomic strata, the two groups would have very different sociodemographic characteristics, thus threatening validity of comparisons between the groups.

The second study design, which was ultimately used, involved evaluating persons whose demographic characteristics were relatively homogeneous and determining whether their variations in literacy skills were associated with variations in health status. Thus, the study population itself provided some inherent control over sociodemographic covariables. Statistical methods were then used to provide additional adjustment for these covariables.

Physical Health

The results of this research suggest that there is a relation between poor literacy skills and poor health in the United States, just as there is in nonindustrialized nations. The relation was statistically significant even after controlling for confounding covariables. The relation of literacy skills to physical health was particularly strong ($P < 0.002$) and was due primarily to the poor health status of individuals with extremely poor reading skills.

The results are of even greater interest given a research design that could have biased the results against finding a relation between illiteracy and poor health. One example of this bias is that the study population consisted mostly of young adults, a group with a limited range of medical

problems and less severe illness in comparison with older populations. The smaller range of medical problems decreased the range of SIP scores and, therefore, lessened the ability of statistical tests to detect a relation between SIP scores and illiteracy.

A second factor that could have biased the study against detecting a relation between health status and illiteracy is that the SIP test seemed to lack sensitivity for demonstrating health status differences among subjects: more than one-half of the subjects had a physical SIP score of 0.0. Here again, the decreased variation in SIP scores reduced the ability of statistical tests to detect a relation between illiteracy and poor health.

A third methodological bias was right truncation of the study population (i.e., exclusion of persons with reading levels above grade 12). If such individuals had been tested, differences in SIP scores between high- and low-level readers might have been more marked.

Psychosocial Health

The apparent poor psychosocial health of study participants is worthy of special note. Our subjects' psychosocial SIP scores were in the range generally found among persons with serious psychosocial dysfunction (Table 2).

Previous research has shown that, in general, persons from lower socioeconomic strata are at higher risk for depression and other psychiatric morbidity.⁴⁰⁻⁴⁶ The poor psychosocial health of subjects in this study thus could be partly attributable to their socioeconomic status. After adjusting for socioeconomic variables, however, psychosocial SIP scores were still related to literacy, suggesting that lack of adequate reading skills can itself play a role in poor psychosocial health.

The subjects in this study were a special group of self-selected adults who had the individual personal motivation to seek additional education. It is possible that psychosocial health impairment among other illiterate adults can be even more significant and contribute to their failure to seek remedial literacy education.

Implications for Clinicians

The mechanisms by which literacy and health are interrelated have not been clarified, nor has causation been established. That is, low literacy skills might or might not directly cause poor health

status. Nonetheless, the results of this study indicate that patients with extremely poor literacy skills (particularly at grade levels 0-1) are at increased risk for poor health, regardless of other sociodemographic characteristics.

There are several plausible mechanisms by which illiteracy could affect health status.⁴⁷ Illiterate patients might fail to obtain or understand information regarding their personal medical care. Illiterate individuals also might not understand how to use the health care system properly, potentially resulting in inappropriate overuse or underuse of services. Both mechanisms could increase costs and rates of adverse medical outcomes.

The association between illiteracy and health status found in this research suggests that improving literacy skills of US residents might improve the population's health status. If confirmed by other studies, there would be medical justification for initiatives to improve the literacy skills of our populace.

We thank the following members of the Pima County Adult Education program staff for their invaluable contributions to this research project: Peggy Altfater, Deborah Cassady, Kathleen Eichnorst, Pamela Hennessy, Maria Martinez, Cynthia Meir, Lois Miller, Emily Ravenscroft, Jackie O'Rourke, Ellen Shepherd, and Amy Stein.

References

1. UNESCO. Statistical yearbook 1988. Paris: United Nations Educational, Scientific and Cultural Organization, 1988:15-25.
2. Caldwell JC. Education as a factor in mortality decline: an examination of Nigerian data. *Popul Stud* 1979; 33:395-413.
3. Cochrane SH, O'Hara DJ, Leslie J. The effects of education on health: a background study for world development report. World Bank Staff Working Paper No. 405. Washington, DC: World Bank, 1980.
4. Foege WH, Henderson DA. Selective primary health care. XXV. Management priorities. *Rev Infect Dis* 1986; 8:467-75.
5. Flegg AT. Inequality of income, illiteracy and medical care as determinants of infant mortality in underdeveloped countries. *Popul Studies* 1982; 36:441-58.
6. Socioeconomic differentials in child mortality in developing countries. New York: United Nations Department of Social and Economic Affairs, 1985.
7. Bairagi R. Is income the only constraint on child nutrition in Bangladesh? *Bull World Health Organ* 1980; 58:767-72.
8. Choe MK, Retherford RD, Gubhaju BB, Thapa S. Ethnic differentials in early childhood mortality in Nepal. *J Biosoc Sci* 1989; 21:223-33.

9. Esrey SA, Habicht JP. Maternal literacy modifies the effect of toilets and piped water on infant survival in Maylasia. *Am J Epidemiol* 1988; 127: 1079-87.
10. Cleland JG, Van Ginneken JK. Maternal education and child survival in developing countries: the search for pathways of influence. *Soc Sci Med* 1988; 27:1357-68.
11. Grosse RN, Auffrey C. Literacy and health status in developing countries. *Ann Rev Public Health* 1989; 10:281-97.
12. Hunter CS. Adult illiteracy in the United States. New York: McGraw-Hill, 1985.
13. Kozol J. *Illiterate America*. Garden City, NY: Anchor Press/Doubleday, 1985.
14. Harman D. Turning illiteracy around: an agenda for national action: working paper no. 12. New York: Business Council for Effective Literacy, 1985.
15. The adult basic education program: progress in reducing illiteracy and improvements needed. Office of Education, Department of Health Education and Welfare. Report to Congress. Washington, DC: US General Accounting Office, 1975.
16. Kelso CR, Northcutt N, Selz N, Shelton E, Nyer L, Hickok D. Adult functional competency. A report to the Office of Education Dissemination Review Panel. Austin, TX: Division of Extension, University of Texas, 1975.
17. Berger J. Price of illiteracy translates into poverty and humiliation. *New York Times*. September 6, 1988; Sect. A1, Sect. B8.
18. Digest of education statistics 1988. Washington, DC: Government Printing Office, 1988:332.
19. Unlocking the future: adult literacy in Arizona: report of the Governors and Superintendent's Joint Task Force on Adult Illiteracy. Phoenix: Government Publication, 1986.
20. Literacy and the marketplace: improving the literacy skills of low-income single mothers: a report on a meeting of practitioners, policymakers, researchers, and funders. New York: Rockefeller Foundation, 1989.
21. Kirsch IS, Jungeblut A. Literacy: profiles of America's young adults. Report No. 16-P1-02. Princeton, NJ: Educational Testing Service, 1986.
22. Tests of adult basic education. Monterey, CA: CTB/McGraw-Hill, 1976.
23. Chapman BE, Copeman KL, Schulz L, Schulz C. The Mott basic language skills programs. Niles, MI: Allied Educational Press, 1981.
24. Carter WB, Bobbitt RA, Bergner M, Gilson BS. Validation of interval scaling: the sickness impact profile. *Health Serv Res* 1976; 11:516-28.
25. Gilson BS, Bergner M, Bobbitt RA, Carter WB. The sickness impact profile: final development and testing, 1975-1978. Final report to the National Center for Health Services Research. Seattle: Department of Health Services, School of Public Health and Community Medicine, University of Washington, 1979.
26. Deyo RA, Inui TS, Leininger JD, Overman SS. Measuring functional outcomes in chronic disease: a comparison of traditional scales and a self-administered health status questionnaire in patients with rheumatoid arthritis. *Med Care* 1983; 21: 180-92.
27. Bergner L, Hallstrom AP, Bergner M, Eisenberg MS, Cobb LA. Health status of survivors of out-of-hospital cardiac arrest and of myocardial infarction controls. *Am J Public Health* 1985; 75: 1321-3.
28. Ott CR, Sivarajan ES, Newton KM, Almes MJ, Bruce RA, Bergner M, et al. A controlled randomized study of early cardiac rehabilitation: the Sickness Impact Profile as assessment tool. *Heart Lung* 1983; 12:162-70.
29. Longstreth WT Jr, Inui TS, Cobb LA, Copass MK. Neurologic recovery after out-of-hospital cardiac arrest. *Ann Intern Med* 1983; 98:588-92.
30. Sugarbaker PH, Barofsky I, Rosenberg SA, Gianola FJ. Quality of life assessment of patients in extremity sarcoma clinical trials. *Surgery* 1982; 91:17-23.
31. Johnson JE, King KB, Murray RA. Measuring the impact of sickness on usual functions of radiation therapy patients. *Oncol Nurs Forum* 1983; 10:36-9.
32. Bergner M, Hudson LD, Conrad DA, Patmont CM, McDonald GJ, Perrin EB, et al. The cost and efficiency of home care for patients with chronic lung disease. *Med Care* 1988; 26:566-79.
33. McSweeney AJ, Grant I, Heaton RK, Adam KM, Timms RM. Life quality of patients with chronic obstructive pulmonary disease. *Arch Intern Med* 1982; 142:473-8.
34. Nocturnal Oxygen Therapy Trial Group. Continuous or nocturnal oxygen therapy in hypoxemic chronic obstructive lung disease: a clinical trial. *Ann Intern Med* 1980; 93:391-8.
35. Deyo RA, Diehl AK. Measuring physical and psychosocial function in patients with low back pain. *Spine* 1983; 8:635-42.
36. Deyo RA, Walsh NE, Martin DC, Schoenfeld LS, Ramamurthy S. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *N Engl J Med* 1990; 322:1627-34.
37. Rockey PH, Griep RJ. Behavioral dysfunction in hyperthyroidism. Improvement with treatment. *Arch Intern Med* 1980; 140:1194-7.
38. Bergner M, Bobbitt RA, Carter W, Gilson BS. The Sickness Impact Profile: development and final revision of a health status measure. *Med Care* 1981; 19:787-805.
39. Conn J, Bobbitt R, Bergner M. Administration procedures and interviewer training for the Sickness Impact Profile. Seattle: University of Washington Department of Health Services, 1979. Manual.
40. Huxley P, Raval H, Koror J, Jacob C. Psychiatric morbidity in the clients of social workers: clinical outcome. *Psychol Med* 1989; 19:189-97.
41. Kennedy GJ, Kelman HR, Thomas C, Wisniewski W, Metz H, Bijur PE. Hierarchy of characteristics

- associated with depressive symptoms in an urban elderly sample. *Am J Psychiatry* 1989; 146:220-5.
42. Maurer FA. Acute depression: treatment and nursing strategies for this affective disorder. *Nurs Clin North Am* 1986; 21:413-27.
 43. Heller TA, Larson EB, LoGerfo JP. Quality of ambulatory care of the elderly: an analysis of five conditions. *J Am Geriatr Soc* 1984; 32: 782-8.
 44. Williams DH. The epidemiology of mental illness in Afro-Americans. *Hosp Community Psychiatry* 1986; 37:42-9.
 45. Hall LA, Williams CA, Greenberg RS. Supports, stressors, and depressive symptoms in low-income mothers of young children. *Am J Public Health* 1985; 75:518-22.
 46. Whitley MP, Osborne OH, Godfrey MA, Johnston K. A point prevalence study of alcoholism and mental illness among downtown migrants. *Soc Sci Med* 1985; 20:579-83.
 47. Weiss BD, Hart G, Pust R. Relationship between illiteracy and health. *J Health Care Poor Under-served* 1991; 1:43-55.

ABFP ANNOUNCEMENT

Geriatrics Certificate of Added Qualification (CAQ)

The last opportunity to qualify for the American Board of Family Practice (ABFP) Certificate of Added Qualification in Geriatric Medicine via a nonfellowship pathway will be April 1994.

Applications will be available beginning July 1, 1993. All applications must be returned to the Board office by November 1, 1993.

RESERVE YOUR APPLICATION TODAY

Send a written request on letterhead stationery and your application will be automatically sent to you in July 1993.

Send your written request for application materials to:

Geriatrics CAQ
American Board of Family Practice
2228 Young Drive
Lexington, Kentucky 40505