

Dermatologic Manifestations Of Giardiasis

Jerry T. McKnight, M.D., and Paul E. Tietze, M.D.

Giardia lamblia is the most common intestinal parasite in the United States and is worldwide in distribution.¹ Approximately 4 percent of stool specimens submitted to public health laboratories in this country contain *Giardia* cysts.² The usual symptoms of acute giardiasis include diarrhea, abdominal cramps, nausea, and weight loss. Many, if not most, individuals with *Giardia* infection are asymptomatic. Giardiasis can be an acute self-limiting diarrheal illness, or it can lead to chronic diarrhea and malabsorption.³ Children are affected more often than adults, and person-to-person transmission has been documented.^{4,5} There have been reports of allergic symptoms associated with giardiasis. This article describes a case of dermatitis associated with *Giardia* infection and reviews dermatologic manifestations of *Giardia lamblia* infection.

Case Report

A 48-year-old woman complained of an 8-month history of atopic dermatitis. The patient had been cared for by a dermatologist, who prescribed triamcinolone ointment 0.1 percent topically twice daily, hydroxyzine hydrochloride 25 mg four times daily, and doxepin hydrochloride 50 mg at bedtime. This treatment provided some symptomatic relief but had not cleared the markedly pruritic chronic dermatitis. There was no history of atopy, nor was there any history of dermatologic conditions. The patient requested her medications be changed because of worsening of the rash.

The patient's medical history was unremarkable except one previous hospitalization for an uncomplicated childbirth. The social history was pertinent for a move to east Tennessee 10 months earlier from her native country of Germany to live with her daughter. The dermatitis developed approximately 2 months after moving.

The physical examination was remarkable only for an eczematous-type dermatitis with erythema, xerosis, and several fine papular and vesicular lesions on the extremities. The rash was confined primarily to the trunk and extremities and was located predominately on the flexor surfaces. This pattern was consistent with the previous diagnosis of atopic dermatitis. The involved skin was excoriated, and lichenification was present.

The patient's complete blood count, chemical analysis of serum, urinalysis, and thyroid profile were normal. The serum immunoglobulin E (IgE) was 48 $\mu\text{g/L}$ (20 U/mL), normal 24–240 $\mu\text{g/L}$ (10–100 U/mL).

The medications were changed to fluocinonide ointment 0.05 percent twice daily and terfenadine 60 mg twice daily. This medication regimen decreased symptoms somewhat; however, the dermatitis persisted with the pruritus being especially bothersome.

Several weeks after changing medications, the patient developed the new complaint of abdominal cramping without nausea, diarrhea, or weight loss. Further questioning revealed that cramping had been an intermittent problem starting soon after coming to live with her daughter. A stool culture was negative for enteric pathogens; however, a microscopic examination showed numerous *Giardia* cysts. Treatment with metronidazole, 250 mg three times a day for 7 days, relieved the abdominal complaints. Additionally, the dermatitis completely resolved within 2 weeks. All medications were discontinued, and the rash had not recurred after 18 months of follow-up care.

The suspected source of infection was an underground well, which was the only water source for the family. The local health department believed it was impractical to filter mass quantities of water for detection of *Giardia*. The other family members were asked to have stool specimens for parasites. No members of the household were symptomatic, and they declined. The family was asked to boil water used for consumption.

Submitted, revised, 2 October 1991.

From the Department of Family Medicine, University of Alabama, Tuscaloosa. Address reprint requests to Jerry T. McKnight, M.D., Department of Family Medicine, University of Alabama, 107 Educational Tower, Tuscaloosa, AL 35401.

Discussion

This case illustrates a possible complication of giardiasis. An extensive review of the medical literature, including foreign sources, revealed several references to dermatologic manifestations, specifically urticaria, associated with giardiasis.⁶⁻⁸ Infection with *Giardia* has been associated with urticaria, pruritus, and possibly angioedema.^{9,10} Other allergic manifestations, such as reactive arthritis, have also been reported.¹¹ *Giardia* is not usually associated with elevated IgE levels or eosinophilia.^{5,12} Appropriate treatment of this parasite resulted in complete resolution of skin manifestations in all cases described. The cases described in the literature are presented in Table 1.

Additional studies not tabulated include a study in 1977 of 66 Argentinian children, aged 2 to 15 years, who came to a rheumatology clinic with joint symptoms. Goobar¹¹ reported that these children had joint symptoms of pain accompanied by synovitis. Sixty-four of these children had giardiasis. The 66 patients were selected because of gastrointestinal symptoms, including diarrhea, nausea or vomiting, abdominal cramps, or bloody stools. Sixty percent had what were described as allergic symptoms: nasal itching, generalized pruritus, and anal itching. Urticaria or dermatitis was not specifically reported. The report indicated that after treatment with either metronidazole or quinacrine, 90 percent of the articular and extra-articular signs and symptoms disappeared.¹¹

Another specific report of note is that by Chirila, et al.,¹⁰ who in 1981 studied 434 patients who came to the Department of Allergy at the Institute of Internal Medicine in Bucharest. Of these patients, 186 had chronic urticaria, 120 had angioedema, and 50 had both urticaria and angioedema. *Giardia* was found in 74 (40 percent) of patients with chronic urticaria, in 42 (35 percent) of patients with angioedema, and in 48 (96 percent) of patients who had both chronic urticaria and angioedema. The authors indicated clinical symptoms were rarely suggestive of giardiasis. *Giardia* was found in these allergic patients three times more frequently than the normal population of Rumania. The authors did not report dermatologic response to treatment in these patients.

There have been many cases of urticaria associated with *Giardia lamblia* infection. Previous approximations of nine total cases by Hamrick and Moore¹⁹ in 1983 and 20 cases by Clyne and Eliopoulos²⁰ in 1989 were underestimations. The total number of cases reported is 33, not counting

Table 1. Literature Review of Giardiasis Associated with Urticaria or Pruritus.

Year	Cases	Ages (yr)	Other Symptoms	Treatment	Outcome
1949	1	16	Diarrhea, abdominal pain	Quinacrine	All symptoms resolved ⁶
1957	2	34, 42	Diarrhea, weight loss	Quinacrine	All symptoms resolved ⁷
1958	6*	NR	Gastrointestinal symptoms	Quinacrine	Urticaria resolved ⁸
1969	5†	NR	NR	NR	NR ¹³
1976	1	28	Diarrhea, weight loss	Metronidazole	All symptoms resolved ¹⁴
1978	4	NR	Diarrhea	Metronidazole	All symptoms resolved ¹⁵
1979	1	30s	None	Metronidazole	Urticaria resolved ¹⁶
1980	3	1, 13, 35	Anorexia, diarrhea, weight loss	Metronidazole	All symptoms resolved ¹⁷
1983	6‡	5-33	50% had gastrointestinal symptoms	Metronidazole	All symptoms resolved ¹⁸
1983	1	4	None	Metronidazole	All symptoms resolved ¹⁹
1983	1	6	Diarrhea, arthritis	Metronidazole	All symptoms resolved ⁹
1989	1	28	Diarrhea, fever	Metronidazole	All symptoms resolved ²⁰
1989	1	53	Diarrhea, fever	Metronidazole	Urticaria resolved ²¹
1990	1§	NR	Mild gastrointestinal symptoms	Metronidazole	All symptoms resolved ²²

NR = not reported.

*6 of 32 patients with giardiasis.

†5 of 500 patients with giardiasis.

‡6 of 50 patients with chronic urticaria.

§Self-report, allergist, pruritus only.

additional cases unreported by Kennou, et al.¹⁷ and the Rumanian study,¹⁰ which did not report response to treatment. Goobar¹¹ also did not quantitate the exact number with pruritus or the specific response to treatment in each incident.

Some of these patients had few if any gastrointestinal symptoms. This finding is not surprising in view of the known history of giardiasis.³ The types of gastrointestinal presentations of giardiasis range from minimal or no symptoms to diarrhea and weight loss.³ Fever is also occasionally seen.^{20,23,24} The number of reports and the differences in clinical presentations, coupled with the geographic diversity, make clear that this problem is not unusual worldwide. The rather consistent response to treatment leaves little doubt there is an association between *Giardia* and urticaria, pruritus, and possibly other allergic symptoms, such as other skin rashes and synovitis.

Our patient had what appeared to be atopic dermatitis, but it occurred in an adult who had no personal or family history of atopy and who had a normal IgE level. Atopic dermatitis in association with giardiasis has not previously been described. It is, however, not unreasonable to postulate a similar pathogenesis, as this condition is clearly allergy related.

The appearance of the dermatitis after exposure to untreated well water was significant. The dermatologic response to treatment for giardiasis was dramatic and, although anecdotal, is highly suggestive of a relation between the two. It is possible this rash was secondary to another cause; however, complete resolution of this dermatitis after a single course of metronidazole and the lack of recurrence make other causes unlikely.

The immune response to *Giardia lamblia* is not completely understood. The immune system generates both a humoral and cellular response to *Giardia*.^{25,26} Serum antibodies to *Giardia* of IgG and IgM classes have been found.²⁷⁻²⁹ Secretory IgA antibodies to *Giardia* antigens have been found in breast milk.³⁰ Giardiasis has been associated with reduced intestinal secretory IgA, and patients with immunodeficiency syndromes have more severe gastrointestinal syndromes.^{31,32}

The mechanism of urticaria associated with giardiasis has not been elucidated. Although urticaria can be caused by immunologic and nonimmunologic mechanisms, it is most likely

the association with *Giardia* is immunologically based. This reaction, however, might not be a classical IgE-mediated phenomenon.³³ Although urticaria is not a usual manifestation of *Giardia*, it should be recognized, especially in view of the ubiquitous nature of the organism. Because chronic urticaria is a relatively common dermatologic disorder and results in a frequently frustrating and unfruitful evaluation, it would be reasonable to include *Giardia* in the differential diagnosis and to search for *Giardia* by either microscopic stool analysis or by the newer and perhaps more sensitive stool antigen detection method.

Detection of *Giardia lamblia* can be difficult. Microscopic identification is dependent on the skill of the examiner and appropriate sample collection and preparation. The sensitivity of microscopic stool analysis varies but is approximately 50 to 60 percent.^{34,35} The sensitivity of stool antigen detection methods has been reported as 88 to 96 percent and the specificity as 95 to 100 percent.³⁶⁻³⁸

Summary

We have presented a case of atopic dermatitis associated with *Giardia lamblia* infection, which has not been previously described. Review of the world literature shows an association between giardiasis and urticaria. Other allergic phenomena, such as angioedema and possibly arthropathy, also might be associated with this infection. When confronted with these clinical problems, giardiasis should be included in the differential diagnosis. As these complications respond to specific therapy, identification of this organism as its cause can be particularly rewarding.

References

1. Craft JC. *Giardia* and giardiasis in childhood. *Pediatr Infect Dis* 1982; 1:196-211.
2. Center for Disease Control. Intestinal parasite surveillance—United States, 1976. *MMWR* 1978; 27(20):167-8.
3. Dupont HL, Sullivan PS. Giardiasis: the clinical spectrum, diagnosis and therapy. *Pediatr Infect Dis* 1986; 5(Suppl):S131-8.
4. Black RE, Dykes AC, Sinclair SP, Wells JG. Giardiasis in day-care centers: evidence of person-to-person transmission. *Pediatrics* 1977; 60:486-91.
5. Burke JA. Giardiasis in childhood. *Am J Dis Child* 1975; 129:1304-10.

6. Harris RH, Mitchell JH. Chronic urticaria due to *Giardia lamblia*. Arch Dermatol Syphilology 1949; 59:587-9.
7. Wilhelm RE. Urticaria associated with *Giardia lamblia*. J Allergy 1957; 28:351-3.
8. Webster BH. Human infection with *Giardia lamblia*: an analysis of thirty-two cases. Am J Dig Dis 1958; 3:64-71.
9. Farthing MJ, Chong SK, Walker-Smith JA. Acute allergic phenomena in giardiasis [letter]. Lancet 1983; 2:1428.
10. Chirila M, Panaitescu D, Capraru T. Frequency of *Giardia lamblia* in certain allergic syndromes. Med Interne 1981; 19:367-72.
11. Goobar JP. Joint symptoms in giardiasis [letter]. Lancet 1977; 1:1010-1.
12. Geller M, Geller M, Flaherty DK, Black P, Madruga M. Serum IgE levels in giardiasis. Clin Allergy 1978; 8:69-71.
13. Felix H, Freyria J, Maillard A, Legendre M, Douchet C, Le Vagueresse R. [Apropos of 500 cases of lambliasis in adults]. J Med Lyon 1969; 50:163-73.
14. Dellamonica P, Le Fichoux Y, Monnier B, Duplay H. [Dysenteric syndrome and urticaria in giardiasis (letter)]. Nouv Presse Med 1976; 5:1913.
15. La Rosa F, Caruso R, Bertuccio O. [Urticarial manifestations during lambliasis]. Policlinico Sez Med 1978; 85:77-81.
16. Weisman BL. Urticaria and *Giardia lamblia* infection. Ann Allergy 1979; 42(2):91.
17. Kennou MF, Jenayah S, Rekhis M, Bayar N, Necibi S. [Skin manifestations of giardiasis. Some clinical cases]. Arch Inst Pasteur Tunis 1980; 57:257-60.
18. Veronesi S, Palmerio B, Negosanti M, Tosti A. Urticaria and giardiasis. Dermatologica 1983; 166:42-3.
19. Hamrick HJ, Moore GW. Giardiasis causing urticaria in a child. Am J Dis Child 1983; 137:761-3.
20. Clyne CA, Eliopoulos GM. Fever and urticaria in acute giardiasis. Arch Intern Med 1989; 149:939-40.
21. Juergens UR, Predel HG, Vetter H. [Urticaria, pruritus (clinical conference)]. Schweiz Rundsch Med Prax 1989; 78:403-5.
22. Spaulding HS Jr. Pruritus without urticaria in acute giardiasis [letter]. Ann Allergy 1990; 65:161.
23. Pickering LK, Engelkirk PG. *Giardia lamblia*. Pediatr Clin North Am 1988; 35:565-77.
24. Meyer EA, Jarroll EL. Giardiasis. Am J Epidemiol 1980; 111:1-12.
25. Smith PD, Elson CO, Keister DB, Nash TE. Human host response to *Giardia lamblia*. I. Spontaneous killing by mononuclear leukocytes in vitro. J Immunol 1982; 128:1372-6.
26. Visvesvara GS, Smith PD, Healy GR, Brown WR. An immunofluorescence test to detect serum antibodies to *Giardia lamblia*. Ann Intern Med 1980; 93:802-5.
27. Ridley MJ, Ridley DS. Serum antibodies and jejunal histology in giardiasis associated with malabsorption. J Clin Path 1976; 29:30-4.
28. Smith PD, Gillin FD, Brown WR, Nash TE. IgG antibody to *Giardia lamblia* detected by enzyme-linked immunosorbent assay. Gastroenterology 1981; 80:1476-80.
29. Goka AK, Rolston DD, Mathan VI, Farthing MJ. Diagnosis of giardiasis by specific IgM antibody enzyme-linked immunosorbent assay. Lancet 1986; 2:184-6.
30. Miotti PG, Gilman RH, Pickering LK, Ruiz-Palacios G, Park HS, Yolken RH. Prevalence of serum and milk antibodies to *Giardia lamblia* in different populations of lactating women. J Infect Dis 1985; 152:1025-31.
31. Ament ME. Immunodeficiency syndromes and gastrointestinal disease. Pediatr Clin North Am 1975; 22:807-25.
32. Zinneman HH, Kaplan AP. The association of giardiasis with reduced intestinal secretory immunoglobulin A. Am J Dig Dis 1972; 17:793-7.
33. Monroe EW, Jones HE. Urticaria. An updated review. Arch Dermatol 1977; 113:80-90.
34. Kamath KR, Murugasu R. A comparative study of four methods for detecting *Giardia lamblia* in children with diarrheal disease and malabsorption. Gastroenterology 1974; 66:16-21.
35. Danciger M, Lopez M. Numbers of *Giardia* in the feces of infected children. Am J Trop Med Hyg 1975; 24:237-42.
36. Janoff EN, Craft JC, Pickering LK, Novotny T, Blaser MJ, Knisley CV, et al. Diagnosis of *Giardia lamblia* infections by detection of parasite-specific antigens. J Clin Microbiol 1989; 27:431-5.
37. Rosoff JD, Sanders CA, Sonnad SS, DeLay PR, Hadley WK, Vincenzi FF, et al. Stool diagnosis of giardiasis using a commercially available enzyme immunoassay to detect *Giardia*-specific antigen 65 (GSA 65). J Clin Microbiol 1989; 27:1997-2002.
38. Knisley CV, Engelkirk PG, Pickering LK, West MS, Janoff EN. Rapid detection of *Giardia* antigen in the stool with the use of enzyme immunoassays. Am J Clin Path 1989; 91:704-8.