

For first-line therapy in mild-to-moderate hypertension

**Discover the classic benefits of a beta-blocker  
and a diuretic...now at low doses for a  
side-effect profile comparable to placebo<sup>1\*</sup>**



**ZIAC controls mild-to-moderate hypertension  
in up to 80% of patients<sup>1†</sup>**

**ZIAC controls blood pressure for a full 24 hours  
for true once-a-day dosing<sup>2</sup>**

**ZIAC minimizes traditional beta-blocker- and  
HCTZ-associated metabolic effects (hypokalemia,  
hyperuricemia, hypercholesterolemia, hyperglycemia)<sup>1</sup>**

\*The two most common side effects — dizziness and fatigue — occurred at rates comparable to placebo.

<sup>†</sup>Clinical trial response rates were: 2.5 mg—61%; 5 mg—73%; 10 mg—80%.

ZIAC is contraindicated in patients in cardiogenic shock, overt cardiac failure (see WARNINGS section of full Prescribing Information), second- or third-degree AV block, marked sinus bradycardia, anuria, and hypersensitivity to either component of this product or to other sulfonamide-derived drugs.

Please see Brief Summary of Prescribing Information on adjacent page.

First-line therapy option

**ZIAC™**

(bisoprolol fumarate-hydrochlorothiazide)  
2.5, 5, & 10 mg Tablets with 6.25 mg HCTZ

First-line therapy option  
**ZIAC™**  
 (bisoprolol fumarate-hydrochlorothiazide)  
 2.5, 5, & 10 mg Tablets with 6.25 mg HCTZ

- References:**  
 1. DeQuattro V, Weir MR. Bisoprolol fumarate/hydrochlorothiazide 6.25 mg: a new, low-dose option for first-line antihypertensive therapy. *Adv Ther.* 1993;10:197-206.  
 2. Lewin AJ, Lueg MC, Targum S, et al. A clinical trial evaluating the 24-hour effects of bisoprolol/hydrochlorothiazide 5 mg/6.25 mg combination in patients with mild to moderate hypertension. *Clin Cardiol.* 1993;16:732-736.

**Brief Summary**

**ZIAC™ (Bisoprolol Fumarate and Hydrochlorothiazide) Tablets**

FOR FULL PRESCRIBING INFORMATION, PLEASE CONSULT PACKAGE INSERT.

**DESCRIPTION**

ZIAC (bisoprolol fumarate and hydrochlorothiazide) is indicated for the treatment of hypertension. It combines two antihypertensive agents in a once-daily dosage: a synthetic beta<sub>1</sub>-selective (cardioselective) adrenoceptor blocking agent (bisoprolol fumarate) and a benzothiadiazine diuretic (hydrochlorothiazide).

**CLINICAL PHARMACOLOGY**

At doses  $\geq$  20 mg bisoprolol fumarate inhibits beta<sub>1</sub>-adrenoceptors located in bronchial and vascular musculature. To retain relative selectivity, it is important to use the lowest effective dose.

**CONTRAINDICATIONS**

Cardiogenic shock, overt cardiac failure (see WARNINGS), second or third degree AV block, marked sinus bradycardia, anuria, and hypersensitivity to either component of this product or to other sulfonamide-derived drugs.

**WARNINGS**

**Cardiac Failure:** Beta-blocking agents should be avoided in patients with overt congestive failure.

**Patients Without a History of Cardiac Failure:** Continued depression of the myocardium with beta-blockers can precipitate cardiac failure. At the first signs or symptoms of heart failure, discontinuation of ZIAC should be considered.

**Abrupt Cessation of Therapy:** Abrupt cessation of beta-blockers should be avoided. Even in patients without overt coronary artery disease, it may be advisable to taper therapy with ZIAC over approximately 1 week with the patient under careful observation. If withdrawal symptoms occur, beta-blocking agent therapy should be reinstated, at least temporarily.

**Peripheral Vascular Disease:** Beta-blockers should be used with caution in patients with peripheral vascular disease.

**Bronchospastic Disease:** PATIENTS WITH BRONCHOSPASTIC PULMONARY DISEASE SHOULD, IN GENERAL, NOT RECEIVE BETA-BLOCKERS.

**Anesthesia and Major Surgery:** If used perioperatively, particular care should be taken when anesthetic agents that depress myocardial function, such as ether, cyclopropane, and trichloroethylene, are used.

**Diabetes and Hypoglycemia:** Beta-blockers may mask some of the manifestations of hypoglycemia, particularly tachycardia. Patients subject to spontaneous hypoglycemia, or diabetic patients receiving insulin or oral hypoglycemic agents, should be cautioned. Also, latent diabetes mellitus may become manifest and diabetic patients given thiazides may require adjustment of their insulin dose.

**Thyrotoxicosis:** Beta-adrenergic blockade may mask clinical signs of hyperthyroidism. Abrupt withdrawal of beta-blockade may be followed by an exacerbation of the symptoms of hyperthyroidism or may precipitate thyroid storm.

**Renal Disease:** Cumulative effects of the thiazides may develop in patients with impaired renal function. In such patients, thiazides may precipitate azotemia. In subjects with creatinine clearance less than 40 mL/min, the plasma half-life of bisoprolol fumarate is increased up to threefold, as compared to healthy subjects.

**Hepatic Disease:** ZIAC should be used with caution in patients with impaired hepatic function or progressive liver disease.

**PRECAUTIONS**

**General: Electrolyte and Fluid Balance Status:** Periodic determination of serum electrolytes should be performed, and patients should be observed for signs of fluid or electrolyte disturbances. Thiazides have been shown to increase the urinary excretion of magnesium; this may result in hypomagnesemia. Hypokalemia may develop. Hypokalemia and hypomagnesemia can provoke ventricular arrhythmias or sensitize or exaggerate the response of the heart to the toxic effects of digitalis. Dilutional hyponatremia may occur in edematous patients in hot weather; appropriate therapy is water restriction rather than salt administration, except in rare instances when the hyponatremia is life-threatening. In actual salt depletion, appropriate replacement is the therapy of choice.

**Parathyroid Disease:** Calcium excretion is decreased by thiazides, and pathologic changes in the parathyroid glands, with hypercalcemia and hypophosphatemia, have been observed in a few patients on prolonged thiazide therapy. **Hyperuricemia:** Hyperuricemia or acute gout may be precipitated in certain patients receiving thiazide diuretics. Bisoprolol fumarate, alone or in combination with HCTZ, has been associated with increases in uric acid.

**Drug Interactions:** ZIAC may potentiate the action of other antihypertensive agents used concomitantly. ZIAC should not be combined with other beta-blocking agents. In patients receiving concurrent therapy with clonidine, if therapy is to be discontinued, it is suggested that ZIAC be discontinued for several days before the withdrawal of clonidine.

ZIAC should be used with caution when myocardial depressants or inhibitors of AV conduction or anti-arrhythmic agents are used concurrently.

**Bisoprolol Fumarate:** Concurrent use of ritampin increases the metabolic clearance of bisoprolol fumarate, shortening its elimination half-life. Pharmacokinetic studies document no clinically relevant interactions with other agents given concomitantly, including thiazide diuretics, digoxin and cimetidine. There was no effect of bisoprolol fumarate on prothrombin times in patients on stable doses of warfarin.

While taking beta-blockers, patients with a history of severe anaphylactic reaction may be more reactive to repeated challenge, either accidental, diagnostic, or therapeutic and may be unresponsive to the usual doses of epinephrine used to treat allergic reactions.

**Hydrochlorothiazide:** The following drugs may interact with thiazide diuretics. Alcohol, barbiturates, or narcotics—potentiation of orthostatic hypotension may occur. Dosage adjustment of the antidiabetic drugs (oral agents and insulin) may be required. Other antihypertensive drugs—additive effect or potentiation. Cholestyramine and colestipol resins—single doses of cholestyramine and colestipol resins bind the hydrochlorothiazide and reduce its absorption in the gastrointestinal tract by up to 85 and 43 percent, respectively. Corticosteroids, ACTH—intensity of electrolyte depletion, particularly hypokalemia. Possible decreased response to pressor amines but not sufficient to preclude their use. Possible increased responsiveness to muscle relaxants, nondepolarizing. Generally, lithium should not be given with diuretics. Diuretic agents reduce the renal clearance of lithium and add a high risk of lithium toxicity. The administration of a nonsteroidal anti-inflammatory agent can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing and thiazide diuretics.

In patients receiving thiazides, sensitivity reactions may occur with or without a history of allergy or bronchial asthma. Photosensitivity reactions and possible exacerbation or activation of systemic lupus erythematosus have been reported in patients receiving thiazides. The antihypertensive effects of thiazides may be enhanced in the post-sympathectomy patient.

**Laboratory Test Interactions:** Based on reports involving thiazides, ZIAC may decrease serum levels of protein-bound iodine without signs of thyroid dysfunction. Because it includes a thiazide, ZIAC should be discontinued before carrying out tests for parathyroid function (see PRECAUTIONS—Parathyroid Disease).

**ADVERSE REACTIONS**

**ZIAC:** Bisoprolol fumarate/H6.25 mg is well tolerated in most patients. Most adverse effects (AEs) have been mild and transient. In more than 85,000 patients treated worldwide with bisoprolol fumarate, occurrences of bronchospasm have been rare. Discontinuation rates for AEs were similar for B/H6.25 mg and placebo-treated patients.

In the United States, 252 patients received bisoprolol fumarate (2.5, 5, 10, or 40 mg)/H6.25 mg and 144 patients received placebo in two controlled trials. In Study 1, bisoprolol fumarate 5/H6.25 mg was administered for 4 weeks. In Study 2, bisoprolol fumarate 2.5, 10 or 40/H6.25 mg was administered for 12 weeks. All adverse experiences, whether drug-related or not, and drug-related adverse experiences in patients treated with B2.5-10/H6.25 mg, reported during comparable, 4 week treatment periods by at least 2% of bisoprolol fumarate/H6.25 mg-treated patients (plus additional selected adverse experiences) are presented in the following table.

**ZIAC™ (Bisoprolol Fumarate and Hydrochlorothiazide) Tablets**

Body System/ Adverse Experience	% of Patients with Adverse Experiences*			
	All Adverse Experiences		Drug-related Adverse Experiences	
	Placebo† (n=144) %	B2.5-40/H6.25† (n=252) %	Placebo† (n=144) %	B2.5-10/H6.25† (n=221) %
<b>Cardiovascular</b>				
bradycardia	0.7	1.1	0.7	0.9
arrhythmia	1.4	0.4	0.0	0.0
peripheral ischemia	0.9	0.7	0.9	0.4
chest pain	0.7	1.8	0.7	0.9
<b>Respiratory</b>				
bronchospasm	0.0	0.0	0.0	0.0
cough	1.0	2.2	0.7	1.5
rhinitis	2.0	0.7	0.7	0.9
URI	2.3	2.1	0.0	0.0
<b>Body as a Whole</b>				
asthenia	0.0	0.0	0.0	0.0
fatigue	2.7	4.6	1.7	3.0
peripheral edema	0.7	1.1	0.7	0.9
<b>Central Nervous System</b>				
dizziness	1.8	5.1	1.8	3.2
headache	4.7	4.5	2.7	0.4
<b>Musculoskeletal</b>				
muscle cramps	0.7	1.2	0.7	1.1
myalgia	1.4	2.4	0.0	0.0
<b>Psychiatric</b>				
insomnia	2.4	1.1	2.0	1.2
somnolence	0.7	1.1	0.7	0.9
loss of libido	1.2	0.4	1.2	0.4
impotence	0.7	1.1	0.7	1.1
<b>Gastrointestinal</b>				
diarrhea	1.4	4.3	1.2	1.1
nausea	0.9	1.1	0.9	0.9
dyspepsia	0.7	1.2	0.7	0.9

\*Averages adjusted to combine across studies.

†Combined across studies.

Other adverse experiences that have been reported with the individual components are listed below.

**Bisoprolol Fumarate:** In clinical trials worldwide, a variety of other AEs, in addition to those listed above, have been reported. While in many cases it is not known whether a causal relationship exists between bisoprolol and these AEs, they are listed to alert the physician to a possible relationship. **Central Nervous System:** Unsteadiness, vertigo, syncope, paresthesia, hyperesthesia, sleep disturbance/vivid dreams, depression, anxiety/restlessness, decreased concentration/memory. **Cardiovascular:** Palpitations and other rhythm disturbances, cold extremities, claudication, hypotension, orthostatic hypotension, chest pain, congestive heart failure. **Gastrointestinal:** Gastric/epigastric/abdominal pain, peptic ulcer, gastritis, vomiting, constipation, dry mouth. **Musculoskeletal:** Arthralgia, muscle/joint pain, back/neck pain, twitching/tremor. **Skin:** Rash, acne, eczema, psoriasis, skin irritation, pruritus, purpura, flushing, sweating, alopecia, dermatitis, exfoliative dermatitis (very rarely). **Special Senses:** Visual disturbances, ocular pain/pressure, abnormal lacrimation, tinnitus, decreased hearing, earache, taste abnormalities. **Metabolic:** Gout. **Respiratory:** Asthma, bronchitis, dyspnea, pharyngitis, sinusitis. **Genitourinary:** Peyronie's disease (very rarely), cystitis, renal colic, polyuria. **General:** Malaise, edema, weight gain, angioedema.

In addition, a variety of adverse effects have been reported with other beta-adrenergic blocking agents and should be considered potential adverse effects: **Central Nervous System:** Reversible mental depression progressing to catatonia, hallucinations, an acute reversible syndrome characterized by disorientation to time and place, emotional lability, slightly clouded sensorium. **Allergic:** Fever, combined with aching and sore throat, laryngospasm, and respiratory distress. **Hematologic:** Agranulocytosis, thrombocytopenia. **Gastrointestinal:** Mesenteric arterial thrombosis and ischemic colitis. **Miscellaneous:** The oculomucocutaneous syndrome associated with the beta-blocker practolol has not been reported with bisoprolol fumarate during investigational use or extensive foreign marketing experience.

**Hydrochlorothiazide:** The following adverse experiences, in addition to those listed in the above table, have been reported with hydrochlorothiazide (generally with doses of 25 mg or greater). **General:** Weakness. **Central Nervous System:** Vertigo, paresthesia, restlessness. **Cardiovascular:** Orthostatic hypotension (may be potentiated by alcohol, barbiturates, or narcotics). **Gastrointestinal:** Anorexia, gastric irritation, cramping, constipation, jaundice (intrahepatic cholestatic jaundice), pancreatitis, cholecystitis, sialadenitis, dry mouth. **Musculoskeletal:** Muscle spasm. **Hypersensitive Reactions:** Purpura, photosensitivity, rash, urticaria, necrotizing angitis (vasculitis and cutaneous vasculitis), fever, respiratory distress including pneumonitis and pulmonary edema, anaphylactic reactions. **Special Senses:** Transient blurred vision, xanthopsia. **Metabolic:** Gout. **Genitourinary:** Sexual dysfunction, renal failure, renal dysfunction, interstitial nephritis.

**LABORATORY ABNORMALITIES**

**ZIAC:** Because of the low dose of hydrochlorothiazide in ZIAC, adverse metabolic effects with B/H6.25 mg are less frequent and of smaller magnitude than with HCTZ 25 mg.

Treatment with both beta-blockers and thiazide diuretics is associated with increases in uric acid. Mean increases in serum triglycerides were observed in patients treated with bisoprolol fumarate and hydrochlorothiazide 6.25 mg. Total cholesterol was generally unaffected, but small decreases in HDL cholesterol were noted.

Other laboratory abnormalities that have been reported with the individual components are listed below.

**Bisoprolol Fumarate:** In clinical trials, the most frequently reported laboratory change was an increase in serum triglycerides, but this was not a consistent finding.

Sporadic liver test abnormalities have been reported. In the U.S. controlled trials experience with bisoprolol fumarate treatment for 4 to 12 weeks, the incidence of concomitant elevations in SGOT and SGPT of between 1 to 2 times normal was 3.9%, compared to 2.5% for placebo. No patient had concomitant elevations greater than twice normal.

In the long-term, uncontrolled experience with bisoprolol fumarate treatment for 6-18 months, the incidence of one or more concomitant elevations in SGOT and SGPT of between 1-2 times normal was 6.2%. The incidence of multiple occurrence was 1.9%. For concomitant elevations in SGOT and SGPT of greater than twice normal, the incidence was 1.5%. The incidence of multiple occurrences was 0.3%. In many cases these elevations were attributed to underlying disorders, or resolved during continued treatment with bisoprolol fumarate.

Other laboratory changes included small increases in uric acid, creatinine, BUN, serum potassium, glucose, and phosphorus and decreases in WBC and platelets. There have been occasional reports of eosinophilia. These were generally not of clinical importance and rarely resulted in discontinuation of bisoprolol fumarate.

As with other beta-blockers, ANA conversions have also been reported on bisoprolol fumarate. About 15% of patients in long-term studies converted to a positive titer, although about one-third of these patients subsequently converted to a negative titer while on continued therapy.

**Hydrochlorothiazide:** Hyperglycemia, glycosuria, hyperuricemia, hypokalemia and other electrolyte imbalances (see PRECAUTIONS), hyperlipidemia, hypercalcemia, leukopenia, agranulocytosis, thrombocytopenia, aplastic anemia, and hemolytic anemia have been associated with HCTZ therapy.

See **DOSE AND ADMINISTRATION** section in package insert for complete dosing and precautionary information.

 ADVANTUS PHARMACEUTICALS and LEDERLE LABORATORIES DIVISION  
 American Cyanamid Company  
 Pearl River, NY 10965

Under license of E. MERCK, Darmstadt, Germany

 **Advantus™**  
 Pharmaceuticals

REV 3/93  
 L-32073-93

 LEDERLE LABORATORIES  
 A Division of American Cyanamid Company  
 Wayne, New Jersey 07470

Under license of E. MERCK, Darmstadt, Germany

Copromoted by  
 **Advantus™ Procter & Gamble**  
 PHARMACEUTICALS

# ATTENTION

## DIPLOMATES OF THE ABFP ADDRESS CHANGE FORM

***** 3-DIGIT-080 0028 <b>12345</b> 76/88 ABFP BEVERLY J. CLARK MD 24 LEXINGTON ST AKRON OH 44309
---

5-digit ABFP Identification Number

The Board prefers the use of *professional addresses*, because the address given will become your "address of record" with the Board and will be published in our Directory of Diplomates.

Current addresses for all Diplomates are necessary for communication from the Board relating to the Examinations, up-dated Recertification information, etc., as well as to ensure the receipt of *The Journal of the American Board of Family Practice*.

Name \_\_\_\_\_

### Current Address

### New Address

Street \_\_\_\_\_

Street \_\_\_\_\_

City/State \_\_\_\_\_

City/State \_\_\_\_\_

Zip Code \_\_\_\_\_

Zip Code \_\_\_\_\_

Effective Date of Change \_\_\_\_\_

Signature of Diplomate \_\_\_\_\_

ABFP Identification Number \_\_\_\_\_

(5-digit number above name on mailing label)

Year of Certification or Recertification \_\_\_\_\_

**Return to:**

**Ann Stockham  
The American Board of Family Practice  
2228 Young Drive  
Lexington, KY 40505**

# Release the grip of tension headache



**Non-scheduled\*** **Esgic**

Butalbital 50mg (Warning: May be habit forming)  
/Acetaminophen 500mg/Caffeine 40mg

**Over 50% more analgesic power than the leading products in its class.**

**Well tolerated — Without aspirin-related side effects such as GI irritation and GI bleeding.<sup>1-5</sup> The most frequent adverse reactions are drowsiness and dizziness.**

# Esgicplus™ tablets

Butalbital 50mg (Warning: May be habit forming)  
/Acetaminophen 500mg/Caffeine 40mg

References: 1. Benson GD. Hepatotoxicity following the therapeutic use of antipyretic analgesics. *Am J Med*. 1983;75(suppl 5A):85-93. 2. Jick H. Effects of aspirin and acetaminophen in gastro-intestinal hemorrhage. *Arch Intern Med*. 1981;141:316-321. 3. Mielke CH Jr. Comparative effects of aspirin and acetaminophen on hemostasis. *Arch Intern Med*. 1981;141:305-310. 4. Hansten PD. *Drug Interactions*. 5th ed. Philadelphia, PA: Lea & Febiger; 1985. p. 95. 5. Insel PA. Analgesic-antipyretics and antiinflammatory agents; drugs employed in the treatment of rheumatoid arthritis and gout. In: Gilman AG, Rall TW, Nies AS, Taylor P, eds. *The Pharmacological Basis of Therapeutics*. 8th ed. New York, NY: Pergamon Press; 1990:638-681.

## ESGIC-PLUS™ Tablets

(Butalbital, Acetaminophen and Caffeine Tablets, USP)

50mg/500mg/40mg

Brief Prescribing Information: (Please see package insert for full prescribing information) Each Esgic-plus™ Tablet contains: Butalbital, USP 50 mg. WARNING: May be habit forming. Acetaminophen, USP 500 mg, Caffeine, USP 40 mg. In addition each tablet contains the following inactive ingredients: microcrystalline cellulose, croscarmellose sodium, colloidal silicon dioxide and stearic acid.

**CONTRAINDICATIONS:** This product is contraindicated under the following conditions: • Hypersensitivity or intolerance to any component of this product. • Patients with porphyria. **WARNINGS:** Butalbital is habit-forming and potentially abusive. Consequently, the extended use of this product is not recommended.

**PRECAUTIONS: General:** Esgic-plus™ Tablets should be prescribed with caution in certain special-risk patients, such as the elderly or debilitated, and those with severe impairment of renal or hepatic function, or acute abdominal conditions. **Information for Patients:** This product may impair mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery. Such tasks should be avoided while taking this product. Alcohol and other CNS depressants may produce an additive CNS depression, when taken with this combination product, and should be avoided. Butalbital may be habit-forming. Patients should take the drug only for as long as it is prescribed, in the amounts prescribed, and no more frequently than prescribed. **Laboratory Tests:** In patients with severe hepatic or renal disease, effects of therapy should be monitored with serial liver and/or renal function tests. **Drug Interactions:** The CNS effects of butalbital may be enhanced by monoamine oxidase (MAO) inhibitors. Esgic-plus™ Tablets may enhance the effects of: other narcotic analgesics, alcohol, general anesthetics, tranquilizers such as chloridazepoxide, sedative-hypnotics, or other CNS depressants, causing increased CNS depression. **Drug/Laboratory Test Interactions:** Acetaminophen may produce false-positive test results for urinary 5-hydroxyindoleacetic acid. **Carcinogenesis, Mutagenesis, Impairment of Fertility:** No adequate studies have been conducted in animals to determine whether acetaminophen or butalbital have a potential for carcinogenesis, mutagenesis, or impairment of fertility. **Pregnancy: Teratogenic Effects:** Pregnancy Category C. Animal reproduction studies have not been conducted with this combination product. It is also not known whether Esgic-plus™ Tablets can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. This product should be given to a pregnant woman only when clearly needed. **Nonteratogenic Effects:** Withdrawal seizures were reported in a two-day-old male infant whose mother had taken a butalbital-containing drug during the last two months of pregnancy. Butalbital was found in the infant's serum. The infant was given phenobarbital 5 mg/kg, which was tapered without further seizure or other withdrawal symptoms. **Nursing Mothers:** Caffeine, barbiturates and acetaminophen are excreted in breast milk in small amounts, but the significance of their effects on nursing infants is not known. Because of potential for serious adverse reactions in nursing infants from Esgic-plus™ Tablets, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother. **Pediatric Use:** Safety and effectiveness in children below the age of 12 have not been established.

**ADVERSE REACTIONS: Frequently Observed:** The most frequently reported adverse reactions are drowsiness, light-headedness, dizziness, sedation, shortness of breath, nausea, vomiting, abdominal pain, and intoxicated feeling. **Infrequently Observed:** All adverse events tabulated below are classified as infrequent. **Central Nervous:** headache, shaky feeling, tingling, agitation, fainting, fatigue, heavy eyelids, high energy, hot spells, numbness, sluggishness, seizure. Mental confusion, excitement or depression can also occur due to intolerance, particularly in elderly or debilitated patients, or due to overdose of butalbital. **Autonomic Nervous:** dry mouth, hyperhidrosis. **Gastrointestinal:** difficulty swallowing, heartburn, flatulence, constipation. **Cardiovascular:** tachycardia. **Musculoskeletal:** leg pain, muscle fatigue. **Genitourinary:** diuresis. **Miscellaneous:** pruritus, fever, earache, nasal congestion, tinnitus, euphoria, allergic reactions. Several cases of dermatological reactions, including toxic epidermal necrolysis and erythema multiforme, have been reported. The following adverse drug events may be borne in mind as potential effects of the components of this product. Potential effects of high dosage are listed in the OVERDOSAGE section. **Acetaminophen:** allergic reactions, rash, thrombocytopenia, agranulocytosis. **Caffeine:** cardiac stimulation, irritability, tremor, dependence, nephrotoxicity, hyperglycemia.

**DRUG ABUSE AND DEPENDENCE: Abuse and Dependence:** Butalbital: Barbiturates may be habit-forming; Tolerance, psychological dependence, and physical dependence may occur especially following prolonged use of high doses of barbiturates. The average daily dose for the barbiturate addict is usually about 1500 mg. As tolerance to barbiturates develops, the amount needed to maintain the same level of intoxication increases; tolerance to a fatal dosage, however, does not increase more than two-fold. As this occurs, the margin between an intoxication dosage and fatal dosage becomes smaller. The lethal dose of a barbiturate is far less if alcohol is also ingested. Major withdrawal symptoms (convulsions and delirium) may occur within 16 hours and last up to 5 days after abrupt cessation of these drugs. Intensity of withdrawal symptoms gradually declines over a period of approximately 15 days. Treatment of barbiturate dependence consists of cautious and gradual withdrawal of the drug. Barbiturate-dependent patients can be withdrawn by using a number of different withdrawal regimens. One method involves initiating treatment at the patient's regular dosage level and gradually decreasing the daily dosage as tolerated by the patient.

**OVERDOSAGE:** Following an acute overdose of Esgic-plus™ Tablets, toxicity may result from the barbiturate or the acetaminophen. Toxicity due to caffeine is less likely, due to the relatively small amounts in this formulation. **Signs and Symptoms:** Toxicity from barbiturate poisoning includes drowsiness, confusion, and coma; respiratory depression; hypotension; and hypovolemic shock. In acetaminophen overdose: dose-dependent, potentially fatal hepatic necrosis is the most serious adverse effect. Renal tubular necroses, hypoglycemic coma and thrombocytopenia may also occur. Early symptoms following a potentially hepatotoxic overdose may include: nausea, vomiting, diaphoresis and general malaise. Clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours post-ingestion. In adults, hepatic toxicity has rarely been reported with acute overdoses of less than 10 grams, or fatalities with less than 15 grams. **Acute caffeine poisoning** may cause insomnia, restlessness, tremor, and delirium, tachycardia and extrasystoles. **Treatment:** A single or multiple overdose with this combination product is a potentially lethal polydrug overdose, and consultation with a regional poison control center is recommended. Immediate treatment includes support of cardiorespiratory function and measures to reduce drug absorption. Vomiting should be induced mechanically, or with syrup of ipecac, if the patient is alert (adequate pharyngeal and laryngeal reflexes). Oral activated charcoal (1 g/kg) should follow gastric emptying. The first dose should be accompanied by an appropriate cathartic. If repeated doses are used, the cathartic might be included with alternate doses as required. Hypotension is usually hypovolemic and should respond to fluids. Pressors should be avoided. A cuffed endotracheal tube should be inserted before gastric lavage of the unconscious patient and, when necessary, to provide assisted respiration. If renal function is normal, forced diuresis may aid in the elimination of the barbiturate. Alkalinization of the urine increases renal excretion of some barbiturates, especially phenobarbital. Meticulous attention should be given to maintaining adequate pulmonary ventilation. In severe cases of intoxication, peritoneal dialysis, or preferably hemodialysis may be considered. If hypoprothrombinemia occurs due to acetaminophen overdose, vitamin K should be administered intravenously. If the dose of acetaminophen may have exceeded 140 mg/kg, acetylcysteine should be administered as early as possible. Serum acetaminophen levels should be obtained, since levels four or more hours following ingestion help predict acetaminophen toxicity. Do not await acetaminophen assay results before initiating treatment. Hepatic enzymes should be obtained initially, and repeated at 24-hour intervals. Methemoglobinemia over 30% should be treated with methylene blue by slow intravenous administration.

**Toxic Doses (for adults):** Butalbital: toxic dose 1g (20 tablets); Acetaminophen: toxic dose 10g (20 tablets); Caffeine: toxic dose 1g (25 tablets). CAUTION: Federal law prohibits dispensing without prescription.

Manufactured by: MIKART, INC., Atlanta, GA 30318

Distributed by: FOREST PHARMACEUTICALS, INC., Subsidiary of Forest Laboratories, Inc., St. Louis, MO 63045

Revised 3/94

Code 374A00

FOREST PHARMACEUTICALS, INC.  
UAD LABORATORIES  
St. Louis, Missouri 63045

# Hate to Wait?



## Don't get aggravated.


Use your waiting time to read *Journal Watch*, the newsletter that monitors over 20 top journals. It provides summaries of the most significant advances so you're briefed on them in just minutes.

You can take *Journal Watch* anywhere. Read it on the go and make profitable use of otherwise wasted time.

## Get Updated.

Written by doctors for doctors, this 8-page newsletter published twice monthly brings you news of clinical advances while it's still fresh.

Order your introductory subscription today.



Try *JOURNAL WATCH* Today — Satisfaction Guaranteed

Credit card orders, call toll-free 1-800-843-6356  
FAX orders, 617-893-0413 or mail coupon below.

Yes, send me 24 issues of *Journal Watch*, mailed first class for just \$79. If not satisfied, I may cancel anytime for a full refund

NAME \_\_\_\_\_

ADDRESS \_\_\_\_\_

CITY \_\_\_\_\_

STATE \_\_\_\_\_ ZIP \_\_\_\_\_

Payment enclosed\*  Please send invoice

Charge my:  VISA  MasterCard  AmEx

CREDIT CARD NO. \_\_\_\_\_ CARD EXP. DATE \_\_\_\_\_

SIGNATURE \_\_\_\_\_

\*Make check payable to *Journal Watch*.

Mail to: *Journal Watch*, P.O. Box 9085, Waltham, MA 02254-9085.

Allow 4-6 weeks for delivery. ISSN: 0896-7210.

Send me more information on obtaining 50 CME credits.

SOJH1

# INFORMATION FOR AUTHORS

The *Journal of the American Board of Family Practice* welcomes for editorial review manuscripts that contribute to family practice as a clinical scientific discipline. High priority is given to reports of clinically relevant studies that have practical implications for improved patient care. Manuscripts are considered in relation to the extent to which they represent original work, their significance to the advancement of family medicine, and their interest to the practicing family physician. Some papers that are accepted by the *Journal* will be selected for an accompanying guest editorial or concurrent commentary by other invited authors addressing issues raised by the papers. The *Journal* publishes the following features:

**Original Articles.** Reports of original research, usually dealing with a clinical, health services, or other clinically relevant study.

**Medical Practice.** Scholarly articles that relate directly to clinical topics useful in everyday family practice, whether dealing with diagnostic or therapeutic roles of the family physician or reporting studies of what family physicians do in practice.

**Clinical Review.** In-depth reviews of specific clinical problems, disease entities, or treatment modalities; comprehensive and critical analysis of the literature is required (usual maximum length 5000 words).

**Clinical Guidelines and Primary Care.** Summaries of major clinical guidelines proposed by various specialty, governmental, or health care organizations, with critical commentary from a primary care perspective.

**Family Practice and the Health Care System.** Articles reporting studies and scholarly commentary on changing trends and patterns of care in family practice, primary care, and the health care system.

**Special Articles.** Articles in other areas that may relate to the role of the family physician, education for family practice, or other subjects important to family practice as a clinical specialty.

**Brief Reports.** Short reports of pilot studies or case reports with a teaching point of clinical relevance (usual length 1000–1500 words).

**Family Practice—World Perspective.** Papers reporting developments related to the practice or education of family physicians in various countries

around the world (usual length 1200–1800 words).

**Reflections in Family Practice.** Papers in narrative or essay format that illuminate qualitative aspects of family practice, including such areas as ethical issues, the physician-patient relationship, or the diverse roles of the family physician.

**Editorial.** Focused opinion or commentary that bears on an issue relevant to the field. May or may not accompany an original article in the same issue (usual length 1000–1500 words).

**Letters to the Editor.** Observations, opinion, or comment on topics under discussion in the *Journal*, usually not to exceed 500 words.

**Book Reviews.** Books for review and book reviews should be sent to Dr. John P. Geyman, Editor, the *Journal of the American Board of Family Practice*, Department of Family Medicine (HQ-30), School of Medicine, University of Washington, Seattle, WA 98195.

The following guidelines are in accordance with the "Uniform Requirements for Manuscripts Submitted to Biomedical Journals." The current (fourth) edition was published in the February 7, 1991, issue of the *New England Journal of Medicine*.

## MANUSCRIPT SUBMISSION

Manuscripts containing original material are accepted for consideration with the understanding that neither the article nor any part of its essential substance, tables, or figures has been or will be published or submitted for publication elsewhere before appearing in the *Journal*. This restriction does not apply to abstracts or press reports published in connection with scientific meetings. Copies of any possibly duplicative manuscripts should be submitted to the Editor along with the manuscript that is to be considered by the *Journal*. The *Journal* strongly discourages the submission of more than one article dealing with related aspects of the same study. In almost all cases, a single study is best reported in a single paper.

Submit an original and 3 copies of the complete manuscript, including text pages, legends, tables, references, and glossy prints of figures. Only typed copy, on standard-sized typewriter paper and double-spaced throughout, with margins of at least 2.5 cm, is

acceptable. Address all submissions to John P. Geyman, M.D., Editor, the *Journal of the American Board of Family Practice*, Department of Family Medicine (HQ-30), School of Medicine, University of Washington, Seattle, WA 98195. A covering letter should identify the person (with the address and telephone number) responsible for negotiations concerning the manuscript; the letter should make it clear that the final manuscript has been seen and approved by all authors. If authors acknowledge by name persons who provided important technical, advisory, or reviewer contributions, the corresponding author should sign the following statement: "I have obtained written permission from all persons named in the acknowledgment."

The *Journal* expects authors to take public responsibility for their manuscripts, including conception and design of the work, data analysis, writing, and review of the paper. Authors are expected to stand behind the validity of their data and, if asked by the Editor, to submit the actual data for editorial review with the manuscript. In most instances authorship should be limited to 8 authors or fewer, all meeting the above criteria for authorship. Exceptions to these guidelines, especially those involving multisite collaborative research projects, should be discussed on a case-by-case basis with the Editor.

The *Journal* also expects authors to disclose any commercial associations that might pose a conflict of interest in connection with the submitted article. Consultancies, stock ownership or other equity interests, patent-licensing arrangements, and other kinds of associations that might involve conflict of interest should be disclosed to the Editor in a covering letter at the time of submission. Such information will be held in confidence while the paper is under review and will not influence the editorial decision. If the manuscript is accepted, the Editor will discuss with the authors how best to disclose the relevant information. Questions about this policy should be directed to the Editor.

## MANUSCRIPTS

### Titles and Authors' Names

With the manuscript, provide a page giving the title of the paper; a running foot of fewer than 40 letter spaces; the name(s) of the author(s), including first name(s) and academic degree(s); the name of the department and institution in which the work was done; and the name and address of the author to whom reprint requests should be

addressed. All funding sources supporting the work should be routinely acknowledged on the title page, as should all institutional or corporate affiliations of the authors. Two to four key words should be submitted with the manuscripts to be used for purposes of classification by subject. Use terms from the Medical Subject Headings from *Index Medicus* when possible.

#### Abstracts

Use another page to provide an abstract of not more than 200 words. This abstract should be factual, not descriptive, with its content appropriate to the type of paper. For original articles reporting results of studies, a four-paragraph format should be used labeled Background, Methods, Results, and Conclusions. These should briefly describe, respectively, the object of the study, the methods used, the major results, and the author(s) conclusions. Abstracts are not necessary for Brief Reports or World Perspective papers.

#### Abbreviations

Except for units of measurement, abbreviations are discouraged. Consult the *Council of Biology Editors Style Manual* (Fifth edition. Bethesda, MD: Council of Biology Editors, 1983) for lists of standard abbreviations. The first time an abbreviation appears, it should be preceded by the words for which it stands.

#### Drug Names

Generic names should, in general, be used. If an author so desires, brand names may be inserted in parentheses.

#### Inclusive Language

Sex bias should be avoided and gender-inclusive language used whenever possible.

#### References

References must be typed in double spacing and numbered consecutively as they are cited. References first cited in tables or figure legends must be numbered so that they will be in sequence with references cited in the text. The style of references is that of the *Index Medicus*. List all authors when there are 6 or fewer; when there are 7 or more, list the first 6, then "et al." Sample references are as follows:

#### Standard Journal Article

(List all authors, but if the number exceeds 6, give 6 followed by et al. Note that month and issue number are omitted when a journal has continuous pagination throughout a volume.)

Morrow JD, Margolies GR, Rowland J, Roberts LJ 2nd. Evidence that histamine is the causative toxin of

scombroid-fish poisoning. *N Engl J Med* 1991; 324:716-20.

#### Organization as Author

Clinical Experience Network (CEN). A large-scale, office-based study evaluates the use of a new class of nonse dating antihistamines. A report from CEN. *J Am Board Fam Pract* 1990; 3:241-58.

#### Book

Rakel RE. Textbook of family practice. 4th ed. Philadelphia: WB Saunders, 1990.

#### Chapter in Book

Haynes RC Jr. Agents affecting calcification: calcium, parathyroid hormone, calcitonin, vitamin D, and other compounds. In: Gilman AG, Rall TW, Nies AS, Taylor P, editors. Goodman and Gilman's the pharmacological basis of therapeutics. 8th ed. New York: Pergamon Press, 1990.

#### Government Agency

Schwartz JL. Review and evaluation of smoking cessation methods: the United States and Canada, 1978-1985. Bethesda, MD: Department of Health and Human Services, 1987. (NIH publication no. 87-2940.)

#### Personal Communications

Numbered references to personal communications, unpublished data, and manuscripts either "in preparation" or "submitted for publication" are unacceptable (see "Permissions"). If essential, such material may be incorporated in the appropriate place in the text.

#### Tables

Type tables in double spacing on separate sheets, and provide a title for each. For footnotes, use the following symbols, in this sequence: \*, †, ‡, §, ¶, \*\*, ††, etc. Excessive tabular data are discouraged. If an article is accepted, the *Journal* will arrange to deposit extensive tables of important data with the National Auxiliary Publications Service (NAPS); we will pay for the deposit and add an appropriate footnote to the text. This service makes microfiche or photocopies of tables available at moderate charges to those who request them.

#### Illustrations

Figures should be professionally designed. Glossy, black-and-white photographs are requested. Symbols, lettering, and numbering should be clear, and these elements should be large enough to remain legible after the figure has been reduced to fit the width of a single column.

The back of each figure should include the sequence number, the name of the author, and the proper orientation (e.g., "top"). Do not mount the figure on cardboard. Photomicrographs should be cropped to a width of 8 cm, and electron photomicrographs should have internal scale markers.

If photographs of patients are used, either the subjects should not be identifiable or their pictures must be accompanied by written permission to use the figure. Permissions forms are available from the Editor.

Legends for illustrations should be type-written (double-spaced) on a separate sheet and should not appear on the illustrations.

Color illustrations are used from time to time. Send both transparencies and prints for this purpose.

#### Permissions

Every effort (short of changing the patient data) should be made by the authors to protect the anonymity of patients (and relatives) in any published work. If identification is unavoidable, informed consent should be obtained and attached with the submitted letter; in the case of minors or incompetent patients, consent should be obtained from relatives or guardians.

Materials taken from other sources must be accompanied by a written statement from both author and publisher giving permission to the *Journal* for reproduction. Obtain permission in writing from at least one author of papers still in press, of unpublished data, and of personal communications.

#### REVIEW AND ACTION

Manuscripts are examined by the editorial staff and are usually sent to outside reviewers. Authors will remain anonymous to outside reviewers and vice versa. External statistical review will be accomplished where appropriate. Every effort will be made to complete the review process as expeditiously as possible.

#### Copyright Transfer Forms

Transfer of copyright to the *Journal* is requested upon acceptance of the material for publication. Copyright transfer is required of all materials to be published in the *Journal*, including Letters to the Editor and Book Reviews.

#### Reprints

Authors will receive reprint information and rates when they are sent their galley proofs. Reprints ordered at that time will be shipped about 3 weeks after the publication date.

# INFORMATION FOR READERS

## The Journal of the American Board of Family Practice

Official Publication of the American Board of Family Practice  
2228 Young Drive, Lexington, KY 40505

### EDITORS

John P. Geyman, M.D., *Editor*  
Paul R. Young, M.D., *Executive Editor*  
Alfred O. Berg, M.D., *Associate Editor* Paul Brucker, M.D., *Associate Editor*  
G. Gayle Stephens, M.D., *Associate Editor*  
Claire Z. Fenwick, *Assistant Editor*  
Ann Stockham, *Copy Editor and Assistant Executive Editor*  
Virginia M. Gessner, *Senior Editorial Assistant*  
Debbie Wilson, *Production Assistant* Mary K. Lowell, *Reference Verification*

### PUBLISHING SERVICES

Publishing Division, Massachusetts Medical Society  
Robert D. Bovenschulte, *Vice President for Publishing*  
William H. Paige, *Executive Director for Operations*  
Christopher R. Lynch, *Executive Director for Product Management*  
Justin R. Spence, *Product Manager*  
Robert Quinn, *Reprints*

#### Advertising Sales and Production

Arthur Wilschek, *Director*  
William Healy, *Manager, Midwest Accounts*  
Lewis Wetzel, *Manager, Eastern Accounts*  
Wayne Wickman, *Manager, Eastern Accounts*  
Christine Miller, *Manager, Recruitment Advertising*  
Mary Kaye Howe, *Production Manager*  
Lynne Brochu

Circulation and Product Marketing  
Laurie Pass, *Director*

#### Electronic Production

Ruth E. Goodman, *Director*  
Tommie Richardson, *Coordinator*  
Sherie Peters, Thomas Gardon,  
Susan McDonough, Laurie Marcaurelle,  
David Lashua

#### Computer Assisted Publishing Support

Leon Barzin, *Director*  
Martha Soule

#### Customer Service and Fulfillment

Elizabeth C. Clancy, *Director*

#### Distribution and Postal Affairs

Mark Davidson, *Director*

#### Printing Services

Joseph Cortese, *Director*

### SUBSCRIPTION INFORMATION AND SERVICES

The *Journal of the American Board of Family Practice* is supplied free of charge to 50,400 Diplomates and Residents of the American Board of Family Practice. For information please contact:

American Board of Family Practice  
2228 Young Drive  
Lexington, KY 40505  
Tel: (606) 269-5626  
FAX: (606) 266-9699

For all other subscribers please contact:

**The Journal of the American Board of Family Practice**  
Subscription Department  
1440 Main Street  
Waltham, MA 02154-1649  
(617) 893-3800, ext. 1199  
Telex: 5106017779 NEJM BOS  
FAX: (617) 893-0413

### SUBSCRIPTION RATES

	Domestic	International*
Institutions	\$58.00	\$60.00
Physicians	\$35.00	\$45.00
Residents/Students	\$20.00	\$45.00

\*Pounds Sterling drawn on U.K. banks accepted and converted at current rate of exchange. U.S. dollars drawn on U.S. banks.

### OTHER SUBSCRIPTION INFORMATION

Diplomates should make address changes on the form accompanying this issue and forward to the Diplomat address listed above. All other subscribers should forward changes to the Waltham, Mass., address listed on this page. Changes must be received at least six weeks in advance of intended move. Please send new address, old address, and expected date of change.

### ISSUES NOT RECEIVED

Missing issues will be replaced for up to three months from the issue date without charge. Diplomates and other subscribers who fail to notify the Lexington, Ky., or the Waltham, Mass., office of address changes will not be eligible for free replacement issues. Claims beyond the three-month limit must be prepaid at the backcopy rates. Claims should be sent to either the Diplomat or regular subscriber address listed on this page.

### BACK COPIES

If you wish to purchase back copies (issues published prior to your effective start date) of the *Journal of the American Board of Family Practice*, there is a charge of \$12.50 per issue. Contact the Waltham, Mass., address listed above for information.

### REPRINTS

Individual copies of articles are available from the Waltham, Mass., office. If you wish to order bulk reprints (minimum order of 100) please contact the Reprint Department (617) 893-3800, ext. 1279, at the Waltham, Mass., office.


### COPYRIGHT

Material appearing in the *Journal of the American Board of Family Practice* is covered by copyright. Copying beyond the quantities permitted under "fair use" as defined by U.S. copyright law is allowed provided the stated fee of \$.20 per page is paid through the Copyright Clearance Center, 21 Congress St., Salem, MA 01970. This consent does not extend to other copying, such as copying for advertising or promotional purposes. Single copies for personal or internal use are allowed at no charge. Nonprofit institutions may make copies provided they obtain prior consent from the *Journal of the American Board of Family Practice*, Rights and Permissions Department, 1440 Main Street, Waltham, MA 02154-1649, (617) 893-3800, ext. 1413.

### INDEXING AND MICROFORM

The *Journal of the American Board of Family Practice* is indexed in *Index Medicus* and is available in microform from University Microfilms International.

# Who's a candidate?





**BRIEF SUMMARY**

**TABLETS**  
**PLENDIL®**  
 (FELODIPINE)  
 EXTENDED-RELEASE TABLETS

**INDICATIONS AND USAGE**

PLENDIL® is indicated for the treatment of hypertension. PLENDIL may be used alone or concomitantly with other antihypertensive agents.

**CONTRAINDICATIONS**

PLENDIL is contraindicated in patients who are hypersensitive to this product.

**PRECAUTIONS**

**General**

**Hypotension:** Felodipine, like other calcium antagonists, may occasionally precipitate significant hypotension and rarely syncope. It may lead to reflex tachycardia which in susceptible individuals may precipitate angina pectoris. (See ADVERSE REACTIONS.)

**Heart Failure:** Although acute hemodynamic studies in a small number of patients with NYHA Class II or III heart failure treated with felodipine have not demonstrated negative inotropic effects, safety in patients with heart failure has not been established. Caution therefore should be exercised when using PLENDIL in patients with heart failure or compromised ventricular function, particularly in combination with a beta blocker.

**Elderly Patients or Patients with Impaired Liver Function:** Patients over 65 years of age or patients with impaired liver function may have elevated plasma concentrations of felodipine and may therefore respond to lower doses of PLENDIL. These patients should have their blood pressure monitored closely during dosage adjustment of PLENDIL and should rarely require doses above 10 mg. (See CLINICAL PHARMACOLOGY and DOSAGE AND ADMINISTRATION sections of complete Prescribing Information.)

**Peripheral Edema:** Peripheral edema, generally mild and not associated with generalized fluid retention, was the most common adverse event in the clinical trials. The incidence of peripheral edema was both dose- and age-dependent. Frequency of peripheral edema ranged from about 10 percent in patients under 50 years of age taking 5 mg daily to about 30 percent in those over 60 years of age taking 20 mg daily. This adverse effect generally occurs within 2-3 weeks of the initiation of treatment.

**Information for Patients**

Patients should be instructed to take PLENDIL whole and not to crush or chew the tablets. They should be told that mild gingival hyperplasia (gum swelling) has been reported. Good dental hygiene decreases its incidence and severity.

**NOTE:** As with many other drugs, certain advice to patients being treated with PLENDIL is warranted. This information is intended to aid in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects.

**Drug Interactions**

**Beta-Blocking Agents:** A pharmacokinetic study of felodipine in conjunction with metoprolol demonstrated no significant effects on the pharmacokinetics of felodipine. The AUC and C<sub>max</sub> of metoprolol, however, were increased approximately 31 and 38 percent, respectively. In controlled clinical trials, however, beta blockers including metoprolol were concurrently administered with felodipine and were well tolerated.

**Cimetidine:** In healthy subjects pharmacokinetic studies showed an approximately 50 percent increase in the area under the plasma concentration time curve (AUC) as well as the C<sub>max</sub> of felodipine when given concomitantly with cimetidine. It is anticipated that a clinically significant interaction may occur in some hypertensive patients. Therefore, it is recommended that low doses of PLENDIL be used when given concomitantly with cimetidine.

**Digoxin:** When given concomitantly with felodipine the peak plasma concentration of digoxin was significantly increased. There was, however, no significant change in the AUC of digoxin.

**Anticonvulsants:** In a pharmacokinetic study, maximum plasma concentrations of felodipine were considerably lower in epileptic patients on long-term anticonvulsant therapy (e.g., phenytoin, carbamazepine, or phenobarbital) than in healthy volunteers. In such patients, the mean area under the felodipine plasma concentration-time curve was also reduced to approximately six percent of that observed in healthy volunteers. Since a clinically significant interaction may be anticipated, alternative antihypertensive therapy should be considered in these patients.

**Other Concomitant Therapy:** In healthy subjects there were no clinically significant interactions when felodipine was given concomitantly with indomethacin or spironolactone.

**Interaction with Food:** See CLINICAL PHARMACOLOGY, Pharmacokinetics and Metabolism section of complete Prescribing Information.

**Carcinogenesis, Mutagenesis, Impairment of Fertility**

In a two-year carcinogenicity study in rats fed felodipine at doses of 7.7, 23.1 or 69.3 mg/kg/day (up to 28 times' the maximum recommended human dose on a mg/m<sup>2</sup> basis), a dose related increase in the incidence of benign interstitial cell tumors of the testes (Leydig cell tumors) was observed in treated male rats. These tumors were not observed in a similar study in mice at doses up to 138.6 mg/kg/day (28 times' the maximum recommended human dose on a mg/m<sup>2</sup> basis). Felodipine, at the doses employed in the two-year rat study, has been shown to lower testicular testosterone and to produce a corresponding increase in serum luteinizing hormone in rats. The Leydig cell tumor development is possibly secondary to these hormonal effects which have not been observed in man.

In this same rat study a dose-related increase in the incidence of focal squamous cell hyperplasia compared to control was observed in the esophageal groove of male and female rats in all dose groups. No other drug-related esophageal or gastric pathology was observed in the rats or with chronic administration in mice and dogs. The latter

species, like man, has no anatomical structure comparable to the esophageal groove.

Felodipine was not carcinogenic when fed to mice at doses of up to 138.6 mg/kg/day (28 times' the maximum recommended human dose on a mg/m<sup>2</sup> basis) for periods of up to 80 weeks in males and 99 weeks in females.

Felodipine did not display any mutagenic activity *in vitro* in the Ames microbial mutagenicity test or in the mouse lymphoma forward mutation assay. No clastogenic potential was seen *in vivo* in the mouse micronucleus test at oral doses up to 2500 mg/kg (506 times' the maximum recommended human dose on a mg/m<sup>2</sup> basis) or *in vitro* in a human lymphocyte chromosome aberration assay.

A fertility study in which male and female rats were administered doses of 3.8, 9.6 or 26.9 mg/kg/day showed no significant effect of felodipine on reproductive performance.

**Pregnancy**

**Pregnancy Category C**

**Teratogenic Effects:** Studies in pregnant rabbits administered doses of 0.46, 1.2, 2.3 and 4.6 mg/kg/day (from 0.4 to 4 times' the maximum recommended human dose on a mg/m<sup>2</sup> basis) showed digital anomalies consisting of reduction in size and degree of ossification of the terminal phalanges in the fetuses. The frequency and severity of the changes appeared dose-related and were noted even at the lowest dose. These changes have been shown to occur with other members of the dihydropyridine class and are possibly a result of compromised uterine blood flow. Similar fetal anomalies were not observed in rats given felodipine.

In a teratology study in cynomolgus monkeys no reduction in the size of the terminal phalanges was observed but an abnormal position of the distal phalanges was noted in about 40 percent of the fetuses.

**Nonteratogenic Effects:** A prolongation of parturition with difficult labor and an increased frequency of fetal and early postnatal deaths were observed in rats administered doses of 9.6 mg/kg/day (4 times' the maximum human dose on a mg/m<sup>2</sup> basis) and above.

Significant enlargement of the mammary glands in excess of the normal enlargement for pregnant rabbits was found with doses greater than or equal to 1.2 mg/kg/day (equal to the maximum human dose on a mg/m<sup>2</sup> basis). This effect occurred only in pregnant rabbits and regressed during lactation. Similar changes in the mammary glands were not observed in rats or monkeys.

There are no adequate and well-controlled studies in pregnant women. If felodipine is used during pregnancy, or if the patient becomes pregnant while taking this drug, she should be apprised of the potential hazard to the fetus, possible digital anomalies of the infant, and the potential effects of felodipine on labor and delivery, and on the mammary glands of pregnant females.

**Nursing Mothers**

It is not known whether this drug is secreted in human milk and because of the potential for serious adverse reactions from felodipine in the infant, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

**Pediatric Use**

Safety and effectiveness in children have not been established.

**ADVERSE REACTIONS**

In controlled studies in the United States and overseas approximately 3000 patients were treated with felodipine as either the extended-release or the immediate-release formulation.

The most common clinical adverse experiences reported with PLENDIL® (felodipine) administered as monotherapy in all settings and with all dosage forms of felodipine were peripheral edema and headache. Peripheral edema was generally mild, but it was age- and dose-related and resulted in discontinuation of therapy in about 4 percent of the enrolled patients. Discontinuation of therapy due to any clinical adverse experience occurred in about 9 percent of the patients receiving PLENDIL, principally for peripheral edema, headache, or flushing.

Adverse experiences that occurred with an incidence of 1.5 percent or greater during monotherapy with PLENDIL without regard to causality are compared to placebo in the table below.

**Percent of Patients with Adverse Effects in Controlled Trials of PLENDIL as Monotherapy (incidence of discontinuations shown in parentheses)**

Adverse Effect	PLENDIL® N = 730	Placebo % N = 283
Peripheral Edema	22.3 (4.2)	3.5
Headache	18.6 (2.1)	10.6
Flushing	6.4 (1.0)	1.1
Dizziness	5.8 (0.8)	3.2
Upper Respiratory Infection	5.5 (0.1)	1.1
Asthenia	4.7 (0.1)	2.8
Cough	2.9 (0.0)	0.4
Paresthesia	2.5 (0.1)	1.8
Dyspepsia	2.3 (0.0)	1.4
Chest Pain	2.1 (0.1)	1.4
Nausea	1.9 (0.8)	1.1
Muscle Cramps	1.9 (0.0)	1.1
Palpitation	1.8 (0.5)	2.5
Abdominal Pain	1.8 (0.3)	1.1
Constipation	1.6 (0.1)	1.1
Diarrhea	1.6 (0.1)	1.1
Pharyngitis	1.6 (0.0)	0.4
Rhinorrhea	1.6 (0.0)	0.0
Back Pain	1.6 (0.0)	1.1
Rash	1.5 (0.1)	1.1

In the two dose response studies using PLENDIL as monotherapy, the following table describes the incidence (percent) of adverse experiences that were dose-related. The incidence of discontinuations due to these adverse experiences are shown in parentheses.

The incidence of discontinuations due to these adverse experiences are shown in parentheses.

Adverse Effect	Placebo N = 121	2.5 mg N = 71	5.0 mg N = 72	10.0 mg N = 123	20 mg N = 50
Peripheral Edema	2.5 (1.6)	1.4 (0.0)	13.9 (2.8)	19.5 (2.4)	36.0 (10.0)
Palpitation	0.8 (0.8)	1.4 (0.0)	0.0 (0.0)	2.4 (0.8)	12.0 (8.0)
Headache	12.4 (0.0)	11.3 (1.4)	11.1 (0.0)	18.7 (4.1)	28.0 (18.0)
Flushing	0.0 (0.0)	4.2 (0.0)	2.8 (0.0)	8.1 (0.8)	20.0 (8.0)

In addition, adverse experiences that occurred in 0.5 up to 1.5 percent of patients who received PLENDIL® (felodipine) in all controlled clinical studies (listed in order of decreasing severity within each category) and serious adverse events that occurred at a lower rate and were found during marketing experience (those lower rate events are in italics) were: *Body as a Whole:* Facial edema, warm sensation; *Cardiovascular:* Tachycardia, myocardial infarction, hypotension, syncope, angina pectoris, arrhythmia; *Digestive:* Vomiting, dry mouth, flatulence; *Hematologic:* Anemia; *Musculoskeletal:* Arthralgia, arm pain, knee pain, leg pain, foot pain, hip pain, myalgia; *Nervous/Psychiatric:* Depression, anxiety disorders, insomnia, irritability, nervousness, somnolence; *Respiratory:* Bronchitis, influenza, sinusitis, dyspnea, epistaxis, respiratory infection, sneezing; *Skin:* Contusion, erythema, urticaria; *Urogenital:* Decreased libido, impotence, urinary frequency, urinary urgency, dysuria.

Felodipine, as an immediate release formulation, has also been studied as monotherapy in 680 patients with hypertension in U.S. and overseas controlled clinical studies. Other adverse experiences not listed above and with an incidence of 0.5 percent or greater include: *Body as a Whole:* Fatigue; *Digestive:* Gastrointestinal pain; *Musculoskeletal:* Arthritis, local weakness, neck pain, shoulder pain, ankle pain; *Nervous/Psychiatric:* Tremor; *Respiratory:* Rhinitis; *Skin:* Hyperhidrosis, pruritus; *Special Senses:* Blurred vision, tinnitus; *Urogenital:* Nocturia.

**Gingival Hyperplasia:** Gingival hyperplasia, usually mild, occurred in <0.5 percent of patients in controlled studies. This condition may be avoided or may regress with improved dental hygiene. (See PRECAUTIONS, Information for Patients.)

**Clinical Laboratory Test Findings**

**Serum Electrolytes:** No significant effects on serum electrolytes were observed during short- and long-term therapy.

**Serum Glucose:** No significant effects on fasting serum glucose were observed in patients treated with PLENDIL in the U.S. controlled study.

**Liver Enzymes:** One of two episodes of elevated serum transaminases decreased once drug was discontinued in clinical studies; no follow-up was available for the other patient.

**OVERDOSAGE**

Oral doses of 240 mg/kg and 264 mg/kg in male and female mice, respectively and 2390 mg/kg and 2250 mg/kg in male and female rats, respectively, caused significant lethality.

In a suicide attempt, one patient took 150 mg felodipine together with 15 tablets each of atenolol and spironolactone and 20 tablets of nitrazepam. The patient's blood pressure and heart rate were normal on admission to hospital; he subsequently recovered without significant sequelae.

Overdosage might be expected to cause excessive peripheral vasodilation with marked hypotension and possibly bradycardia.

If severe hypotension occurs, symptomatic treatment should be instituted. The patient should be placed supine with the legs elevated. The administration of intravenous fluids may be useful to treat hypotension due to overdosage with calcium antagonists. In case of accompanying bradycardia, atropine (0.5-1 mg) should be administered intravenously. Sympathomimetic drugs may also be given if the physician feels they are warranted.

It has not been established whether felodipine can be removed from the circulation by hemodialysis.

**DOSAGE AND ADMINISTRATION**

The recommended initial dose is 5 mg once a day. Therapy should be adjusted individually according to patient response, generally at intervals of not less than two weeks. The usual dosage range is 5-10 mg once daily. The maximum recommended daily dose is 20 mg once a day. That dose in clinical trials showed an increased blood pressure response but a large increase in the rate of peripheral edema and other vasodilatory adverse events (see ADVERSE REACTIONS). Modification of the recommended dosage is usually not required in patients with renal impairment.

PLENDIL should be swallowed whole and not crushed or chewed.

**Use in the Elderly or Patients with Impaired Liver Function:** Patients over 65 years of age or patients with impaired liver function, because they may develop higher plasma concentrations of felodipine, should have their blood pressure monitored closely during dosage adjustment (see PRECAUTIONS). In general, doses above 10 mg should not be considered in these patients.



For more detailed information, consult your Astra/Merck Specialist or see complete Prescribing Information.  
 Astra/Merck Group of Merck & Co., Inc.  
 725 Chesterbrook Boulevard, Wayne, PA 19087

PLENDIL is a product of Astra/Merck Research  
 Copyright ©1994 by Astra/Merck Group of Merck & Co., Inc.  
 All rights reserved  
 7650203  
 3000-0492/93/12-09/Ppl

\*Registered trademark of AB Astra  
 †Based on patient weight of 50 kg